Lynemouth, Northumberland (ironically less than ten miles from IRD), by as much as £1 million.

MOON FLIGHTS

Vinogradov and Apollo

from a Correspondent

ALTHOUGH Russian space research seems to have been tied, from the beginning, to the sequence of projects envisaged by Tsiolkovskii, which gives the creation of an orbital station priority over any voyages beyond the Earth's orbit, there is no doubt that the Apollo projects have evoked considerable scientific and popular interest throughout the Soviet Union. In a recent interview (*Pravda*, November 27) Academician A. P. Vinogradov, vice-president of the Academy of Sciences of the USSR, gives what would be called, in military terms, an "appreciation" of the Apollo results to date.

After outlining the surface experiments carried out by the landing parties of Apollo 11 and 12, he emphasizes that neither crew included a geologist, and hence that the rock samples collected would be entirely random, although the longer spell of the Apollo 12 crew outside the rocket would result in samples being "more diverse and richer". The chemical and geological findings of the Apollo 11 specimens are outlined, with especial note of their similarity to achondritic meteorites. Cautiously, Vinogradov states: "I do not therefore consider it to be an absurd idea that the basalt meteorites found on Earth are of lunar origin". He stresses, however, that for a proper appraisal, it would be necessary to have selenological specimens from different parts of the lunar surface, in particular from the highlands.

The problem is raised of the greater dustiness of the Ocean of Storms in comparison with the Sea of Tranquillity; this, says Vinogradov, is a "riddle", and he contents himself with restating the basic problem as "Is the presence of lunar dust connected with the nature of the surface or do any other factors play a part here?"

Finally, when asked to comment on the contribution of manned flights to lunar studies, he chose as a particularly noteworthy example the fact that the Moon is now supposed to have a definite crust and not to be the homogeneous body previously imagined. Rather strangely, no indication is given as to how this result was inferred. From Vinogradov's article it might be assumed that this new model of the lunar structure had been obtained either from the analysis of specimens or from the readings under natural conditions of the seismometer, magnetometer and spectrometer which he enumerates at the beginning of his appreciation. In extending the congratulations of Soviet scientists to the American "cosmonauts, scientists and engineers", he makes no mention whatever of the Intrepid crashdown nor of its world-shaking (or moon-shaking) consequences.

AGRICULTURE

Fast Growth at Nottingham

from our Botany Correspondent

THE University of Nottingham seems to be defying those pessimists who hold that the fruits of university

agricultural research are inferior to those that come out of special institutions. Nottingham has a fast expanding school of agriculture and financial backing to prove that many people think that its activities are very valuable. The latest annual report of the school of agriculture (price 10s.) records that the 1968-69 session, which marked the fiftieth anniversary of the teaching of agriculture on the site at Sutton Bonington, has seen the virtual completion of the new teaching and research block. This will house the departments of agricultural economics, agriculture and horticulture and the animal physiology group. There has also been a new building for avian physiology, and animal accommodation which has been built jointly with the new medical school.

Perhaps it is cooperation and diversification that are the secret of Nottingham's success in agriculture, for there is a generous admixture of pure and applied work in progress with no isolation from the other biological departments of the university. The spirit of diversity represented by the constituent departments of agricultural economics, agriculture and horticulture, applied biochemistry and nutrition and physiology and environmental studies is carried over into the undergraduate courses. The honours BSc degree depends on a "triad" system in which a student takes three honours units, selected to form an effective combination, and writes a dissertation in one of them. The wide scope of this system is partly a consequence of cooperation between faculties, so that topics such as embryology and entomology can be studied in the department of zoology. On the other hand, students of zoology can go to Sutton Bonington to learn about genetics, while lawyers and social scientists are instructed in agricultural economics.

Signs of approval of the research effort are recorded in the form of grants from government, industry and other organizations, which have recently been supplementing the contribution from the university's own funds. The department of applied biochemistry and nutrition, for example, has received £55,300 from diverse sources, most of it to cover one year's activities. The department of agricultural economics has been spending £134,000, which it is receiving in several instalments. This sum includes £30,000 over three years for joint work with the University of Zambia, involving an examination of factors affecting labour productivity. The rest of the school of agriculture has fared equally well, and no doubt feels confident in expanding its activities.

WATER RESOURCES

Icebergs for Sale

from our Geomagnetism Correspondent

THE idea of towing icebergs from Antarctica to irrigate deserts farther north sounds at first as if it might have been invented by Spike Milligan. But with the shortage of fresh water in many areas, a serious evaluation of the idea is being carried out by Dr William Campbell of the US Geological Survey and Dr Wilford Weeks of the US Army's Cold Regions Research and Engineering Laboratory. More than 80 per cent of the world's fresh water, equivalent to about sixty years' precipitation over the entire globe, is tied up in Arctic and Antarctic glaciers. Even in North America the volume of water stored as snow and ice exceeds that

of all lakes, rivers and reservoirs on the continent. Natural ice is a vast resource as yet barely tapped.

The economic use of ice meltwater is not new either in theory or practice. The earliest known large scale example was the commercial shipping of Alaskan lake ice to San Francisco during the 1850s with its replacement by glacier ice when lake ice was in short supply. Since then even larger scale use of glaciers for water has been suggested many times but has never (except perhaps in the Soviet Union) received a critical analysis.

The first problem is to find a supply of icebergs of the correct shape. In practice this means icebergs of roughly square or circular cross-section to avoid the hazards of rolling during towing. The sources of such "tabular" icebergs in the Arctic are not common, but Antarctica is more plentifully endowed especially in the large ice shelves which fringe the continent. The most obvious sites are the Ross Ice Shelf, which would be particularly suitable for supplying the west coast of South America, and the Amery Ice Shelf, conveniently located to satisfy Australian demands. A further possibility is the Filchner Ice Shelf which could supply icebergs to the Namib Desert area on the west coast of Africa. The selection of icebergs from these regions will be aided during the next few years by photography from polar-orbiting satellites capable of identifying individual objects with dimensions of about one hundred metres.

But would any ice arrive at its destination, or would it have melted away? For icebergs from the Antarctic, surface water temperatures at the northern ends of their journeys will be above 20° C; and for more than half of the transit distances temperatures will be above 5° C. Campbell and Weeks have examined three journeys ranging from 3,240 to 4,680 miles which, assuming an average towing speed of one knot, would last between 120 and 160 days, and conclude that large amounts of ice would remain. For example, an iceberg with lateral dimensions of 1,000 metres and 250 metres thick making a journey from Amery to Australia would end up with lateral dimensions of 760 metres and a thickness of 130 metres, or 30 per cent of the original. Longer journeys are less favourable; but 14 per cent of a Ross iceberg would reach the Atacama Desert.

And what of the economics of the operation? The only cost involved in the transit is that of the power required for towing. The resistance of an iceberg being pulled through water varies as the square of the velocity so that lower speeds are the most economic in terms of power. But longer towing times mean more melting so that a compromise must be made between melting losses and towing costs. Expenses could perhaps be reduced by slow towing along suitable ocean current paths.

Preliminary calculations show that an Amery iceberg of dimensions 2,700 × 2,700 × 250 metres towed at half a knot would reach Australia as one of 2,460 × 2,460 × 130 metres. The 207 billion gallons of water involved would be worth about \$5.5 million, or about 10 per cent of the cost of the same quantity of desalinated water, compared with towing costs of about \$1 million. A similar iceberg from the Ross Ice Shelf would arrive at the Atacama Desert as 101 billion gallons worth \$2.7 million with transport costs of \$1.3 million. None of these figures includes the costs involved in iceberg

selection, "docking", melting or the transport of the resulting water; but the economics seem favourable enough to warrant more detailed study even with water at its present price.

MATHEMATICS

Jamboree for Number Crunchers

THE academic session 1970-71 is to be Numerical Analysis Year. This name has been announced by Professor D. S. Jones of the University of Dundee, where the Department of Mathematics has just been awarded a grant of £23,000 by the Science Research Council. The money will be spent on a year-long round of conferences, symposia and summer schools, and Professor Jones hopes that at the end, numerical analysis in Britain will be stronger than before. British numerical analysts, Professor Jones believes, are lagging behind their opposite numbers in Europe, the United States and the Soviet Union. The trouble is that news of time-saving developments in numerical analysis is not getting through to the practising mathematicians working in computer departments. During Numerical Analysis Year, mathematicians from abroad will visit Dundee to spread the word and, Professor Jones hopes, generate interest and awareness among the steady flow of people from British universities, colleges, and industry who will visit the department. So far, thirty missionaries from overseas have agreed to come for all or part of the year, and at some point they should all be in Dundee together. Professor A. R. Mitchell, also of Dundee, is in charge of the organization, and a by-product of the concentration of talent could be the encouragement of more universities to do work in this important field.

FISHERIES

Scientific Farming

THE basic biology which must underpin any attempt to rationalize the fishing industry is providing the Ministry of Agriculture's marine scientists with a wide range of problems to solve. In his annual report for 1968, which has just appeared (obtainable free of charge on application to the Fisheries Laboratory, Lowestoft), the Director of Fishery Research, Dr H. A. Cole, depicts a scene of healthy progress, both in the physiology, genetics and ecology of sea fish—studies which form the foundation of fish farming projects such as those of the White Fish Authority—and in more peripheral investigations of marine pollution and the disposal of radioactive wastes into the sea. The research is supported by records and predictions of the fishing conditions in the areas exploited by British fishing fleets, including a computer model that analyses productivity in the North Atlantic since 1946.

Commercial considerations have obviously determined many of the research topics. Pigment abnormalities, for example, can make a fish on the shop counter look unattractive to customers, and if plaice are reared in a tank they are often deficient in colour. This deficiency has been correlated with the size of the plaice larvae. Pigment peculiarities in sole are rarer, but have been similarly associated with size. Attempts to cross-breed different kinds of fish aim to generate new characteristics of commercial value,