Gauss normal epoch, two further anomalies, designated *W* and *X*, frequently occur either as a pair close together or as a single anomaly. These have never been identified with any event or events on the continental scale. But if the rate of spreading between 0.9 and 3.32 million years is assumed constant, *W* turns out to have an average age of 1.99 million years, very close to the measured age of the Olduvai (1.96 million years). Anomaly *X*, with an average age of 2.3 million years, still remains unexplained but is less important in the sense that whenever only one anomaly appears in the region it is always *W*.

A further objection to the variable rate spreading interpretation is that the Gilsa event is left without a corresponding ocean magnetic anomaly. The important result, however, is that with the removal of the larger spreading rate discontinuity at 2 million years, the rate of spreading over the whole of the past 3.32 million years can be regarded as constant within experimental error—an altogether simpler solution which has the merit of explaining more of the observations than does variable rate spreading.

COMETS

OAQ views Tago-Sato-Kosaka

by our Astronomy Correspondent

INTERESTING data continue to flow down the telemetry channels from the Orbiting Astronomical Observatory (OAQ 2) which NASA launched in December 1968. By a happy chance, the brightest comet for several years has now appeared in the sky, and the OAQ is being used to secure the first observations of a comet in the ultraviolet. This follows surprising results obtained by the OAQ last year and released by the Americans in December (*Nature*, **224**, 1058; 1969) which seem to show that a few galaxies are unexpectedly bright in the ultraviolet.

Ultraviolet light below about 3000 Å is, of course, absorbed in the atmosphere, which among other things makes the important Lyman α line of hydrogen (1216 Å) inaccessible from the ground. With orbiting ultraviolet telescopes now available in the OAQ, one of the results should be a better knowledge of the distribution of hydrogen in space.

In this spirit NASA announced last week that Lyman α observations of Tago-Sato-Kosaka by the University of Wisconsin instruments on OAQ 2 show that the comet is surrounded by a cloud of hydrogen gas which is said to be comparable in size to the Sun. The hydrogen is presumably being ejected from the comet by solar heating, and will be interpreted in terms of Whipple's "icy-conglomerate" theory. This has it that the nucleus of a comet is made up of mineral particles bound up in a matrix of frozen gases such as water and ammonia—a flying gravel bank, in other words. Some of the gases vaporize when the comet approaches the Sun, to form a cloud known as the coma, but the size of the coma is small compared with the cloud of hydrogen which is now reported.

Following the usual practice, Tago-Sato-Kosaka is named after the three Japanese amateurs who independently discovered it on October 10 last year. (The Japanese have of course become dab hands at spotting undiscovered comets.) By December the comet was visible by eye in the southern hemisphere, moving

north, but it is now retreating from the Sun and unlikely to be visible by eye into February. On February 2 it will be in Aries, near Alpha Arietis, at about magnitude five—the naked eye limit.

The Wisconsin instruments acquired Tago-Sato-Kosaka on January 14, and the plan is to follow the hydrogen cloud for as long as possible. The chief experimenter is Dr Arthur D. Code, and the schedule seems to be to observe the comet for 10 per cent of each 100 minute orbit.

CELL DIVISION

Retine for Inhibition

from our Cytogenetics Correspondent

THE suggestion that dehydroascorbic acid is an inhibitor of cell division is a recent contribution to the discussion of how this process is controlled. Cells of a living organism may be actively dividing or they may be quiescent. How is the transition from one state to the other brought about? And how do cancer cells differ in their regulation of cell division? The task of answering these questions is formidable. One line of attack has been opened up by Szent-Györgyi and his colleagues. They argue that two regulators of division are involved; there is an inhibitor, "retine", and a promoter, "promine" (*Science*, **149**, 34; 1965). Szent-Györgyi's group identified retine as a keto-aldehyde, the simplest member of which is methylglyoxal. Keto-aldehydes are good electron acceptors and inhibit cell division and their effect can be reversed by thiols (molecules with —SH groups) which are good electron donors and so behave as promines (Együd, *Currents Mod. Biol.*, **1**, 14; 1967 and **2**, 128; 1968). These properties led Szent-Györgyi to propose (*Science*, **161**, 988; 1968) that the decision as to whether a cell would divide or not was controlled by the relative balance of retine and promine molecules. Stimuli to cell division, such as wounding of a tissue, cause the removal of retine molecules. In cancer cells it may be that retine is not produced and so the biochemistry of the cell favours cell division.

Much work remains to be done to support such concepts. Now, J. A. Edgar joins the fray with more ideas (*Experientia*, **25**, 1514; 1969). He accepts Szent-Györgyi's scheme but suggests that dehydroascorbic acid (DHA) may be a retine substance. DHA is widely found in animal and plant cells; it is an electron acceptor and is reversibly associated with ascorbic acid (AA). The reaction $DHA \rightleftharpoons AA$ is mediated by glutathione (GSH), an electron donor that for many years has been thought to be implicated with cell division, in the following way

Edgar summarizes a large body of evidence which indicates that the balance between GSH and DHA may control the ability of cells to divide. For example, in plants it has been found that in cells leaving the dormant state (as in seed germination) the amount of DHA decreases while the amount of AA and GSH