nucleotide bases and nicotinamide.

Professor M. Eigen (Göttingen) then advanced his recently developed theory of evolution during the phase of self-organization after chemical evolution had occurred and before Darwinian evolution began. This physical theory of evolution, starting at the molecular level, stresses the close linkage between the process of the recognition of the information properties of nucleic acids and the functional properties of proteins—information acquires a meaning only when coding for function, so that feedback processes occur between both information and function. Thus self-duplicating systems could not have been random but must have been directed.

Eigen has developed an equation which has the properties of selection, for it includes a value factor which would decide which molecular species would be selected to grow and which to decay. The equation is interesting because it has a vectorial function which would select not just for a single molecular species but for many, so resulting in more and more coupling. But there is also a moving threshold value, so that more and more species would be below the threshold value and thus decay, which means that the autocatalytic properties of the system would take care of selection. The equation also recognizes that the system cannot be at equilibrium because the growth properties would then disappear; however, the steady state of the total system must be constant with "fluxes" and "forces" to maintain the state. Briefly, the theory includes the ability of competitive growth and selection, sharp selection, evolutionary development from the initially established cycle (the universal code) and selection against "parasites".

Professor I. Prigogine (Brussels) talked on instabilities and structure and explained the phenomena of natural boundaries using models from hydrodynamics and the Jabotinski reaction (oxidation of malonic acid in the presence of cerium ions). Instabilities are created at boundaries resulting in "dissipative" chemical structures which can be stable for some time before abruptly disappearing. Professor Prigogine developed the idea that the dissipative structures occur at conditions which are far from equilibrium, thus favouring autocatalysis.

Dr S. Spiegelman (Columbia, NY) emphasized that nucleic acids are the only molecules which can duplicate themselves and have the possibility of transmitting errors, these being the two prime requirements for directed evolution of living systems. He discussed his work on Q β replicase and the search for variants with low molecular weights. Variants had been obtained which had discarded 85 per cent of their molecule leaving only a strand of about 550 nucleotides. These variants could adapt

to certain changes in environmental conditions by altering the allostery of their enzymes. Recently, mini-variants containing only 180 nucleotides have been obtained which can now be made to add about fifty nucleotides to themselves.

TUMOUR VIRUSES

A Transforming Gene

from our Cell Biology Correspondent

A LONG expectant audience has received its reward; in the latest issue of the *Proceedings of the US National Academy of Sciences* (67, 1775; 1970), Dulbecco and Eckhart report some of the properties of *ts3*, a mutant of polyoma virus isolated by M. Vogt, with a temperature sensitive lesion in one of the transforming genes of this virus. Polyoma virus and the other small DNA tumour virus, Simian Virus 40, can infect certain cells and cause them to be stably transformed so that their phenotype resembles in many respects that of tumour cells. These two viruses contain only 3×10^6 daltons of DNA sufficient to code for about 200,000 daltons of polypeptide or six to eight proteins.

On the face of things it should be straightforward to define such a simple genome, including those genes responsible for the establishment and maintenance of transformation, by isolating temperature sensitive mutants. An understanding of the molecular mechanisms of

virus transformation should also throw light on the essential differences between normal and cancerous cells. In 1965 Fried isolated the *tsa* mutant of polyoma virus which carries a temperature sensitive lesion in a gene required for the establishment but not the maintenance of transformation. But we have had to wait five years for the characterization of a mutation in a polyoma gene which controls the maintenance of transformation.

Hamster cells of the BHK line transformed by *ts3* and maintained at the permissive temperature, 32° C, have all the characteristics of BHK cells transformed by wild type polyoma virus; unlike untransformed cells, *ts3*-BHK cells grow in agar, are not subject to topoinhibition, that is, the inhibition of cell cellular DNA synthesis by such factors as the proximity of other cells, and their serum requirement differs from that of untransformed BHK cells. When *ts3*-BHK cells are shifted from 32° C to 39° C, however, although the cells' ability to grow in agar and their serum requirements are unchanged they become sensitive once again to topoinhibition. BHK cells transformed by wild type virus, by contrast, are insensitive to topoinhibition at both temperatures. This indicates that the loss of topoinhibition, which characterizes transformation by polyoma virus, depends on the continued production of a functional polyoma-gene product.

Optical Frequency Breakdown in Gases

ONE of the most surprising effects produced by powerful laser light is that it can transform gases which are normally transparent to the long wavelength red and infrared radiation of lasers into opaque, intensely bright and highly conducting plasmas in times ~ 1 ns. If the laser light is very powerful, say ~ 10 GW obtainable from multistage oscillator amplifier systems, a series of small discrete plasma regions separated by centimetres extending over several tens of metres along the unfocused beam can readily be produced in air at atmospheric pressure. Focused beams from much less powerful single stage lasers, giving intensities around 10^{11} W cm $^{-2}$, can also create multiple collinear plasmas, but on a very much smaller scale extending over millimetres; the newly discovered high pressure CO $_2$ lasers, the so called TEA (Transverse Excitation, Atmospheric) devices, operating in the infrared, can cause optical frequency breakdown of gases at smaller intensities.

There has been much discussion about the mechanisms responsible for discharge initiation and the processes which lead to the subsequent growth of ionization and the nature of the complex plasmas formed. Some recent research into the phenomenon is reported by W.

Zernik in next Monday's *Nature Physical Science* (229, 46; 1971). The laser flash itself evidently releases the first electron from a gas atom even though the value of the quantum of energy associated with the light is typically, for a ruby laser, only 10 per cent of the ionization potential. This seems to imply a multiphoton absorption mechanism which is considered by many as a satisfactory explanation of initiation (H. B. Bebb and A. Gold, *Phys. Rev.*, 143, 1; 1966), but accepted only with reserve by others (R. G. Tomlinson, *Phys. Rev. Lett.*, 14, 489; 1965).

Another surprising feature is the success with which an extrapolation of the classical microwave breakdown theory of gases to optical frequencies can account for many of the observed phenomena. The extrapolation is based on the view that ionization growth and plasma formation occur, following initiation, by inelastic collisions between atoms and electrons in which the interaction energy is drawn from the electromagnetic field of the laser light. This at once leads to the concept of an effective electric field, oscillating with the optical frequency, which produces the same energy transfer between electrons and atoms as a static field to which it is related by the light

frequency ω and electron-atom collision frequency. By this means excitation, ionization and other transport properties, readily measurable under static fields, may be used to deduce optical frequency counterparts. In this way it is possible to derive expressions for the critical radiation intensity required to cause breakdown in gases exposed to laser light; it turns out to be approximately proportional to ω^2/τ where τ is the duration of the laser flash (F. Morgan, L. R. Evans and C. G. Morgan, *J. Phys. D.*, **4**, in the press).

Comparison between measurement and prediction is remarkably good when the experimental complexities and theoretical approximations are taken into account, but it has been argued that because the classical oscillation energy of the electron in the laser field is small compared to the photon energy the classical description is inapplicable.

Zernik contends that this reasoning is incorrect and he applies a simple extrapolation of the classical theory in the case of breakdown in helium with some success over a wide pressure range.

The *ts3* mutation also affects the response of permissive cells supporting the replication of the virus. Usually when polyoma virus replicates in mouse cells, host DNA synthesis is induced concomitant with the replication of the viral genome and the cells are induced to move. At 32° C the *ts3* mutant behaves in these respects like wild type virus, but at 39° C both the induction of cell movement and cell DNA synthesis by *ts3* are impaired. It seems unlikely that the *ts3* gene product directly controls all these responses in both transformed hamster cells and lytically infected mouse cells. More likely the *ts3* gene may specify a molecule which changes the chemistry of the surface of the infected cell. Dulbecco, Eckhart and Burger have evidence (as yet unpublished), that this is indeed the case, but that is another story about which we shall no doubt shortly be reading.

Apart from inducing cellular DNA synthesis, polyoma virus and SV40 replicating in permissive hosts induce the scission of the cell's genome and as a result pieces of cellular DNA of appropriate size are enclosed in capsids of the virus. The possibility that such pseudovirions, as they are called, might be useful as vehicles for the transduction of cells has been widely and often uncritically discussed. The analogy between pseudovirions and certain transducing phage particles remains, however, and Grady, Axelrod and Trilling (*Proc. US Nat. Acad. Sci.*, **67**, 1886; 1970) are among the groups pursuing the possibility of showing transduction of eukaryotic cells. They report that the pseudovirions produced when SV40 replicates in the VERO line of monkey cells contain repeated and unique sequences of monkey DNA in the

same proportions as in the total VERO cell genome. This indicates, of course, that particular portions of the genome are not preferentially incorporated into pseudovirions. There is therefore no hope of using pseudovirions as analogues of lambda phage for specialized transduction, that is, the transduction at high frequencies of specific portions of a cell's genome. A putative general transducing potential remains, however, and Grady *et al.* have been able to show that about 10 per cent of the SV40 pseudovirions taken up by mouse embryo cells enter the nucleus carrying their monkey DNA undegraded. Whether such foreign DNA is replicated and expressed, however, is the crucial question which remains unanswered.

AAAS

Model Biology

by our Special Correspondent

THE increasing infiltration of biology by physical scientists and mathematicians was plainly reflected in the contribution of the American Mathematical Society to this year's meeting of the American Association for the Advancement of Science (AAAS) in Chicago. The section was devoted entirely to biology; in particular to two of its most fascinating and intractable problems—control of development and central nervous function.

Dr B. C. Goodwin (University of Sussex) gave a somewhat apologetic talk

Synthesis of Bacterial Cell Walls

FUNCTIONS for the teichoic acids of Gram positive bacteria are proposed in three articles in next Wednesday's *Nature New Biology* (229, 1971) by a team in the Microbiological Chemistry Research Laboratory at the University of Newcastle upon Tyne. Teichoic acids are polymers composed either of glycerol phosphate or ribitol phosphate, and the principal chain has substituents on it which vary in different strains and include D-alanine, D-glucose, D-galactose or amino sugars. The Newcastle team (J. Baddiley *et al.*), who first discovered these polymers, suggested in 1961 (*Nature*, **191**, 570) that they participate in ion-exchange reactions in the outer regions of the bacterial cell; the phosphate of the teichoic acids bestows on the isolated cell walls the ability to bind magnesium ions. Baddiley *et al.* have now isolated functioning cell wall-cell membrane fragments and cell membrane fragments without any cell wall. These fragments catalyse the synthesis of teichoic acids from UDP-glucose and CDP-glycerol and require high concentrations of Mg^{2+} ions. Pre-incubation of the fragments of wall membrane made the membrane enzymes independent of Mg^{2+} concentration; the pure membrane fractions required 10 mM Mg^{2+} for maximal activity. This is a direct demonstration of the function of teichoic acids in binding Mg^{3+} ions and making them available to the membrane.

Chloramphenicol inhibits the biosynthesis of the teichoic acid of a bacterial cell wall in a cell-free system of fragmented cell membrane from *Bacillus subtilis*. The chloramphenicol seems to inhibit the synthesis of the teichoic acid directly and is unrelated to protein synthesis. The antibiotic inhibits the transfer of glucose from UDP-glucose to the polymer. The inhibition is specific for glucose and does not occur with other nucleotide sugars. The polymer, whose synthesis is inhibited, is unusual in that the glucose is

in the principal chain, rather than attached to it.

The biosynthesis of several bacterial wall polymers uses lipid intermediates that aid in the transport of sugar residues and related components from intracellular nucleotide precursors through the membrane to the polymer chains in the wall. Such lipid intermediates are used in the biosynthesis of both teichoic acids and peptidoglycan, which is a polymer containing N-acetylmuramic acid, N-acetylglucosamine, alanine, lysine, glutamic acid and glycine.

The Newcastle group has further examined the possibility that a single lipid may be common to both pathways, and that therefore the availability of this lipid could play some part in the regulation of cell wall synthesis. The amount of these lipids in bacteria is very small and their direct study is therefore difficult. Consequently, Baddiley and his colleagues looked for an indirect answer. If in a single organism the identical lipid is involved in the synthesis of the two polymers, competition for the limited amount of lipid would probably occur. They therefore examined the relationship between the biosynthesis of peptidoglycan and teichoic acid in cell membrane fragments from *Staphylococcus lactis* I3. Inhibition of teichoic acid synthesis is achieved by the addition of peptidoglycan precursors. The antibiotics bacitracin and vancomycin inhibit the formation of peptidoglycan and prevent the return of the lipid intermediate to the pool; the antibiotics have no direct effect on teichoic acid biosynthesis. When peptidoglycan precursors and the antibiotic are added together, however, thereby effectively trapping the lipid intermediate, a sharp inhibition of teichoic acid synthesis results. It follows that the same lipid carrier molecules are used for transporting precursors of both peptidoglycan and teichoic acid.