cause the formation of a crust impregnated with organic matter in the surface sand. When this is fragmented by further usage, erosion results. Dr Liddle has experimented with a variety of remedial operations in the early stages of damage and has found that harrowing of the compressed surface accompanied by the addition of fertilizers significantly improves recovery by encouraging the rapid growth of meadow grasses (*Poa pratensis* and *P. annua*).

Experiments of this type may enable descriptive recreational ecology to develop into informed and effective environmental surgery.

PROVIRUSES

Site of Synthesis

from our Cell Biology Correspondent

MOST stocks of most viruses seldom contain more than one infectious particle per ten physical particles and more often contain only one infectious unit per hundred or thousand physical particles. It is therefore almost impossible to determine unambiguously how an infectious virus particle enters a cell and brings about an infectious cycle. One can, however, use the electron microscope coupled with autoradiographic techniques to observe the fate of a sample of the virus particles in a population that is allowed to adsorb to any particular population of cells and assume—this assumption is usually tacit—that the minority of infectious units behave like the majority of physical particles. Dales, Hanafusa and Huima (*Virology*, **50**, 440; 1972) have just done this in a painstaking way using Schmidt Ruppin Rous sarcoma virus and an avian leukosis virus (RAV 2) labelled with ^3H -uridine and secondary cultures of chick embryo fibroblasts. Their conclusions, in particular about the site of synthesis of the DNA provirus, are at variance with those of Hatanaka *et al.* who last year reported somewhat similar experiments with murine RNA tumour viruses and mouse cells.

Dales *et al.* allowed the virus particles in the presence of DEAE dextran to adsorb to the cells for 1 h at 4° C or 10° C during which time from 500–2,000 virions irreversibly adsorbed to each cell. Electron micrographs of these cells revealed two interesting points. First, cells comprising about 4–18 per cent of the population which have a distinct morphology different from the majority of fibroblasts adsorb many more virions than typical fibroblasts. Second, the adsorption sites at the surfaces of fibroblasts have a distinct structure; they consist of a reticulum of fibrils and resemble gap junctions.

When cells are warmed from 4° C or 10° C to 37° C, a minority of the virions

are phagocytosed and transported in phagocytic vacuoles to the vicinity of the nucleus and, by 10 min after the temperature shift, labelled RNA molecules can be detected by autoradiography in the nuclei of the cells. The amount of label in the nucleus reaches a maximum about an hour after the temperature shift, and a simple piece of arithmetic based on several not unreasonable assumptions indicates that if all the ^3H -uridine remains in viral RNA there must be between twenty-five and seventy-five genome equivalents per infected cell nucleus after an hour at 37° C. This, of course, is only a small fraction of the number of virions that adsorb to the cells. Aware of the problem that viral RNA may be degraded and the ^3H -uridine re-utilized in the synthesis of cell RNA, Dales *et al.* performed appropriate control experiments which eliminated this possibility.

Clearly the infection of chick cells by these avian tumour viruses involves the rapid transport of viral RNA to the cell nucleus, and this finding suggests, of course, that the synthesis of a DNA provirus by reverse transcription may well occur in the cell nucleus, which is what one might have predicted. But Hatanaka *et al.* have reported that soon after mouse cells are infected with mouse sarcoma or leukaemia virus, DNA synthesis can be detected by ^3H -thymidine autoradiography in the cytoplasm and this synthesis, they suggested, was the synthesis of DNA proviruses. By contrast Dales *et al.* apparently found no evidence of cytoplasmic DNA synthesis above background in chick cells infected with avian viruses. They therefore believe that at least in the avian system proviruses are made within the cell nucleus, not in the cytoplasm.

The complete discrepancy between the conclusion of Dales *et al.* and that of Hatanaka *et al.* merits resolution. The generous way to account for the difference is simply to suggest that the avian and murine systems are quite different, but *a priori* that seems most

unlikely. Perhaps Dales *et al.* could be persuaded to repeat their experiments using murine viruses and cells.

EUTROPHICATION

Is pH the Clue?

from a Correspondent

THE changes in fauna and flora—eutrophication—which result from the addition to lakes or estuaries of large quantities of inorganic nitrates or phosphates from sewage and agricultural run-off are often deleterious, especially to fisheries. The mechanisms of these changes, however, are not always clear. O'Brien and de Noyelles (*Ecology*, **53**, 607; 1973) now show that pH changes may often be implicated.

The initial response to over-fertilization is usually a "bloom" of one or more species of alga. Intense blooms are followed by the disappearance of many animal species, those with high oxygen requirements often seeming to be most susceptible. These disappearances have been ascribed to toxins produced by the algae, to low oxygen concentrations resulting from bacterial decomposition of dead algae, and to high oxygen concentrations caused by rapid photosynthesis.

O'Brien and de Noyelles applied one of three fertilization treatments to each of six experimental ponds of equal size, with two remaining as unfertilized controls. For two summers, various environmental parameters were monitored, together with the densities of plant and animal species. In the ponds with the most fertilizer, summer increases in phytoplankton were followed by disappearances of crustacean zooplankton. These disappearances occurred at different plant densities but at almost identically high pH values. Furthermore, water from these ponds could be detoxified by 'Millipore' filtration, which caused a slight reduction in pH with no change in oxygen tension.

The authors present further evidence which, though not conclusive, suggests

Nuclear Giant RNA Contains Messenger Sequences

THERE has recently been a great deal of indirect evidence that heterogeneous nuclear RNA is a precursor of mRNA. Direct proof of this is provided in last Wednesday's edition of *Nature New Biology* (**241**, 66; 1973) by Williamson, Drewienkiewicz and Paul, of the Beatson Institute of Cancer Research, Glasgow. These authors report that when heterogeneous RNA with a high molecular weight from mouse reticulocytes was injected into *Xenopus* oocytes along with labelled histidine, and the soluble fraction was then analysed on a 'Sephadex G-100' column, radioactivity

was located in the haemoglobin region. Material from this region co-chromatographed with mouse globin chains when examined on CM cellulose. Similar labelled material from a control experiment did not chromatograph as globin. When brain heterogeneous RNA, isolated in the presence of a reticulocyte lysate, was injected into *Xenopus* oocytes no globin synthesis occurred.

These results eliminate the possibility that aggregation of heterogeneous RNA and cytoplasmic globin mRNA had occurred during the preparation of the heterogeneous RNA fractions.