also indicated by the X-ray and radio data (ref. 11 and S. L. Shapiro, A. P. Lightman and D. M. Eardly, unpublished).

We thank J. Katz for discussions.

TSVI PIRAN
JACOB SHAHAM

Racah Institute of Physics,
The Hebrew University, Jerusalem, Israel

Received March 10; accepted May 21, 1975.

- ¹ Ruderman, M., *Proc. seventh Texas Symposium* (in the press).
- ² Piran, T., Shalom, J., and Katz, J., *Astrophys. J. Lett.*, **196**, L107-L108 (1975).
- ³ Thorne, K. S., *Astrophys. J.*, **191**, 597-619 (1974).
- ⁴ Cline, T. L., and Desai, U. D., *Astrophys. J. Lett.*, **196**, L43-L46 (1975).
- ⁵ Tananbaum, H., et al., *Astrophys. J. Lett.*, **177**, L5-L10 (1972).
- ⁶ Pringle, J. E., and Rees, M. J., *Astr. Astrophys.*, **21**, 1-9 (1972).
- ⁷ Lightman, A. P., and Eardly, D. M., *Astrophys. J. Lett.*, **187**, L1-L3 (1974).
- ⁸ Lamb, D. Q., Lamb, F. K., and Pines, D., *Nature phys. Sci.*, **246**, 52-54 (1973).
- ⁹ Palumbo, G. G. C., et al., *Astrophys. J. Lett.*, **189**, L9-L11 (1974).
- ¹⁰ Strong, I. B., and Klebesadel, R. W., *Nature*, **251**, 396-397 (1974).
- ¹¹ Thorne, K. S., and Price, R. H., *Astrophys. J. Lett.*, **195**, L101-L106 (1975).

Mother and baby paradox

THE 'twin paradox' of relativity theory describes the difference in the elapsed time between the twin who stays at home and the one who goes on a space flight and returns. No genuine paradox is involved; elapsed time is not in general independent of the state of motion. In this case the difference in the elapsed time is related to a difference in the (locally measurable) acceleration histories of the twins; the space traveller must suffer an acceleration, however brief, in order to return.

But no paradox is involved, even if there is no locally measurable effect to distinguish the two timekeepers. General relativity provides such possibilities, one of which I describe here.

A mother, desiring her baby to remain young for as long as possible, decided on the following plan of action. Each night, when the baby was asleep, she went out and collected masses from afar, symmetrically, and arranged them in a spherical shell around the baby's crib (herself staying outside and some distance away from the completed shell). The baby's sleep was not disturbed, no acceleration being measurable inside the shell. In the morning before the baby woke up, she dismantled the shell and stored the masses again far away. Doing this carefully, with spherical symmetry, again no acceleration could be felt by the baby. But, in the 12-h period of the night experienced by the mother, the baby had aged only $12 \text{ h} \times (1 - GM/Rc^2)$ (for small values of GM/Rc^2 , where M and R refer to the mass and radius of the shell, and G and c are the gravitational constant and speed of light respectively). In the daytime the components of the shell were stored so far apart that the large value of R made the value of GM/Rc^2 negligibly small.

By this procedure the mother's and her baby's timekeeping were caused to be different; yet, unlike the case of the 'twin paradox' of special relativity, without any necessity for a different acceleration history.

T. GOLD

Center for Radiophysics and Space Research,
Space Sciences Building, Cornell University,
Ithaca, New York 14853

Received May 13; accepted May 23, 1975.

Trapped helium and argon and the formation of the atmosphere by degassing

I REPORT here some radiogenic ^4He and ^{40}Ar data measured in the glass margins of deep sea pillow basalts, and comment on their applicability to discussions of formation of the Earth's atmosphere by degassing from a solid Earth.

Discussions of such degassing generally assume either catastrophic or continuous degassing (see refs 1, 2 and references therein). In fact, there is no reason why these two sets of models should be considered to be mutually exclusive. The history of the Earth may include both a catastrophic event with an associated 'big burp', and later continuous degassing.

Ozima and Kudo² and Ozima³ have presented calculations which indicate the evolution of the terrestrial $^{40}\text{Ar}/^{36}\text{Ar}$ ratio as a function of time, in either catastrophic or continuous degassing models of atmospheric formation. In particular, Ozima³ has shown that the present $^{40}\text{Ar}/^{36}\text{Ar}$ ratio in the mantle is a sensitive indicator of the validity of the catastrophic model. In such models, the degassing is defined by any two of these four quantities: The time of degassing; the fraction of Ar degassed; the K content of the Earth; the present value of the $^{40}\text{Ar}/^{36}\text{Ar}$ ratio within the Earth. Ozima showed sets of graphical solutions for these quantities which satisfy the total Ar abundance and the present atmospheric ratio of 295.5.

The rare gases found in deep sea basaltic glasses are clearly those trapped within the rock on its eruption⁴⁻⁶, due to the high hydrostatic pressures and rapid chilling, and as such represent mantle-ambient gases. A characteristic of those trapped gases is the presence of "excess" radiogenic ^4He and ^{40}Ar , in amounts greater than can be accounted for by the decay of U, Th and K subsequent to their eruption⁴⁻⁷. Similar "excess" ^{40}Ar contents have been measured in ultramafic intrusives⁸ and in continental rock, but their origin is not as clear.

Ozima therefore accepted the $^{40}\text{Ar}/^{36}\text{Ar}$ trapped in submarine basalts as representing the isotopic ratio in the upper mantle; the published ratio available to him, $\sim 1,000$, could not be reconciled with a catastrophic degassing model of the Earth's atmosphere without the assumption of an unacceptably high K content³. He noted, however, following Schwartzman¹⁶, that if the ultramafic values for this ratio ($\sim 10,000$) were indicative of mantle gases, then the catastrophic model could not be ruled out. The $^{40}\text{Ar}/^{36}\text{Ar}$ values shown here in Table 1, measured in the glassy margins of deep sea basalts from both the Atlantic and Pacific Oceans, on both ridge and non-ridge samples, reach values as great as $\sim 15,000$. If these values are accepted, Ozima's argument against the catastrophic model must be relinquished. The argument can be replaced, however, with one based on the $^4\text{He}/^{40}\text{Ar}$ data of Table 1.

Schwartzman predicted a $^4\text{He}/^{40}\text{Ar}$ ratio in the present-day mantle of between 1.36 and 2.23, depending on the

Table 1 ^4He , ^{40}Ar , and ^{36}Ar data in glassy margins of deep sea basalts. The rocks are described in refs 4 and 17. All units are 10^{-8} s.c.c. g^{-1} . The ^{40}Ar reported is total measured ^{40}Ar ; correcting for atmospheric contamination does not change the $^4\text{He}/^{40}\text{Ar}$ results significantly

Sample	^{40}Ar	^{36}Ar	^4He	$^{40}\text{Ar}/^{36}\text{Ar}$	$^4\text{He}/^{40}\text{Ar}$
Vema Fracture Zone, Atlantic Ocean					
72	218	0.0175	—	12,500	—
	77	0.024	1,150	3,210	15
	74	0.0167	1,300	4,430	17
88	290	0.055	2,400	5,200	8.3
	330	0.075	2,400	4,400	7.3
92	235	0.025	1,600	9,400	6.8
	204	0.028	2,370	7,300	12
East Pacific Rise, $\sim 10^\circ\text{S}$					
32	62	0.0174	84	3,600	1.35
	81	0.067	180	1,210	2.2
40	81	0.0104	1,250	7,800	15
	38	0.032	500	1,200	13
44	210	0.0133	1,800	15,800	8.6
	250	0.016	1,970	15,600	7.9
East Pacific Seamount, $\sim 10^\circ\text{S}$ — 98°W					
50	157	0.042	2,800	3,740	18