ity for a series of different orientations in the same animal allowed Bonhoeffer and Grinvald⁶ (using intrinsic signals of activity from upper layers of cortex⁷) to show that the pinwheel model is correct.

There are many models of how the cortex develops to achieve this pattern⁸. Cats and macaques show a pattern of orientation selectivity as soon as the eyes open shortly after (or at) birth. But ferrets, which are born at an earlier stage of development, show little orientation selectivity when their eyes open. Instead, orientation selectivity develops to adult levels during the fifth and sixth weeks after birth, under the influence of nerve-cell activity⁹.

Wolf and Geisel¹ now suggest that all models for the development of orientation selectivity are subject to symmetry principles. The authors first assume that columns with a similar orientation preference tend to be a particular distance apart. Their second assumption is that patterns resulting from (i) shifting the pattern parallel to the cortical surface; (ii) rotating the pattern parallel to the cortical surface; and (iii) shifting all orientation preferences by the same angle, will arise with the same probability. Their final assumption is that the pattern arises from many random factors, and that Gaussian statistics apply. Their mathematics then shows that the expected density of pinwheels is given by

$$\rho = \frac{\pi(1+\alpha)}{\Lambda^2}$$

where Λ is the average spacing of columns, and α is a number that describes the structure of spatial correlations. One therefore expects to find at least π pinwheels in an area of size Λ^2 .

In fact, the density of pinwheels varies from 3.75 in an area of size Λ^2 in macaque, to 2.1–2.6 in tree shrew, with intermediate values for squirrel monkey, ferret and cat. Thus, either there is some process that prevents development from being carried to completion in the cat and tree shrew, where the value is substantially below π , or the density of pinwheels develops to a value greater than π , then decreases through annihilation of pinwheels. Wolf and Geisel propose that such an annihilation could occur through melding of a pair of pinwheels — one with clockwise chirality and another with anticlockwise chirality.

The annihilation of pinwheels is an intriguing possibility. But it is not supported by the few observations available, which show a series of orientation maps that does not change much with age⁸. Areas with longer stretches of parallel slabs — which are predicted to arise when two pinwheels merge — do not seem to appear much more frequently in the older patterns. So, Wolf and Geisel's predictions still need to be rigorously tested. Moreover, the mathematics

assumes that there are no intrinsic places in the cortex that would attract the formation of pinwheels. This has been suggested as a factor in positioning of the blobs for colour in the macaque visual cortex¹⁰.

Wolf and Geisel have used mathematical approaches that are common in physics but rare in biology. In turbulent flow, for example, aspects of a system's behaviour can be predicted from basic symmetries without knowing the detailed nature of the interactions between components of the system. Because the nervous system is made up of billions of cells and trillions of synapses, with detailed interactions that are unknowable in practice, mathematical approaches may have a future in predicting the overall behaviour of the system. □

RNA processing

A tale of two tails

David Bentley

The fate of RNAs in the nucleus is determined by the polymerase that made them. For example, the same primary transcript that will give rise to a messenger RNA if made by RNA polymerase II (pol II), will never mature if it is made by pol I, pol III or a bacteriophage RNA polymerase^{1,2}. How can we explain this link between the machines that make and process mRNAs? On page 93 of this issue, Hirose and Manley³ bring us one step closer to an answer. They report that the carboxy-terminal domain (CTD) — a protein domain unique to pol II — acts as part of the protein complex that fashions the 3' end of the mature mRNA by cleavage and polyadenylation.

The formation of 3' ends is a two-step process in which the RNA transcript is cleaved downstream of the sequence AAUAAA, then a poly(A) tail is added at the cut end. RNAs made *in vivo* by pol II lacking the CTD — a conserved repeat of the

Nigel W. Daw is in the Department of Ophthalmology and Visual Science, Yale University School of Medicine, 330 Cedar Street, New Haven, Connecticut 06520-8061, USA.

e-mail: Daw@biomed.med.yale.edu

1. Wolf, F. & Geisel, T. *Nature* **395**, 73–78 (1998).
2. Mountcastle, V. B. *J. Neurophysiol.* **20**, 408–434 (1957).
3. Hubel, D. H. & Wiesel, T. N. *J. Physiol. (Lond.)* **160**, 106–154 (1962).
4. Hubel, D. H. & Wiesel, T. N. *J. Comp. Neurol.* **158**, 267–294 (1974).
5. Braitenberg, V. & Braitenberg, C. *Biol. Cybern.* **33**, 179–188 (1979).
6. Bonhoeffer, T. & Grinvald, A. *Nature* **353**, 429–431 (1991).
7. Grinvald, A., Lieke, E. E., Frostig, R. D., Gilbert, C. D. & Wiesel, T. N. *Nature* **324**, 361–364 (1986).
8. Erwin, E., Obermayer, K. & Schulten, K. *Neural Comp.* **7**, 425–468 (1995).
9. Chapman, B., Stryker, M. P. & Bonhoeffer, T. *J. Neurosci.* **16**, 6443–6453 (1996).
10. Kuljis, R. O. & Rakic, P. *Proc. Natl Acad. Sci. USA* **87**, 5303–5306 (1990).

sequence YSPTSPS, found at the carboxy terminus of the pol II large subunit — are not efficiently capped, spliced or cleaved at the poly(A) site^{4,5}. This suggests that the CTD helps to target mRNA processing factors to pol II transcripts^{6,7}. Hirose and Manley have now discovered that the CTD facilitates the 3' RNA cleavage reaction, even in the absence of transcription. They suggest that, in effect, the CTD is a cofactor for 3' processing.

Production of a mature mRNA 3' end can be reconstituted *in vitro* with cleavage/polyadenylation specificity factor (CPSF), cleavage-stimulation factor (CstF), cleavage factors CFI and CFII, and poly(A) polymerase (PAP)⁸. The association of CPSF and CstF with the CTD *in vitro*⁴, and the colocalization of CstF and phosphorylated pol II *in vivo* (Fig. 1), indicate that a stable 'mRNA factory' complex of pol II with processing factors carries out synthesis and

Figure 1 Colocalization of the cleavage-stimulation factor CstF and phosphorylated RNA polymerase II (pol II). Immunofluorescence of *Drosophila* polytene chromosomes with anti-pol II monoclonal antibody (left) and rabbit antibody against Suppressor of forked, the *Drosophila* homologue of CstF (right). (Courtesy of M. Sikes and A. Beyer, University of Virginia.)

Figure 2 Model for activation of 3' processing by the pol II carboxy-terminal domain (CTD). Hirose and Manley³ have shown that the CTD is part of the protein complex that carries out cleavage and polyadenylation to produce a mature messenger RNA transcript. CPSF, cleavage/polyadenylation specificity factor; CstF, cleavage-stimulation factor; CFI, cleavage factor I; CFII, cleavage factor II; PAP, poly(A) polymerase.

maturation of the primary transcript⁹.

Hirose and Manley provide an insight into how the mRNA factory operates. They followed up the curious observation that, in a purified system, cleavage at the poly(A) site is stimulated by creatine phosphate and by phosphoamino acids. They reasoned that these compounds may mimic a phosphoprotein that allosterically activates the cleavage reaction. An attractive candidate phosphoprotein is pol II — the CTD undergoes a cycle of hyperphosphorylation and dephosphorylation (mainly at residues two and five of the YSPTSPS repeat¹⁰) as pol II cycles through transcriptional initiation, elongation and termination. Sure enough, the authors found that phosphorylated pol II and recombinant CTD stimulate cleavage *in vitro*. Unexpectedly, however, the unphosphorylated CTD was also effective.

These experiments show that the CTD stimulates 3' processing in the absence of transcription, although it is not clear whether the CTD is acting in the same way as creatine phosphate. Hirose and Manley also found that the cleavage reaction was inhibited when they immunodepleted pol II from a crude nuclear extract, after dissociating poly(A) factors from the CTD with high salt. Moreover, they could restore cleavage activity by adding back pure pol II. In other words, the CTD stimulates cleavage in a crude extract, as well as in a purified system.

But how does the CTD stimulate cleavage in the absence of transcription? Does it promote the formation of a stable complex (within which cleavage can then occur), or does it contact polyadenylation factors only fleetingly, triggering cleavage? Assembly of a complex — which might comprise CstF, CPSF, CFI, CFII and PAP, together with the RNA (Fig. 2) — may be rate limiting both *in vivo* and *in vitro*. The CTD could accelerate assembly of the complex by acting as a scaffold that positions the proteins optimally, all in the same place at the same time. A scaffold might even impose an order of assembly of the polyadenylation machine.

Hirose and Manley suggest that the CTD is not just a passive scaffold but an active participant in the cleavage reaction. According

to this model, which is more consistent with the effect of creatine phosphate, the CTD behaves as a cofactor or allosteric activator. One possible target for allosteric activation is CstF, which binds the CTD *in vitro*⁴. The importance of this interaction for scaffolding or allosteric activation could be tested using CTD mutants that do not bind CstF.

It is now clear that pol II stimulates 3' processing, both *in vivo* and *in vitro*, through the CTD. Conversely, poly(A) factors may affect pol II by influencing the decision to terminate or elongate the RNA chain. Termination depends on transcription of the poly(A) signal¹¹, poly(A) factors¹² and the CTD (ref. 4). Signalling from the polyadenylation machine to the polymerase may ensure that the chain is terminated only after pol II has reached the end of a gene. So, a two-way line of communication between pol II and the 3' processing factors may be built into the mRNA factory.

Working out how the machines in the mRNA factory interact with one another is a big challenge. Hirose and Manley³ have provided the first *in vitro* system to study one aspect of the complex communication system that integrates different nuclear processes through protein–protein interactions. We can look forward to the development of more *in vitro* systems, which will reveal the details of how the CTD affects splicing and capping as well as polyadenylation. □

David Bentley is in the Department of Biochemistry and Molecular Genetics, UCHSC, 4200 East 9th Avenue, Denver, Colorado 80262, USA.
e-mail: David.Bentley@UCHSC.edu

1. Sisodia, S. S., Sollner, W. B. & Cleveland, D. W. *Mol. Cell. Biol.* **7**, 3602–3612 (1987).
2. Smale, S. T. & Tjian, R. *Mol. Cell. Biol.* **5**, 352–362 (1985).
3. Hirose, Y. & Manley, J. L. *Nature* **395**, 93–96 (1998).
4. McCracken, S. et al. *Nature* **385**, 357–361 (1997).
5. McCracken, S. et al. *Genes Dev.* **11**, 3306–3318 (1997).
6. Greenleaf, A. L. *Trends Biochem. Sci.* **18**, 117–119 (1993).
7. Yuryev, A. et al. *Proc. Natl Acad. Sci. USA* **93**, 6975–6980 (1996).
8. Keller, W. & Minvielle-Sebastia, L. *Curr. Opin. Cell Biol.* **9**, 329–336 (1997).
9. Jackson, D. A., Hassan, A. B., Errington, R. J. & Cook, P. R. *EMBO J.* **12**, 1059–1065 (1993).
10. Dahmus, M. E. *J. Biol. Chem.* **271**, 19009–19012 (1996).
11. Logan, J., Falck-Pederson, E., Darnell, J. & Shenk, T. *Proc. Natl Acad. Sci. USA* **84**, 8306–8310 (1987).
12. Birse, C., Minvielle-Sebastia, L., Lee, B., Keller, W. & Proudfoot, N. *Science* **280**, 298–301 (1998).

Daedalus

Zero-tolerance policing

Millions of potentially lethal cell mutations occur in each of us every day. We notice nothing; the immune system arrests and executes the offenders instantly. But with advancing age, its vigilance slackens. Mutations start to go unchecked; and we succumb to the melancholy neoplastic or degenerative diseases of old age.

So Daedalus is extending his scheme of last week for priming the immune system against all normal biological proteins. He now wants to prime it against abnormal proteins as well — in particular, the abnormal ones expressed on the surfaces of malfunctioning human cells.

DREADCO volunteers are now providing small biopsy tissue samples of blood, lung, gut, and so on, to be cultured in the usual way. The cultures will then be attacked with mutagens, ultraviolet light, X-rays, and such, to induce as many mutations in them as possible. The battered and misfiring cells will express all sorts of crazy and altered proteins on their surfaces. These will be extracted, and injected back into the volunteer from whom the cells came. This 'auto-immunization' will prime his immune system against such cell mutations. Should a cell in his gut or liver start to go crazy in one of those pre-identified ways, the immune system will spot the abnormality at once. It will stamp on the rogue cell instantly.

Sadly, auto-immunization can never be a perfect defence. There are too many different body tissues, each with too many ways of going wrong, for any vaccine to contain all the proteins expressed by all these errors. But not all errors are equally likely, or equally deadly if they do occur. The human genome can suffer about 10 billion possible single-site mutations; yet there are only some 30,000 recognized illnesses of all kinds. The deadly diseases are those in which a mutated cell starts to multiply, or to trigger similar changes in its neighbours. So Daedalus will tune his treatment to his customers. If all your ancestors died of liver cancers, you'll include some liver cells in your tissue sample; if Dr Alzheimer stalks your family tree, you'll make sure that brain cells are well represented in it. And the mutagenic regime itself will be chosen to provoke the most appropriate type of cell damage. The resulting immunization will be an excellent defence against the disease of your fears. Sadly, it won't make you immortal. But you'll last much longer, and finally die of something you weren't expecting.

David Jones