news and views

Destruction of stratospheric ozone?

from Michael A. A. Clyne

STRATOSPHERIC ozone will eventually be depleted significantly if releases of chlorofluoromethanes (CFM) continue at the present rates. Legislation is not necessary now, but if after no more than two years, existing evidence still supports significant ozone depletion, action to restrict manufacture should be taken by governments. These are among the main recommendations of the Committee on Impacts of Stratospheric Change, (*Halocarbons: Environmental effects of chlorofluoromethane release*, National Academy of Sciences, Washington DC, 1976) set up by the US National Academy of Sciences.

In 1974 I reviewed Molina and Rowland's article (*Nature*, **249**, 810) in the same issue of *Nature*, on the effect of chlorofluoromethanes on stratospheric ozone. Then, neither I nor most other scientists, believed that the hypothesis presented by Molina and Rowland, and independently by Stolarki and Cicerone (*Canad. J. Chem.*, **52**, 1610; 1974) would in fact develop into one of the most important recent issues in environmental pollution. Briefly, this theory, which has now largely been supported by an authoritative report of the US National Academy of Sciences (*Halocarbons: Effects on stratospheric ozone, Panel on Atmospheric Chemistry*, National Academy of Sciences, Washington DC, 1976), is as follows. CFM 11 (CFCl₃) and 12 (CF₂Cl₂), which contain chlorine, fluorine and carbon only, are produced in large amounts in all developed countries of the world. Approximately half of the total output is used as propellants in aerosol cans for

personal consumer uses (such as hair-sprays and deodorants). Because of the stability that makes these compounds suitable for aerosol cans and for other uses, CFM 11 and 12 are very inert and no known major sinks for these compounds have yet been identified in the lower atmosphere (troposphere). Hence a large fraction of all past production of CFM 11 and 12, whether used for aerosols, car air conditioners or refrigerators is still present in the troposphere and indeed, these compounds will continue to accumulate there. Diffusion to higher altitudes occurs (Fig. 1), up to the stratosphere where most of the Earth's ozone (O₃) shield is located. The CFM can be photochemically dissociated there by far ultraviolet light at wavelengths around 190 nm. This light is not completely removed at stratospheric altitudes by the ozone and oxygen that normally filters out ultraviolet light, at wavelengths less than 300 nm, from reaching the Earth's surface. Photochemical action releases chlorine atoms from the CFM; and chlorine atoms react rapidly with ozone, giving chlorine monoxide (ClO) free radicals: $\text{Cl} + \text{O}_3 \rightarrow \text{ClO} + \text{O}_2$. . (1). The reaction (1) is then followed at once by the rapid reaction (2) of ClO with the large quantities of the free oxygen atoms present at stratospheric altitudes between 25 and 50 km: $\text{ClO} + \text{O} \rightarrow \text{Cl} + \text{O}_2$ (2). Chlorine atoms are thus recycled, and are able to consume more ozone; this sequence is termed a chain reaction. Finally, the cycle is terminated by removal of chlorine atoms in their

reaction with methane

Most of the HCl formed in reaction (3) is not recycled back to active chlorine. Although many other reactions are involved, some of which are mentioned below, the overall effect of CFM is thus clearly to deplete stratospheric ozone. The NAS reports conclude that this description of the problem, essentially as advanced by Molina and Rowland, cannot be seriously questioned, although there are new factors that probably somewhat diminish the effect on ozone depletion that they calculated.

There is no justification for the total rejection of an effect of CFM on ozone depletion, which had been maintained earlier by a few scientists. In their report, the NAS Committee state that "Selective regulation of CFM uses and releases is almost certain to be necessary at some time and to some degree of completeness". In reaching their conclusions, the NAS were guided by a panel of distinguished experts including one Briton, Brian Thrush, and one German, Dieter Ehhalt. The report is an impressive document, containing a large amount of carefully sifted information that will be invaluable to future workers in atmospheric sciences. The approach is low-key, thoughtful and critical. The evidence accumulated by the panel is a formidable case in favour of an effect of halocarbons on stratospheric ozone. However, it is considered that the risks of a delay of not more than two years in restricting CFM releases are low. Although most scientists will probably agree with these views, some will doubtless point to the admitted wide margin of error in the estimate of a 7% eventual ozone depletion, with CFM 11 and 12 releases at the 1973 level. (According to the Panel, the limits of 2 to 20% ozone depletion have a 95% confidence level).

Some will say that the risk is slight, and that in any case industry is already reducing production of CFM 11 and 12, and is actively seeking substitutes that will have a shorter tropospheric life, and hence be less harmful. It re-

Fig. 1 The atmospheric ClO_x cycle.

mains to be seen whether the search for acceptable substitutes for CFM 11 and 12 is successful; this is not likely to be an easy problem. CFM 22 (CHF_2Cl), among others, has promise as a substitute for CFM 11 and 12 in spray cans.

Other people will maintain that the risk is unacceptable, and that legislation is needed now rather than (possibly) in two years time. They will point to the possibility that delay might lead to politicians and governments losing sight of a problem which is effectively postponed for two years, a time scale comparable to intervals between elections. However, it should be noted that as well as studies by US government authorities, both the EEC and the West German government are initiating substantial investigations into the CFM-ozone problem.

The British government published their report on chlorofluoromethanes and their effect on stratospheric ozone earlier in the summer. This report was prepared by the Department of the Environment (DOE) with considerable assistance and guidance from AERE, Harwell. Their conclusions are broadly in line with those of the NAS report, although they do not recommend any specific regulatory action by governments. The DOE report is a useful summary of results; although clearly the resources that are evident in the NAS report were not available to the DOE compilers. The referencing of the DOE report is rather random and non-specific, and some literature citations are an encouragement to whimsical day-dreaming! For instance: ICI (1976), Imperial Chemical Industries, private communication. Dubreuilh W. (1896), des hyper-keratoses circonscrites; *Ann. Dermatol. Syphilig.* 3rd Series, 7, 1158-1204; quoted by Blum (1956).

In order to predict the effects of CFM on stratospheric ozone, a wide variety of data are required. The scope and uncertainties of these data varies enormously. Some of the basic types of information needed are (1) direct measurements of Cl atom, ClO radical, and other concentrations in the stratosphere; (2) the reaction rates of key elementary chemical reactions, as determined in the laboratory, and applicable to stratospheric temperature ($\approx 220\text{ K}$); (3) knowledge of the transport of air masses from the troposphere to the stratosphere. The first type of data (1), is perhaps the most important, because direct measurement of concentrations is the final check that the correct information under categories (2) and (3) is being used. Unfortunately, data of type (1) are by far the most difficult to obtain and are extremely sparse. The only results (so far) are *in situ* measurements of Cl and ClO

How rigid the nucleotide?

from a Correspondent

CRYSTALLOGRAPHERS have become used to seeing every structural idiosyncrasy they unearth in the crystal structure analysis of molecules of biological interest become a peg upon which the edifice of some molecular biological theory is hung. This has been particularly true for the study of the nucleotide and amino acid monomers from which nucleic acids and proteins are constructed. Beginning with the use of the covalent bond lengths and angles observed in monomers in the design of molecular models of these biological polymers, the results of crystal structure determinations have been extended to provide constraints which might properly be assumed for hydrogen bond geometry, for the conformations which can be expected to be preferred about various single bonds and, for the patterns of arrangement of water molecules and ions around biological polymers. This fertile field has been exploited with ever increasing confidence and perhaps nowhere more than in the analysis of the relationship between the properties and structure of nucleic acids. An important development in this work has been the rigid nucleotide concept

which suggests that mononucleotides have a rather rigid conformation which is conserved when they are joined together to form a polymer. X-ray diffraction studies of nucleotides, nucleosides and crystalline polynucleotides have provided a great deal of evidence for this although a number of recent X-ray single crystal studies of nucleotides have suggested that the situation may well be rather more complicated than was at first thought. The need for caution in applying the rigid nucleotide concept too insensitively is further emphasised by Evans and Sarma in a recent issue of *Nature* (263, 567; 1976). From NMR studies they suggest that adenine-containing nucleotides and the oligomers derived from them are no more rigid than the nucleoside adenosine itself. These studies are of course of particular interest because they relate to the situation in solution. An important application of the results of this work which is considered by the authors lies in the analysis of the structure of yeast phenyl alanine tRNA determined by X-ray single crystal studies and in particular the correlation of nucleotide conformation and base-stacking.

in the stratosphere from two balloon flights carried out by J. G. Anderson (University of Michigan) in summer 1976 at Palestine, Texas. Both concentrations of Cl and ClO were determined by Cl atom atomic resonance fluorescence, which has been successfully used also for O atom and OH radical measurements in the stratosphere, by Anderson (*Geophys. Res. Lett.*, 2, 231; 1975; *Geophys. Res. Lett.*, 3, 165; 1976). ClO radicals were reacted with nitric oxide to give Cl atoms, which were detected by atomic resonance fluorescence. The technique was adapted from flow methods for laboratory measurements of Cl atoms and ClO radicals that were developed recently (see Bemand and Clyne, *J. Chem. Soc. Faraday II*, 71, 1132; 1975). Anderson's first Cl and ClO experiments have not so far led to satisfactory results, although the prospects for the atomic resonance and other methods look promising.

Additional direct verification is in principle obtainable from measurements of ultraviolet light intensity at ground level. Whilst such measurements have been made for a number of years, the natural variation of ultraviolet levels at the ground is large. The effects of CFM would probably only become observable in this way when

the ultraviolet level was already increased by a significant, probably unacceptable amount.

Data of type (2), the rates of chemical reactions, are considered carefully in the NAS panel report. One basic difficulty was to identify correctly the major reactions, of which there are a large number. The recently-proposed role of chlorine nitrate, ClONO_2 , has necessitated the introduction of several new reactions. The elementary reaction rates of chlorine nitrate are not, however, all well-defined. Fortunately, the model for ozone depletion is not very sensitive to these rates, and some confidence may be attached to the NAS panel's conclusion that the ozone depletion is reduced by a factor of about 1.7 by inclusion of chlorine nitrate in the model.

The NAS panel conclude that the largest uncertainty in the ozone depletion model is due to uncertainties in the relevant chemical reaction rate constants. They carefully identify the reactions introducing the greatest uncertainty as

followed by

These reactions are involved in the sink processes by which active chlorine

(Cl and ClO) is removed from the ClO_x cycle. Surprisingly, the two reactions of the ClO_x cycle, that is $\text{Cl} + \text{O}_3 \rightarrow \text{ClO} + \text{O}_2$ and $\text{O} + \text{ClO} \rightarrow \text{Cl} + \text{O}_2$, do not introduce large errors, because their rates have been carefully measured in the laboratory. The first measurements of $\text{Cl} + \text{O}_3$ and $\text{O} + \text{ClO}$ rates by Bemand, Clyne and Watson at Queen Mary College in 1973 (see *J. Chem. Soc., Faraday I*, **69**, 1356; 1973; Clyne and Watson, *J. Chem. Soc. Faraday I*, **70**, 2250; 1974) have been only slightly altered as a result of subsequent work in the USA and in this laboratory.

Significant uncertainty ($\pm$ a factor of 1.7) in the prediction of ozone depletion is also introduced by the one-dimensional model used by the NAS panel to approximate the distribution and transport of the chemical species involved. This model is equivalent to averaging the concentrations, motions and chemical reactions over latitude and longitude, leaving only their dependences upon altitude and time. This approximation is considered to be a

good one. For instance, the equivalents of averaged gas transport rates appear in the model as the vertical eddy mixing or transport coefficient, which depends only on the altitude.

One further very important question concerns the effects of an increased ultraviolet intensity (UV-B, 280 to 320 nm) at ground level on animals and plants. The biological effects of UV-B are not as extensively dealt with by the NAS reports as are other aspects. This may be due to the relative lack of knowledge in this area. It is clearly to be identified as a priority area for future work on the CFM-ozone problem. However, in general terms, the deleterious effects of UV-B radiation on living cells are well known. For example, there is a clear causal connection between the more common (but less serious) forms of skin cancer (basal and squamous cell carcinomas) and UV-B exposure. The connection between the less common, but often fatal malignant melanoma and UV-B, is suggestive but not certain, according to

the NAS report.

Release of pollutants other than CFM, which can also affect ozone levels, is another topic requiring extensive future work. Among halogen compounds, further work on methyl chloride formed by the burning of vegetation, is indicated. Also, Crutzen has suggested that methyl chloroform (CH_3CCl_3), whose manufacture as a degreasing agent is considerable and possibly increasing, may eventually cause ozone depletion comparable to that of CFM 11 and 12. Bromine compounds, released for example as methyl bromide in soil fumigants, are probably not produced in large enough quantities to have major effects. However, Br atoms and BrO enter into a similar, but more effective cycle to that of Cl and ClO.

The rate constants for the bromine reactions are not particularly well known. Nitrous oxide, apparently re-

THE *in vitro* cultivation of malaria parasites has been possible for many years but only one or a few successive cycles of development have been achieved and these successes have been mainly with the malaria parasites of monkeys. The amount of physiological and biochemical knowledge that has accumulated from such studies has been vast and it cannot be disputed that much of it is directly or indirectly applicable to chemotherapeutic and immunological studies aimed at the control of malaria. What is desperately required, however, is the long term continuous cultivation of human malaria parasites which cannot be maintained in standard laboratory animals. For some time now there have been rumours of the successful cultivation of *Plasmodium falciparum* in the United States and now two independent reports have appeared, giving the long-awaited experimental details (Trager and Jensen, *Science*, **193**, 673; 1976; and Haynes *et al.*, page 767 in this issue of *Nature*). There can now be no doubt that the continuous cultivation of *P. falciparum* can be achieved in simple media and that it should shortly be possible to obtain the yields of parasites necessary for the production of experimental vaccines.

Both media are remarkably simple. Trager and Jensen use a modification of an earlier medium (RPMI 1640 with HEPES buffer and serum) for the cultivation of *P. coatneyi* from monkeys (Moore *et al.*, *J. Am. med. Ass.*, **199**, 519; 1967) but substitute human serum for the monkey serum originally used. In essence, the medium

Cultivation of human malaria

from F. E. G. Cox

flows over a shallow bed of red blood cells which are disturbed as little as possible in an atmosphere of low O₂ and high CO₂. The technical details are still being modified and Trager and Jensen began by using 7% CO₂ and 5% O₂ but changed this to 1% O₂ in later experiments with better results. The technique of Haynes *et al.* is more traditional, involving a supplemented medium 199 with 1% foetal bovine serum in an atmosphere of 3% CO₂. No doubt many other ways will be found to grow *P. falciparum*, and Trager and Jensen report success using their medium in a candle jar (a container in which a candle has burned and extinguished).

The most important features of the cultivation techniques described seem to be the high levels of CO₂ and the fact that the number of red cells infected rarely rises above 1%, thus preventing the culture medium from becoming too acidic as a result of the metabolic activities of the parasites, which is a problem inherent in this kind of experiment. Although the multiplication of parasites recorded is low (a four-fold increase per cycle is assumed by Trager and Jensen) the numbers of parasites available for collection can be considerable if parallel cultures are set up and initiated from diluted cultures

at the end of each cycle. Haynes *et al.* obtained eleven successive cultures and Trager and Jensen more than twenty-five (and Trager has since reported continuous cultivation over 105 d, implying fifty or more cycles) so there is no reason why such cultures should not be maintained indefinitely.

The technique described by Trager and Jensen looks more promising than that of Haynes *et al.*, but the latter authors have made a major advance by showing that their cultures can be initiated from frozen blood samples. It should therefore be possible to initiate a culture and to freeze stabilates from that culture for the initiation of others. This would save the lives of countless owl monkeys (*Aotus trivirgatus*), which at the moment are the only sources of laboratory maintained *P. falciparum*, and also standardise experiments carried out in different laboratories.

The implications of these experiments are considerable, for as well as the biochemical and physiological observations which are now made feasible it should be possible to grow the numbers of merozoites (the stages that pass from one cell to another) needed for producing vaccines. The only homologous vaccine with real potential, in monkeys at least, is that based on merozoites (Mitchell, Butcher and Cohen, *Immunology*, **29**, 397; 1975) and the production of a suitable vaccine free from viruses is only possible from cultured parasites. With 96 million cases of malaria each year and with *P. falciparum* being the major killer the production and evaluation of such a vaccine should not be delayed.

leased by bacterial action on nitrogenous chemical fertilisers, is fairly stable in the troposphere. The possibility of this compound affecting stratospheric ozone levels should certainly be studied.

All in all, the ozone layer is fragile, and wide-ranging long-term studies of possible effects of pollutants not yet identified are essential. This should include the acquisition of more information on ozone reaction rates. Exclusive concentration on CFM 11 and 12 effects is unwise, since it is quite possible that other deleterious compounds may be present now, or at some future time, due to shifts in industrial production.

I have emphasised here the effects of CFM on the ozone layer and the consequential increase in UV-B radiation at the Earth's surface. It should also be noted that other climatic changes due to CFM cannot be excluded. For instance, Ramanathan (*Science*, **190**, 50; 1975) has stated that CFM 11 and 12 are sufficiently effective absorbers of infrared radiation (near 10 μm) to have a potentially significant effect on the Earth's heat balance.

The role of British science in the CFM-ozone problem is an important one. Much of the basic work leading to the original theories of Stolarski and Cicerone and Molina and Rowland, was carried out in the UK. Lovelock developed a sensitive electron-capture gas chromatograph, with which he was able to demonstrate the presence of various halocarbons arising from man's activities, and to obtain measurements of CFM 11 and 12 at various locations. The programme of work (COMESA) sponsored by the Department of Trade, and carried out partly in university laboratories and partly by the Meteorological Office on possible effects (slight) of Concorde on the stratosphere also

provided valuable input data for the CFM-ozone problem. Current initiatives in the CFM-ozone problem are being taken by industry; research on both CFM 11 and 12 effects, and on substitute CFMs is being carried out. Basic science relevant to stratospheric ozone is being supported by the Science Research Council, and recently the Natural Environment Research Council has announced its intention to concentrate more resources on atmospheric science. The financially small but scientifically significant support over many years in this area by the Royal Society through its Gassiot committee has also been very valuable.

With this expertise available, it is sad therefore that there is no sign at present that the British government is planning any major programme of investigation in stratospheric science related to the CFMs and ozone, although it does contribute through the EEC organisations to the European effort. No more than £100,000 would probably be allocated to such a new investigation, if mounted eventually by the Department of the Environment. Although the restraints of the present economic situation have to be acknowledged, it is a pity, nevertheless that it was not possible to use the experience gained in the COMESA research as the basis of an ongoing and centrally-planned programme of work on atmospheric pollution.

There is a message from all this for those who ask for more "relevant" research programmes, related to current needs of society. Without the undirected, basic research programmes which were carried out, the chances are that the risks to ozone from CFMs would not have been identified until much later, if at all. Identifying the problem is a large part of finding the solution. $\square$

sequences in pure form, by cloning in a bacterial host, has opened up the possibility of studying the sequences flanking the gene which may be important in regulation, and which cannot be obtained sufficiently pure for sequence studies by any other means. The sea urchin (*Psammechinus*) histone gene repeat unit has been cloned in phage lambda and some 25% of the nucleotide sequence determined by a group at the University of Zurich (Walter Schaffner).

The mammalian globin genes are also under determined attack in several laboratories. Charles Weissman (University of Zurich) and Richard A. Flavell (University of Amsterdam) reported the partial purification of the + and - strands respectively of rabbit globin DNA. Weissman and colleagues have isolated the + strand by hybridisation to immobilised globin cDNA; the - strand has been isolated by hybridisation with mercurated mRNA. The final purification step will be to hybridise the two strands, since only complementary sequences should anneal. This reconstituted DNA inserted into a plasmid should provide a source of the complete globin gene sequence including the flanking non-transcribed regions.

Still looking to the future, Bob Williamson (St Mary's Hospital Medical School, London) described a possible use for cloned globin gene sequences as chain-specific probes for the study of certain inherited defects such as the thalassaemias.

Recombinant DNA is also a powerful new tool for probing the immunoglobulin genes. As the first step to tracing changes in the Ig-gene coding sequences predicted by the theory of somatic mutation, Bernard Mach and colleagues (University of Geneva) have constructed plasmids carrying cDNA copies of a mouse myeloma light chain mRNA. The Geneva group has obtained plasmids containing constant and variable region sequences both separately and together. Hybridisation studies have shown that the untranslated region in kappa light chain messenger from different sources is highly conserved.

However, these experiments exemplify some of the conjectural risks which have caused concern about this type of work. The mRNA was not pure and indeed 40% of the initial recombinants contained unknown sequences. More worrying still is that myeloma cells are known to express C-type RNA viruses and that the plasmid carried a drug-resistance marker containing an IS sequence.

This problem is avoided in work from the Laboratory of Molecular Biology in Cambridge reported by Terry Rabbitts. Light chain mRNA

Nuts and bolts of genetic engineering

by Eleanor Lawrence

The Biochemical Society's 9th Harden Conference, entitled the Role of Recombinant DNA in Molecular Biology, was held at Wye College on September 20-24, 1976.

In the present state of the art, a meeting on *in vitro* recombinant DNA must of necessity concern itself more with means than ends. This is partly a result of the need to develop and refine a range of completely new techniques; but it is also a consequence of the almost unique situation in which the novel technology has been applied less

than whole-heartedly, as a result of the voluntary postponement of some experiments until experimental conditions minimising the hypothetical safety risks of this work had been agreed.

But now that experimental guidelines have been published in the United States and Britain, workers are preparing to clone genes from everything from maize to man. A foretaste of the power of these new techniques to investigate otherwise unapproachable problems of the organisation of the eukaryotic genome was given at the meeting.

The ability to obtain specific gene