A GREAT advance in DNA research is reported in this issue: the first complete nucleotide sequence of a DNA genome has been established by Sanger's group in Cambridge (see page 687 of this issue of *Nature*). For the first time it is possible to look at a viral genome as a giant molecule of known structure, all functions of which should be accessible to interpretation in terms of its molecular structure.

The small coliphage Φ X174 is particularly suitable for studies on the structural features underlying the complex processes of gene expression. It has a number of unusual characteristics—its small DNA possesses an unexpectedly high information content and seems to exploit every structural possibility to control gene expression. The single-stranded circular DNA genome of 5,375 nucleotides codes for nine proteins, some of which have been directly sequenced. The amino acid sequence of the others can be deduced from nucleotide sequence data. Genetic and physical maps of the genome have been constructed and have been correlated recently by Weisbeek *et al.* (*Virology* **72**, 61; 1976) who estimated the probable length of each gene. But there is a discrepancy between the calculated coding potential of some regions in the physical map and the proteins produced. Replication and transcription of the genome have also been studied extensively, with some unexpected results. Baas *et al.* (*J. molec. Biol.* **102**, 633; 1976) located the origin of replication in the middle of a structural gene coding for protein A. Hayashi *et al.* (*Proc. natn. Acad. Sci. U.S.A.* **73**, 3519; 1976) reported the presence of two mRNAs covering the A gene region, only one of which was functionally active. The complete sequence has now provided a sound structural basis for these peculiarities.

Φ X174 DNA has been the subject of nucleotide sequence studies from early on. Practically all DNA sequencing techniques have been applied to this molecule, and some have been actually worked out on this DNA. Five years ago classical degradation techniques only allowed the determination of sequences of about 50 nucleotides which made the task of resolving the primary structure of DNA molecules seem much more difficult than the sequencing of RNA. Introduction of new techniques, making use of primed synthesis of highly-labelled complementary strands on Φ X DNA template greatly extended these possibilities. Eventually the new highly efficient 'plus and minus' technique of Sanger and Coulson (*J. molec. Biol.*

Φ X174 sequenced

from Maria Szekely

94, 441; 1975) which allows 150–200 nucleotide long sequences to be read directly, made it possible to produce a complete sequence of the Φ X genome a year after the first (and so far only) complete sequence of an RNA genome had been determined (Fiers *et al.* *Nature* **260**, 500; 1976).

Together with these sequence determinations, information on characteristic structural features of some functional sites of the DNA has also contributed to the construction of a 'chemical map' of the Φ X genome. 5'-terminal sequences of three mRNAs produced *in vitro* have been determined by Smith and Sinsheimer (*J. molec. Biol.* **103**, 699 and 711; 1976); ribosome binding sites on mRNAs corresponding to the initiation of genes B, J, F and G, have been sequenced by Ravetch *et al.* (page 698 of this issue). Homologies with other promoter and terminator sequences as well as the presence of purine tracts at all ribosome recognition sites helped to locate each gene in the sequence. At the same time, these homologies confirmed that the structure suggested for these control signals is generally valid.

The interesting question now arises of how the peculiar properties of Φ X can be interpreted in the knowledge of the primary structure. The most intriguing aspect is how the very high information content can be compressed into a small DNA molecule.

It now seems that this phage has several ways of ensuring the most economical use of its DNA. The most spectacular discovery has already been reported—the same stretch of DNA can code for two proteins which are translated in different reading frames

(*News and Views* **264**, 11; 1976). In 1976 Barrell *et al.* (*Nature* **264**, 34) found that the complete E gene was contained within the D gene region. A similar situation is reported now for genes B and A. Two papers in this issue provide evidence that gene B is contained within the region of gene A: Brown and Smith (page 695) locate the ribosome binding site for gene B within the gene A region, and Smith *et al.* (page 702) assign mutants in gene A and gene B to different reading frames in the overlapping area.

Another DNA stretch which fulfils multiple functions was found at the beginning of the A gene, where the 3'-terminal part of an mRNA which had started at an earlier promoter overlaps with the beginning of the mRNA for the A gene. The sequence of this DNA stretch suggests that a specific hairpin loop structure may be involved in the control mechanism which provides within this overlap a termination signal for one mRNA without interfering with the initiation of transcription of another mRNA. Readthrough at the termination signal may explain Hayashi's finding of two mRNAs covering this area. Two more overlaps occur at the initiation sites of proteins C and J, where the initiation codons overlap with the termination codons of the previous cistrons. The purine tracts preceding the initiation codon serve thus both as recognition sites for ribosomes and as coding sequences for another protein. Gene A seems the most versatile part of the genome: in addition to the multiple functions listed above, it also contains the origin of replication, and thus recognition sites for the proteins involved. The exact site of the origin has not been located, but the region in gene A which contains this site has been defined. Its sequence is less characteristic than could have been anticipated, it shows no symmetry or secondary structure, only a probable specific arrangement of an A–T rich stretch between the G–C rich stretches.

The multifunctional character of Φ X DNA may be a unique property displayed only by a few very small DNA viruses. Overlapping termination and initiation codons have also been detected, however, in the tryptophan messenger of *Escherichia coli* (Platt & Yanofski *Proc. natn. Acad. Sci. U.S.A.* **72**, 2399; 1975), the possibility thus exists that at least some of the above ways to compress more information into DNA may be of more general significance. If this were so, our ideas about the coding potential of DNA genomes would have to be fundamentally altered.