A QUARTER of Yugoslavia's arable land is planted to maize, which vies with wheat as the single most important crop in Yugoslavia. A visit to the Maize Research Institute outside Belgrade proved an eye opener as to how a State Research Institute can be entirely self-supporting from the commercial sale of its products and expertise and at the same time support an active and directed basic research programme which complements its commercial and plant breeding sections. As a novice in this area I may have been over-impressed by the single-mindedness and enthusiasm of the workers at this Research Institute, applied to all stages of the production of a single crop—from research on seed production and gene banks to the extension services on the farm and the sale of seed and expertise (at home and abroad). The increasing interest in growing maize in the UK and Europe both for forage and grain has highlighted the versatility of this plant. It can be grown in a wide range of climates—all that is needed is to find a variety which suits the particular area and the purpose for which it is to be grown.

The Maize Research Institute at Zemun-Polje, about 15 km outside Belgrade, is situated on about 300 hectares of productive land, which is unfortunately now being encroached on by urban and industrial development. Modern well equipped laboratories have a pleasant working atmosphere in extensive fields around a central complex. The Institute employs about 250 people, a quarter of whom have university degrees, and is organised into three different divisions, a science division, an extension service, and an administrative and commercial affairs section. The extension service, which is involved in the controlled production of seed on large cooperative farms and which advises experimental farms, is the crux of the Institute's success. Nearly all the Institute's finance comes from royalties on the sale of its hybrid seeds, both in Yugoslavia and abroad. Nearly a half of its income is from abroad and it has close commercial and research links with more than

Maize in Yugoslavia

from D. O. Hall

20 different countries. Seeds are produced on about 20,000 hectares of good seed-producing land with a suitable climate.

The variability of climate in Yugoslavia from Mediterranean to Continental, its varied geography, and centuries of maize growing, have resulted in a unique maize gene bank. It is claimed that the Maize Institute has more than 100,000 different genetic combinations available for selection of new maize types. The storage of this gene bank is a major part of the Institute's service, since it provides a source of material not only for its own breeding programme but also for research work and hybrid seed production abroad. Yugoslavia seems to have been fortuitously placed in a favourable position for restocking the world's depleted gene banks, especially in North America where the widescale use of hybrid species has resulted in neglect of other varieties selected over the centuries. The Maize Research Institute is making the most of this advantage. Its sales of hybrid seed abroad and patented rights of seed production extend to 18 countries and these hybrids are now grown on over 1.5 million hectares abroad—not bad going for an Institute that only started to specialise in 1959.

Maize in Yugoslavia is mainly fed to cattle. More than 87% of the corn is converted into meat and Yugoslavia is a meat exporter. However fluctuations in the price of meat are resulting in a reassessment of the use of maize and new developments from the United States such as the production of glucose/fructose syrup and the utilisation of the whole maize plant for other types of product are being actively investigated. The potential for greater use of maize seems to be there. Over the past seven years the average yield of maize in Yugoslavia has doubled as a result of the use of the Maize Institute's hybrids which are now grown on two thirds of the total land area cropped to maize. They claim now that they have more than 75 new hybrids which will yield about 10 tonnes of grain per hectare as opposed to the average in Yugoslavia of 4 tonnes per hectare.

Research at the Institute is divided between four sections. Breeding and Genetics naturally employs the largest number of scientists, many of whom have been trained in North America. Research in the Physiology, Biochemistry and Technology Section takes several basic approaches which may have long term application to genetics and maize production. Low temperature growth in maize is being studied in the hope of extending the growing season by earlier planting and of understanding the physiology of low temperature adaptation. Developmental physiology and adaptation to the various climates found in Yugoslavia are also under investigation. Protein biosynthesis is being studied with the aim of increasing the lysine and tryptophan content of the plant. Biophysical assays which can be used for selection during breeding programmes are also being developed. The third section, Maize Protection, deals with fungal and bacterial diseases, insect attack and weed problems. The fourth, Agronomy, is involved in the techniques of maize cultivation. The integration of basic research and seed production and commercial exploitation seems in the case of this Institute to have been particularly successful.

deposition is to be considered an interpreted property of unequal accuracy, dependent on lithology, intensity of remanence, post-depositional history of the layer, the demagnetisation programme and the instruments used.

In these circumstances, the validity of magnetostratigraphic zonation of sedimentary rocks is only proved by reproduction of the results in sequences involving different lithologies,

depositional environments and sedimentation rates. Results of highest confidence are those obtained on stratigraphic successions of effusive volcanic rocks where the danger of remagnetisation is minimal. Magnetostratigraphers are urged to duplicate sampling in parallel sections and in parallel cores wherever possible. Further effort is also needed in studies of rock magnetism dedicated to recog-

nition of original remanence from overprint and to improvement of our understanding of processes through which the remanence of sediments is acquired and destroyed.

Correlation of magnetozones with palaeomagnetic chronology

Magnetostratigraphy is not a dating technique. It only provides globally synchronous correlation planes, whose