10. South, T. L., Blake, P. R., Hare, D. R. & Summers, M. F. C-terminal retroviral-type zinc finger domain from the HIV-1 nucleocapsid protein is structurally similar to the N-terminal zinc finger domain. *Biochemistry* **30**, 6342–6349 (1991).
11. Luisi, B. F. *et al.* Crystallographic analysis of the interaction of the glucocorticoid receptor with DNA. *Nature* **352**, 497–505 (1991).
12. Marmorstein, R., Carey, M., Ptashne, M. & Harrison, S. C. DNA recognition by GAL4: structure of a protein–DNA complex. *Nature* **356**, 408–414 (1992).
13. Everett, R. D. *et al.* A novel arrangement of zinc-binding residues and secondary structure in the C3HC4 motif of an alpha herpes virus protein family. *J. Mol. Biol.* **234**, 1038–1047 (1993).
14. Barlow, P. N., Luisi, B., Milner, A., Elliot, M. & Everett, R. Structure of the C₃HC₄ domain by ¹H-nuclear magnetic resonance spectroscopy. *J. Mol. Biol.* **237**, 201–211 (1994).
15. Borden, K. L. B. *et al.* The solution structure of the RING finger domain from the acute promyelocytic leukaemia proto-oncoprotein PML. *EMBO J.* **14**, 1532–1541 (1995).
16. Phillips, S. E. V. The β-ribbon DNA recognition motif. *Ann. Rev. Biophys. Biomol. Struct.* **23**, 671–701 (1994).
17. Kim, J. L., Nikolov, D. B. & Burley, S. K. Co-crystal structure of TBP recognizing the minor groove of a TATA element. *Nature* **365**, 520–527 (1993).
18. Kim, Y., Geiger, J. H., Hahn, S. & Sigler, P. B. Crystal structure of a yeast TBP–TATA-box complex. *Nature* **365**, 512–520 (1993).
19. Schumacher, M. A., Choi, K. Y., Zalkin, H. & Brennan, R. G. Crystal structure of LacI member, PurR, bound to DNA: minor groove binding by α-helices. *Science* **266**, 763–770 (1994).
20. Flick, K. E. *et al.* Crystallization and preliminary X-ray studies of I-Ppol: a nuclear, intron-encoded homing endonuclease from *Physarum polycephalum*. *Protein Sci.* **6**, 1–4 (1997).
21. Otwinowski, Z. & Minor, W. Processing of X-ray diffraction data collected in oscillation mode. *Methods Enzymol.* **276**, 307–326 (1997).
22. Leslie, A. G. W. in *Joint CCP4 and ESF-EACMB Newsletter on Protein Crystallography* (Daresbury Laboratory, Warrington, UK, 1992).
23. CCP4 The SERC (UK) Collaborative Computing Project No. 4, a Suite of Programs for Protein Crystallography (Daresbury Laboratory, Warrington, UK, 1979).
24. QUANTA96 X-ray Structure Analysis User's Reference (Molecular Simulations, San Diego, 1996).
25. Brünger, A. *XPLOR version 3.1: A System for X-ray Crystallography and NMR* (Yale Univ. Press, New Haven, CT, 1992).
26. Laskowski, R. J., MacArthur, M. W., Moss, D. S. & Thornton, J. M. PROCHECK: a program to check the stereochemical quality of protein structures. *J. Appl. Crystallogr.* **26**, 383–290 (1993).
27. Evans, S. V. SETOR: hardware-lighted three-dimensional solid model representations of macromolecules. *J. Mol. Graphics* **11**, 134–138 (1993).

Acknowledgements. We thank D. McHugh, K. Stephens and J. D. Heath for initial subcloning, purification and crystallization studies; R. Strong, K. Zhang and B. Scott for advice during the crystallographic analysis; and the beamline staff at the Advanced Light Source (NLBL laboratories), beamline 5.0.2, particularly T. Earnest, for assistance. B.L.S. and R.J.M. are funded for this project by the NIH. K.E.F. was supported by a NIH training grant and the American Heart Association. M.S.J. was supported by an NSF fellowship and an NIH training grant.

Correspondence and requests for materials and coordinates should be addressed to B.L.S. (e-mail: bstoddar@fred.hcrcc.org). Coordinates have been deposited in the Brookhaven Protein Data Bank (accession nos 1pp, 1a73, 1a74).

corrections

Emergence of symbiosis in peptide self-replication through a hypercyclic network

David H. Lee, Kay Severin, Yohei Yokobayashi & M. Reza Ghadiri

Nature **390**, 591–594 (1997)

Hypercycles are based on second-order (or higher) autocatalysis and defined by two or more replicators that are connected by

another superimposed autocatalytic cycle. Our study describes a mutualistic relationship between two replicators, each catalysing the formation of the other, that are linked by a superimposed catalytic cycle. Although the kinetic data suggest the intermediary of higher-order species in the autocatalytic processes, the present system should not be referred to as an example of a minimal hypercycle in the absence of direct experimental evidence for the autocatalytic cross-coupling between replicators. □

The complete genome sequence of the hyperthermophilic, sulphate-reducing archaeon *Archaeoglobus fulgidus*

Hans-Peter Klenk, Rebecca A. Clayton, Jean-Francois Tomb, Owen White, Karen E. Nelson, Karen A. Ketchum, Robert J. Dodson, Michelle Gwinn, Erin K. Hickey, Jeremy D. Peterson, Delwood L. Richardson, Anthony R. Kerlavage, David E. Graham, Nikos C. Kyrpides, Robert D. Fleischmann, John Quackenbush, Norman H. Lee, Granger G. Sutton, Steven Gill, Ewen F. Kirkness, Brian A. Dougherty, Keith McKenney, Mark D. Adams, Brendan Loftus, Scott Peterson, Claudia I. Reich, Leslie K. McNeil, Jonathan H. Badger, Anna Glodek, Lixin Zhou, Ross Overbeek, Jeannine D. Gocayne, Janice F. Weidman, Lisa McDonald, Teresa Utterback, Matthew D. Cotton, Tracy Spriggs, Patricia Artiach, Brian P. Kaine, Sean M. Sykes, Paul W. Sadow, Kurt P. D'Andrea, Cheryl Bowman, Claire Fujii, Stacey A. Garland, Tanya M. Mason, Gary J. Olsen, Claire M. Fraser, Hamilton O. Smith, Carl R. Woese & J. Craig Venter

Nature **390**, 364–370 (1997)

The pathway for sulphate reduction is incorrect as published: in Fig. 3 on page 367, adenylyl sulphate 3-phosphotransferase (*cysC*) is not needed in the pathway as outlined, as adenylyl sulphate reductase (*aprAB*) catalyses the first step in the reduction of adenylyl sulphate. The correct sequence of reactions is: sulphate is first activated to adenylyl sulphate, then reduced to sulphite and subsequently to sulphide. The enzymes catalysing these reactions are: sulphate adenylyltransferase (*sat*), adenylylsulphate reductase (*aprAB*), and sulphite reductase (*dsrABD*). We thank Jens-Dirk Schwenn for bringing this error to our attention. □

(that is, on its left side in Fig. 2a and b), a 231-nm-thick $\text{Al}_{0.165}\text{Ga}_{0.835}\text{As}$ spacer layer was grown with two Si δ -doping layers ($1 \times 10^{12} \text{ cm}^{-2}$), one inserted 22 nm and the other 187 nm from the left edge of the deep well. A 10-nm-thick undoped GaAs region capped the structure. The spacer layer thickness was adjusted to preserve the same distance between the δ -doping layers and the double-quantum-well system, and therefore the electrostatic potentials are identical in both structures. The δ -doping provides a two-dimensional electron gas in the deep well with a calculated sheet electron density of $n_s = 4 \times 10^{11} \text{ cm}^{-2}$.

For the absorption measurements, we processed our samples in a multipass (six) 45° wedge waveguide. This geometry allowed us to couple in linearly polarized radiation with a large component of the polarization normal to the layer (50%) as required by the intersub-band absorption selection rule⁹. The absorption was measured with a Fourier-transform infrared spectrometer (FTIR) using a step-scan modulation technique¹⁰ in which the electron gas in the double well is periodically depopulated by a Ti/Au Schottky barrier contact evaporated on the surface of the sample and the two-dimensional electron gas is contacted by indium balls alloyed into the layer.

The absorption measurements at $T = 10 \text{ K}$ for both structures are compared in Fig. 3 with the results of numerical calculations using the coupled Schrödinger's and Poisson's equations. As predicted, the absorption strength at photon energies between the two resonances is strongly suppressed or enhanced by the interference effect depending on the location of the thin barrier, proving that tunnelling through the latter controls the interference effect when the broadening of the states is dominated by tunnelling. However, the finite broadening introduced by interface disorder prevents full quantum interference; this is the main reason for the departure from the calculated profiles and specifically the reason why the absorption does not vanish in the sample with destructive interference. Indeed, linewidth measurements on samples with the same coupled-well structure but with negligible tunnelling to the continuum showed a full-width at half-maximum of the absorption peaks of $\Gamma = 5 \text{ meV}$. This structure consists of an identical double quantum well between two 60-nm-thick $\text{Al}_{0.33}\text{Ga}_{0.67}\text{As}$ barriers. This value is a measure of the non-tunnelling contribution to the broadening of the optical transitions; it is smaller but not negligible compared with the calculated broadening by tunnelling through the 1.5 nm barrier, $\Gamma_1 \cong \Gamma_2 \cong 16 \text{ meV}$.

Destructive interference in intersub-band absorption in a double-well structure coupled by tunnelling to a continuum has recently been inferred from a fit of the absorption lineshape to a model that included the collision broadening in a phenomenological manner¹¹. The present experiment gives more direct evidence of tunnelling-induced quantum interference by showing that tunnelling can be used to control the sign of the interference.

It is important to stress the difference between the phenomena described here and the Fano interference in intersub-band absorption recently reported by us¹². In that work a minimum in the absorption arises because of interference between matrix elements for the ground state to the continuum and to a single resonance coupled by tunnelling to the same continuum. This leads to a strongly asymmetric absorption lineshape. In contrast, in the phenomena studied here interference arises between absorption paths through two resonances coupled to a continuum, and the direct matrix element from the ground state to the continuum is negligible.

These findings are relevant for the design of semiconductor lasers without population inversion (LWI). Such lasing action has so far been observed only in gases^{4,5}. Essential for LWI is nonreciprocity between emission and absorption. A possible semiconductor LWI scheme would use the quantum-well structure of Fig. 2a for the active regions. The latter would be alternated with electron injectors as in quantum cascade lasers¹³. Electrons would be injected from the thick barrier side at an energy between the two resonances where the

absorption cross-section is a minimum, to ensure strong non-reciprocity between intersub-band absorption and emission^{7,8}. Although the realization of such a laser would be scientifically important, its implementation would be difficult and its technological impact limited by the very short lifetime (a few tenths of picoseconds) of the excited state which is required to achieve strong interference¹⁴. □

Received 23 June; accepted 6 October 1997.

1. Fano, U. Effect of configuration interaction on intensity and phase shifts. *Phys. Rev.* **124**, 1866–1878 (1961).
2. Harris, S. E. Lasers without inversion: interference of lifetime-broadened resonance. *Phys. Rev. Lett.* **62**, 1033–1035 (1989).
3. Arkhipkin, V. G. & Heller, Y. I. Radiation amplification without population at transitions to autoionizing states. *Phys. Lett. A* **98**, 12–14 (1983).
4. Zibrov, S. *et al.* Experimental demonstration of laser without population inversion via quantum interference in Rb. *Phys. Rev. Lett.* **75**, 1499–1452 (1995).
5. Padmadandu, G. G. *et al.* Laser oscillation without population inversion in a Sodium atomic beam. *Phys. Rev. Lett.* **76**, 2053–2056 (1996).
6. Boller, K.-J., Imamoglu, A. & Harris, S. E. Observation of electromagnetically induced transparency. *Phys. Rev. Lett.* **66**, 2593–2596 (1991).
7. Imamoglu, A. & Harris, S. E. Lasers without inversion: interference of dressed lifetime-broadened states. *Opt. Lett.* **13**, 1344–1346 (1989).
8. Imamoglu, A. & Ram, R. J. Semiconductor lasers without population inversion. *Opt. Lett.* **19**, 1744–1746 (1994).
9. Levine, B. F., Choi, K. K., Bethea, C. G., Walker, J. & Malik, R. J. New 10 μm infrared detector using intersubband absorption in resonant tunneling GaAlAs superlattices. *Appl. Phys. Lett.* **50**, 1092–1094 (1987).
10. Faist, J. *et al.* Narrowing of the intersubband electroluminescence spectrum in coupled-quantum-well heterostructures. *Appl. Phys. Lett.* **65**, 94–96 (1994).
11. Smidt, H., Campman, K. L., Gossard, A. C. & Imamoglu, A. Tunnel induced transparency: Fano interference in intersubband transitions. *Appl. Phys. Lett.* **70**, 3455–3457 (1997).
12. Faist, J. *et al.* Tunable Fano interference in intersubband absorption. *Opt. Lett.* **21**, 985–987 (1996).
13. Capasso, F., Faist, J., Sirtori, C. & Cho, A. Y. Infrared (4–11 μm) quantum cascade lasers. *Solid State Comm.* **102**, 231–236 (1997).
14. Kurghin, J. B. & Rosencher, E. Practical aspects of lasing without inversion in various media. *IEEE J. Quant. Electron.* **QE-32**, 1882–1890 (1996).

Acknowledgements. We thank S. Harris, A. Imamoglu, R. J. Ram, A. Tredicucci and C. Gmachl for enlightening discussions, and the authors of ref. 11 for making their manuscript available before publication. The work was supported in part by DARPA/ARO.

Correspondence should be addressed to F.C. (e-mail: fc@bell-labs.com or fc@lucent.com).

Emergence of symbiosis in peptide self-replication through a hypercyclic network

David H. Lee, Kay Severin, Yohei Yokobayashi & M. Reza Ghadiri

Departments of Chemistry and Molecular Biology, and the Skaggs Institute for Chemical Biology, The Scripps Research Institute, La Jolla, California 92037, USA

Symbiosis is an association between different organisms that leads to a reciprocal enhancement of their ability to survive. Similar mutually beneficial relationships can operate at the molecular level in the form of a hypercycle, a collective of two or more self-replicating species interlinked through a cyclic catalytic network^{1–5}. The superposition of cross-catalysis onto autocatalytic replication integrates the members of the hypercycle into a single system that reproduces through a second-order (or higher) form of nonlinear autocatalysis. The hypercycle population as a whole is therefore able to compete more efficiently for existing resources than any one member on its own. In addition, the effects of beneficial mutations of any one member are spread over the entire population. The formation of hypercycles has been suggested as an important step in the transition from inanimate to living chemistry⁶, and a large number of hypercycles are expected to be embedded within the complex networks of living systems⁷. But only one naturally occurring hypercycle has been well documented⁸, while two autocatalytic chemical systems may contain vestiges of hypercyclic organization^{9,10}. Here we report a

chemical system that constitutes a clear example of a minimal hypercyclic network, in which two otherwise competitive self-replicating peptides symbiotically catalyse each others' production.

The present design of a minimal hypercycle is based on two self-replicating coiled coil peptides R_1 and R_2 (Fig. 1). The replicator R_1 was recently reported^{11,12} and is produced as the ligation product of the electrophilic peptide fragment E and the nucleophilic fragment N_1 . The replicator R_2 is made from the same electrophilic fragment but a different nucleophilic peptide fragment N_2 . The nucleophilic fragments N_1 and N_2 differ in their sequence at the hydrophobic recognition surface— N_1 is composed of valine and leucine whereas N_2 is made up of isoleucine and leucine residues. This difference in sequence at the hydrophobic core is known to affect profoundly the aggregation state of coiled coils^{13,14}. Furthermore it is known that conservative mutations in this region of the structure can drastically alter the kinetic behaviour of the replicator^{11,12,15}.

The ability of R_2 to self-replicate was determined by observation of characteristics previously established as signatures of self-replication (Fig. 2)^{11,12}. Similar to that of R_1 , the new replicator R_2 also displays a parabolic growth profile. Numerical fitting of the kinetic data obtained for R_2 to the empirical rate equations of von Kiedrowski¹⁶ gave a background rate constant $k_b = 0.072 \pm 0.005 \text{ M}^{-1} \text{ s}^{-1}$ and an apparent autocatalytic rate constant $k_a = 52 \pm 1 \text{ M}^{-3/2} \text{ s}^{-1}$, making R_2 more efficient than its relative R_1 ($k_b = 0.063 \text{ M}^{-1} \text{ s}^{-1}$ and constant $k_a = 29.4 \text{ M}^{-3/2} \text{ s}^{-1}$).

A solution containing all three fragments E, N_1 and N_2 gave a combinatorial synthesis of both replicators. *A priori*, one would

expect a survival-of-the-fittest situation where the more efficient replicator R_2 would overwhelm R_1 by consuming the common fragment E more quickly. At first glance, this expectation seemed to be borne out as R_2 was produced in greater abundance than R_1 (as expected, when molecular interactions are disrupted in the presence of guanidinium hydrochloride, no kinetic preference for R_2 over R_1 was observed). However, the situation is more interesting and complex. When we sought to give R_1 an advantage in this competition by adding 40% R_1 (with respect to the nucleophile concentration) at the start of the reaction, to our surprise the rate of R_1 self-production increased by only 1.7 times over the unseeded reaction but the rate of R_2 formation was enhanced to a greater extent, by 5.4 times (Table 1, Fig. 3). Thus the two replicators are not mutually exclusive in their growth; R_1 catalyses the formation of R_2 as well as itself. Likewise, perturbation of the reaction by seeding it with 45% R_2 not only increased the rate of R_2 production 2.9 times but R_1 as well, by 3.5 times. Thus a cross-catalytic cycle is cooperatively coupled with two self-replicating reactions, making this system one which is hypercyclic in nature. There are four characteristic outcomes expected for such a hypercyclic network, depending on the relative efficiencies of the coupled catalytic and autocatalytic reactions². The observed greater efficiencies of the catalytic reactions over the autocatalytic components of the system are the most desirable outcomes which assure the stability of the hypercycle: production of one species promotes the production of the other to an even greater degree. This particular mode of catalytic coupling prevents one replicator from overwhelming the other and enables the two to reproduce as a single coherent unit.

To verify that R_1 and R_2 catalyse each other's production, the

Figure 1 Schematic diagram of a minimal hypercycle based on two self-replicating peptides. Cycles I and III show the self-producing cycles of replicators R_1 (dark grey/light grey) and R_2 (dark grey/striped) respectively, which pre-organize their constituent fragments thereby promoting peptide ligation. Cycle II, where R_1 promotes R_2 formation, and cycle IV, where R_2 promotes R_1 formation, comprise the catalytic components of the hypercycle and allow the replicators to positively regulate each others' production. The mechanistic details of the present hypercyclic network may be more complex than the minimal system depicted here. Detailed kinetic analyses of the replicator sequences have shown that the autocatalytically productive intermediates involve, at least in part, quaternary complexes in which two template strands pre-organize the reactive peptide fragments (ref. 12 and K. Kumar, D.H.L., M.R.G., unpublished results). The following peptide sequences were employed in this study: replicator 1 (R_1), ArCONH-RMKQLEEKVYELLKSKVA-CLEXEVARLKKLVGE-CONH₂; replicator 2 (R_2), ArCONH-RMKQLEEKVYELLKSKVA-CLEXEVARLKKLVGE-CONH₂; electrophilic fragment (E), ArCONH-RMKQLEEKVYELLKSKVA-COSBn; nucleophilic fragment 1 (N_1), H₂N-CLEXEVARLKKLVGE-CONH₂; nucleophilic fragment 2 (N_2), H₂N-CLEXEVARLKKLVGE-CONH₂. Bn, benzyl; Ar, 4-acetamidophenyl; and X, lysine-ε-NHCO-Ar.

Table 1 Initial rates of product formation

Product	No replicators added	+40% R_1	+45% R_2
R_1	4.8	8.2	17.0
R_2	5.8	31.1	16.9

The data in this table (in units of $10^{-6} \text{ M min}^{-1}$) are for reactions containing the three peptide fragments in the absence and presence of added replicators.

Figure 2 Production of R_2 as a function of time in the presence of various initial concentrations of R_2 . Open circles, in the absence of any added R_2 ; filled diamonds, in the presence of 4.0 μM ; open triangles, in the presence of 21.4 μM ; and filled circles, 42.6 μM of initially added R_2 . Curves were generated by nonlinear least-squares fit of the data to the empirical rate equation of von Kiedrowski using the program SimFit¹⁶. Data are an average of two experiments.

reaction mixtures were simplified to include **E** and only one nucleophile, and then seeded with the template that was not produced *in situ* (Fig. 4). Comparisons with unseeded reactions revealed that even in these simplified systems one template can promote the formation of the other, giving rate enhancements much larger than what would be expected if the reaction mixture were seeded with the autocatalytic template. Reaction mixtures containing **E** and **N**₁ that were seeded with 25% **R**₂ enhanced the initial rate of production of **R**₁ from $3.9 \times 10^{-8} \text{ M min}^{-1}$ to $1.5 \times 10^{-7} \text{ M min}^{-1}$, a 3.8 times increase over the unseeded reaction. Seeding of the same reaction mixture with 25% **R**₁ would improve the rate by only 2.8 times. Similarly, seeding reaction mixtures containing **E** and **N**₂ with 35% **R**₁ gave a 5.4 times rate enhancement over the $5.0 \times 10^{-8} \text{ M min}^{-1}$ rate observed for the reaction without added catalyst. The increase is greater than the 3.6

times enhancement expected for the autocatalytic reaction containing 35% **R**₂.

We now consider the sequence selectivity issues in the formation of the hypercyclic peptide network. The operation of the hypercycle is based on complementary, as well as self-complementary, forms of catalysis. As noted below, there is mounting evidence that both processes are strongly sequence selective. Previously we had shown that in the case of replicator **R**₁, even conservative mutations (Val9Ala—where a valine has been substituted by an alanine at position 9—and Leu26Ala) in the hydrophobic core residues completely abolish the autocatalytic process^{11–12}. In this study we have determined that similar replicator **R**₂ mutations are also autocatalytically infertile. There is also good evidence for high sequence selectivity in the cross-catalytic component of the system. Control studies have indicated that the Leu26Ala **R**₂ mutant cannot cross-catalyse the formation of replicators **R**₁ nor **R**₂. Although in a recent study¹⁵ we have shown that the Val9Ala **R**₁ mutant can efficiently cross-catalyse the formation of **R**₁, we have found it to be ineffective in catalysing **R**₂ production. Moreover, in a related study we have shown that diminution in the initial rate of peptide fragment condensation of more than 3 orders of magnitude can be caused even by electrostatic substitutions at the solvent-exposed *e* and *g* positions of the heptad repeat sequence¹⁷. Although the above studies strongly support high sequence selectivity in the catalytic and autocatalytic components of the hypercyclic network, a significantly large sequence-space must undoubtedly exist that would enable the spontaneous self-organization of even more complex networks. Studies along those lines are under investigation.

The work reported here may have particular relevance to various origin-of-life theories^{1–4,18}. It has been suggested that at the dawn of life the onset of darwinian evolution must have been marked by

Figure 3 Replicators **R**₁ and **R**₂ self-organize into a two-membered hypercyclic network. **a**, Production of **R**₁ (empty circles) and **R**₂ (empty diamonds) as a function of time for reaction mixtures containing **E**, **N**₁ and **N**₂. **b**, Formation of **R**₁ (filled circles) and **R**₂ (filled diamonds) as a function of time for reaction mixtures containing the three fragments and 40% **R**₁. **c**, Formation of **R**₁ (filled circles) and **R**₂ (filled diamonds) as a function of time for reaction mixtures containing the three fragments and 45% **R**₂. In **b** and **c**, production formation in the absence of added templates are shown for comparison. Data are an average of two experiments. Curves are shown to guide the eye.

Figure 4 Replicators **R**₁ and **R**₂ are cross-catalytic. **a**, Formation of **R**₁ as a function of time for the reaction mixture containing only **E** and **N**₁ in the absence (empty circles) and in the presence (filled circles) of 35% **R**₂. **b**, Formation of **R**₂ as a function of time for the reaction mixture containing **E** and **N**₂ in the absence (empty diamonds) and in the presence (filled diamonds) of 25% **R**₁. Data are an average of two experiments. Curves are shown to guide the eye.

selection based on feedback processes of genotype replication¹⁹. It is also likely that molecular genotypes and phenotypes may have been the very same molecules²⁰. Our example of a hypercyclic peptide network supports the idea that peptides could play a role in both hypotheses. □

Methods

Self-replication of R₂. All reactions were done in 0.6 ml Eppendorf tubes at 23 °C. A stock solution containing E, N₂ and the internal standard 4-acetamidobenzoic acid (ABA), were seeded with various amounts of R₂. Benzylmercaptan (1 µl) was then added. Reactions were initiated by adding 3-(N-morpholino)propanesulphonic acid (MOPS) buffer (pH = 7.50, 200 mM, 236 µl), giving a total volume of 300 µl and concentrations of [N₂] = 104.5 µM, [E] = 94.2 µM, [R₂] = 0, 4.0, 21.4 or 42.6 µM. [MOPS] = 157 mM, [ABA] = 40.4 µM. Samples (30 µl) were taken at various time points and quenched with 2% trifluoroacetic acid (TFA) in water (70 µl) then stored at -70 °C. Samples were analysed by high pressure liquid chromatography on a Zorbax C8 column using an acetonitrile/water/0.1% TFA gradient while monitoring at 270 nm. The identity of all peptides was determined by mass spectrometry and verified by coinjection with authentic samples. Experiments were done in duplicate.

Determination of hypercyclic organization in the E/N₁/N₂ mixture. Reactions were done as described above except that the stock solution contained, besides E and ABA, both N₁ and N₂, which was subsequently seeded with either R₁, R₂, or water. Reactions were initiated by adding MOPS buffer (pH = 7.50, 200 mM, 236.6 µl), giving a total volume of 300 µl and concentrations of [N₁] = 112.5 µM, [N₂] = 112.7 µM, [E] = 91.1 µM, [MOPS] = 157.7 mM, [ABA] = 97.1 µM, [R₁] = 45.1 µM, [R₂] = 50.4 µM.

Verification of the catalytic components of the hypercycle. Reactions were performed as described above except only one nucleophile was present in the reaction mixture and the reaction was seeded with the replicator that was not produced *in situ*. Initial concentrations are (1) [E] = 88.9 µM, [N₁] = 98.2 µM, [R₂] = 25.2 µM, [ABA] = 50.5 µM; (2) [E] = 80.4 µM, [N₂₁] = 96.9 µM, [R₁] = 35.3 µM, [ABA] = 36.9 µM.

Received 20 June; accepted 20 October 1997.

1. Eigen, M. & Schuster, P. The hypercycle. A principle of natural self-organization. Part A: emergence of the hypercycle. *Naturwissenschaften* **64**, 541–565 (1977).
2. Eigen, M. & Schuster, P. The hypercycle. A principle of natural self-organization. Part B: the abstract hypercycle. *Naturwissenschaften* **65**, 7–41 (1978).
3. Eigen, M. & Schuster, P. The hypercycle. A principle of natural self-organization. Part C: The realistic hypercycle. *Naturwissenschaften* **65**, 341–369 (1978).
4. Eigen, M. Self-organization of matter and the evolution of biological macromolecules. *Naturwissenschaften* **58**, 465–523 (1971).
5. Müller-Herold, U. What is a hypercycle? *J. Theor. Biol.* **102**, 569–584 (1983).
6. Lee, D. H., Severin, K. & Ghadiri, M. R. Autocatalytic networks: the transition from molecular self-replication to molecular ecosystems. *Curr. Opin. Chem. Biol.* (in the press).
7. Ricard, J. & Noat, G. Electrostatic effects and the dynamics of enzyme reactions at the surface of plant cells. I. A theory of the ionic control of a complex multi-enzyme system. *Eur. J. Biochem.* **155**, 183–190 (1986).
8. Eigen, M., Biebricher, C. K., Gebinoga, M. & Gardiner, W. C. The hypercycle. Coupling of RNA and protein biosynthesis in the infection cycle of an RNA bacteriophage. *Biochemistry* **30**, 11005–11018 (1991).
9. Hong, J.-I., Feng, Q., Rotello, V. & Rebek, J. Jr Competition, cooperation and mutation: improving a synthetic replicator by light irradiation. *Science* **255**, 848–850 (1992).
10. Achilles, T. & von Kiedrowski, G. A self-replicating system from three starting materials. *Angew. Chem. Int. Edn Engl.* **32**, 1198–1201 (1993).
11. Lee, D. H., Granja, J. R., Martinez, J. A., Severin, K. S. & Ghadiri, M. R. A self-replicating peptide. *Nature* **382**, 525–528 (1996).
12. Severin, K. S., Lee, D. H., Martinez, J. A. & Ghadiri, M. R. Peptide self-replication via template-directed ligation. *Chem. Eur. J.* **3**, 1017–1024 (1997).
13. Harbury, P. B., Zhang, T., Kim, P. S. & Alber, T. A switch between two-, three- and four-stranded coiled coils in GCN4 leucine zipper mutants. *Science* **262**, 1401–1407 (1993).
14. Hu, J. C., O'Shea, E. K., Kim, P. S. & Sauer, R. T. Sequence requirements for coiled coils: analysis with λ repressor-GCN4 leucine zipper fusions. *Science* **250**, 1400–1403 (1990).
15. Severin, K., Lee, D. H., Martinez, J. A. & Ghadiri, M. R. Dynamic error-correction in an autocatalytic peptide network. *Angew. Chem. Int. Edn Engl.* (in the press).
16. von Kiedrowski, G. Minimal replicator theory I: parabolic versus exponential growth. *Bioorg. Chem. Front.* **3**, 113–146 (1993).
17. Severin, K., Lee, D. H., Kennan, A. J. & Ghadiri, M. R. A synthetic peptide ligase. *Nature* **389**, 706–709 (1997).
18. Kauffman, S. A. *The Origins of Order* (Oxford Univ. Press, New York, 1993).
19. Küppers, B.-O. *The Origin of Biological Information* (MIT Press, Cambridge, MA, 1990).
20. Joyce, G. F. RNA evolution and the origins of life. *Nature* **338**, 217–224 (1989).

Acknowledgements. We thank K. Kumar for discussions. We also thank the Medical Research Council of Canada for a predoctoral fellowship (D.H.L.), and the Deutsche Forschungsgemeinschaft for a postdoctoral fellowship (K.S.).

Correspondence and requests for materials should be addressed to M.R.G. (e-mail: ghadiri@scripps.edu).

Kinetic limitations on droplet formation in clouds

P. Y. Chuang*, R. J. Charlson† & J. H. Seinfeld‡

* Department of Environmental Engineering Science, MC 138-78, California Institute of Technology, Pasadena, California 91125, USA

† Departments of Atmospheric Sciences and Chemistry, University of Washington, Seattle, Washington 98195, USA

‡ Division of Engineering and Applied Science and Department of Chemical Engineering, MC 210-41, California Institute of Technology, Pasadena, California 91125, USA

The 'indirect' radiative cooling of climate due to the role of anthropogenic aerosols in cloud droplet formation processes (which affect cloud albedo) is potentially large, up to -1.5 W m^{-2} (ref. 1). It is important to be able to determine the number concentration of cloud droplets to within a few per cent, as radiative forcing as a result of clouds is very sensitive to changes in this quantity², but empirical approaches are problematic^{3–5}. The initial growth of a subset of particles known as cloud condensation nuclei and their subsequent 'activation' to form droplets are generally calculated with the assumption that cloud droplet activation occurs as an equilibrium process described by classical Köhler theory^{6,7}. Here we show that this assumption can be invalid under certain realistic conditions. We conclude that the poor empirical correlation between cloud droplet and cloud condensation nuclei concentrations is partly a result of kinetically limited growth before droplet activation occurs. Ignoring these considerations in calculations of total cloud radiative forcing based on cloud condensation nuclei concentrations could lead to errors that are of the same order of magnitude as the total anthropogenic greenhouse-gas radiative forcing¹.

Cloud droplet activation and subsequent treatments of cloud droplet growth in atmospheric models generally rely on the assumption that pre-activation growth is accurately described by an equilibrium model in which the particle diameter is always at equilibrium with the local supersaturation^{6,7}. The equilibrium relationship between supersaturation and particle size for a particle composed of highly soluble inorganic species can be described by the well-known Köhler equation (curve A, Fig. 1)⁸. Cloud droplet nuclei (CDN) activate when they grow larger than their critical diameter, D_{pc} , after which they can grow spontaneously, limited only by growth kinetics. The concept of CDN is distinct from that of CCN in that, whereas CCN are defined as those particles that activate to become cloud droplets within a cloud chamber of fixed or prescribed supersaturation, CDN are those particles that actually activate in the atmosphere under conditions of time-varying supersaturation.

To evaluate the conditions under which the equilibrium activation model is valid, two timescales will be defined. One is the timescale for particle growth that would be required for that particle to remain at equilibrium as the ambient supersaturation ratio increases in a rising air parcel, τ_e . The other is the timescale for actual change in the droplet size resulting from condensational growth, τ_g . Hence, if $\tau_e \gg \tau_g$ then the equilibrium model is reasonable; otherwise, CDN activation, and hence the cloud droplet size distribution, can be accurately predicted only if the kinetics of droplet growth are considered. To calculate τ_e , the rate of change of the droplet diameter that would be required for that droplet to remain at its equilibrium size, dD_{pe}/dt , is determined from the combination of two effects. First, the time rate of change of supersaturation, dS/dt , can be determined using a simple one-dimensional adiabatic parcel model⁹. Next, the rate of change of D_{pe} with respect to supersaturation, dD_{pe}/dS , is determined by differentiating

The complete genome sequence of the hyperthermophilic, sulphate-reducing archaeon *Archaeoglobus fulgidus*

Hans-Peter Klenk*, Rebecca A. Clayton*, Jean-Francois Tomb*, Owen White*, Karen E. Nelson*, Karen A. Ketchum*, Robert J. Dodson*, Michelle Gwinn*, Erin K. Hickey*, Jeremy D. Peterson*, Delwood L. Richardson*, Anthony R. Kerlavage*, David E. Graham†, Nikos C. Kyrpides†, Robert D. Fleischmann*, John Quackenbush*, Norman H. Lee*, Granger G. Sutton*, Steven Gill*, Ewen F. Kirkness*, Brian A. Dougherty*, Keith McKenney*, Mark D. Adams*, Brendan Loftus*, Scott Peterson*, Claudia I. Reich†, Leslie K. McNeil†, Jonathan H. Badger†, Anna Glodek*, Lixin Zhou*, Ross Overbeek‡, Jeannine D. Gocayne*, Janice F. Weidman*, Lisa McDonald*, Teresa Utterback*, Matthew D. Cotton*, Tracy Spriggs*, Patricia Artiach*, Brian P. Kaine†, Sean M. Sykes*, Paul W. Sadow*, Kurt P. D'Andrea*, Cheryl Bowman*, Claire Fujii*, Stacey A. Garland*, Tanya M. Mason*, Gary J. Olsen†, Claire M. Fraser*, Hamilton O. Smith*, Carl R. Woese† & J. Craig Venter*

* The Institute for Genomic Research (TIGR), Rockville, Maryland 20850, USA

† Department of Microbiology, University of Illinois, Champaign-Urbana, Illinois 61801, USA

‡ Mathematics and Computer Science Division, Argonne National Laboratory, Illinois 60439, USA

***Archaeoglobus fulgidus* is the first sulphur-metabolizing organism to have its genome sequence determined. Its genome of 2,178,400 base pairs contains 2,436 open reading frames (ORFs). The information processing systems and the biosynthetic pathways for essential components (nucleotides, amino acids and cofactors) have extensive correlation with their counterparts in the archaeon *Methanococcus jannaschii*. The genomes of these two Archaea indicate dramatic differences in the way these organisms sense their environment, perform regulatory and transport functions, and gain energy. In contrast to *M. jannaschii*, *A. fulgidus* has fewer restriction-modification systems, and none of its genes appears to contain inteins. A quarter (651 ORFs) of the *A. fulgidus* genome encodes functionally uncharacterized yet conserved proteins, two-thirds of which are shared with *M. jannaschii* (428 ORFs). Another quarter of the genome encodes new proteins indicating substantial archaeal gene diversity.**

Biological sulphate reduction is part of the global sulphur cycle, ubiquitous in the earth's anaerobic environments, and is essential to the basal workings of the biosphere. Growth by sulphate reduction is restricted to relatively few groups of prokaryotes; all but one of these are Eubacteria, the exception being the archaeal sulphate reducers in the Archaeoglobales^{1,2}. These organisms are unique in that they are unrelated to other sulphate reducers, and because they grow at extremely high temperatures³. The known Archaeoglobales are strict anaerobes, most of which are hyperthermophilic marine sulphate reducers found in hydrothermal environments^{2,4} and in subsurface oil fields⁵. High-temperature sulphate reduction by *Archaeoglobus* species contributes to deep subsurface oil-well 'souring' by producing iron sulphide, which causes corrosion of iron and steel in oil- and gas-processing systems⁵.

Archaeoglobus fulgidus VC-16 (refs 2, 4) is the type strain of the Archaeoglobales. Cells are irregular spheres with a glycoprotein envelope and monopolar flagella. Growth occurs between 60 and 95 °C, with optimum growth at 83 °C and a minimum division time of 4 h. The organism grows organoheterotrophically using a variety of carbon and energy sources, but can grow lithoautotrophically on hydrogen, thiosulphate and carbon dioxide⁶. We sequenced the genome of *A. fulgidus* strain VC-16 as an example of a sulphur-metabolizing organism and to gain further insight into the Archaea^{7,8} through genomic comparison with *Methanococcus jannaschii*⁹.

General features of the genome

The genome of *A. fulgidus* consists of a single, circular chromosome of 2,178,400 base pairs (bp) with an average of 48.5% G+C content

(Fig. 1). There are three regions with low G+C content (<39%), two rich in genes encoding enzymes for lipopolysaccharide (LPS) biosynthesis, and two regions of high G+C content (>53%), containing genes for large ribosomal RNAs, proteins involved in haem biosynthesis (*hemAB*), and several transporters (Table 1). Because the origins of replication in Archaea are not characterized, we arbitrarily designated base pair one within a presumed non-coding region upstream of one of three areas containing multiple short repeat elements.

Open reading frames. Two independent coding analysis programs and BLASTX¹⁰ searches (see Methods) predicted 2,436 ORFs (Figs 1, 2, Tables 1, 2) covering 92.2% of the genome. The average size of the *A. fulgidus* ORFs is 822 bp, similar to that of *M. jannaschii* (856 bp), but smaller than that in the completely sequenced eubacterial genomes (949 bp). All ORFs were searched against a non-redundant protein database, resulting in 1,797 putative identifications that were assigned biological roles within a classification system adapted from ref. 11. Predicted start codons are 76% ATG, 22% GTG and 2% TTG. Unlike *M. jannaschii*, where 18 inteins were found in coding regions, no inteins were identified in *A. fulgidus*. Compared with *M. jannaschii*, *A. fulgidus* contains a large number of gene duplications, contributing to its larger genome size. The average protein relative molecular mass (M_r) in *A. fulgidus* is 29,753, ranging from 1,939 to 266,571, similar to that observed in other prokaryotes. The isoelectric point (pI) of predicted proteins among sequenced prokaryotes exhibits a bimodal distribution with peaks at pIs of approximately 5.5 and 10.5. The exceptions to this are *Mycoplasma genitalium* in which the distribution is skewed towards high pI

Figure 1 Circular representation of the *A. fulgidus* genome. The outer circle shows predicted protein-coding regions on the plus strand classified by function according to the colour code in Fig. 2 (except for unknowns and hypotheticals, which are in black). Second circle shows predicted protein-coding regions on the minus strand. Third and fourth circles show IS elements (red) and other repeats (green) on the plus and minus strand. Fifth and sixth circles show tRNAs (blue), rRNAs (red) and sRNAs (green) on the plus and minus strand, respectively.

Table 1 Genome features

General

Chromosome size:	2,178,400 bp
Protein coding regions:	92.2%
Stable RNAs:	0.4%

Predicted protein coding sequences:

Identified by database match:	2,436 (1.1 per kb)
putative function assigned:	1,797
homologues of <i>M. jannaschii</i> ORFs:	1,096
conserved hypothetical proteins:	916
No database match:	651
Members of 242 paralogous families:	639
Members of 158 families with known functions:	719
	475

Stable RNAs

16S rRNA:	<i>Coordinates</i>
23S rRNA:	1,790,478–1,788,987
5S rRNA:	1,788,751–1,785,820
7S RNA:	81,144–81,021
RNase P:	798,067–798,376
46 species of tRNA:	86,281–86,032
tRNAs with 15–62 bp introns:	no significant clusters
	Asp ^{GUC} , Glu ^{UUC} , Leu ^{CAA} , Trp ^{CCA} , Tyr ^{GUA}

Distinct G+C content regions

HGC-1, >53% G+C	<i>Coordinates</i>
HGC-2, >53% G+C	1,786,000–1,797,000
LGC-1, <39% G+C	2,158,000–2,159,000
LGC-2, <39% G+C	281,000–284,000
LGC-3, <39% G+C	544,000–550,000
	1,175,000–1,177,000

Short, non-coding repeats

SR-1A, CTTTCAATCCCATTTTGGTCTGATTCAAC	<i>Coordinates</i>
SR-1B, CTTTCAATCCCATTTTGGTCTGATTCAAC	147–4,213
SR-2, CTTTCAATCTCCATTTTCAGGGCCTCCCTTTCTTA	398,368–401,590
	1,690,930–1,694,104

Long, coding repeats

	<i>Length</i>	<i>Copy number</i>
LR-01 NADH-flavin oxidoreductase	1,886 bp	2 copies
LR-02 NifS, NifU + ORF	1,549 bp	2 copies
LR-03 ISA1214 putative transposase + ISORF2	1,214 bp	6 copies
LR-04 ISA1083 putative transposase + ISORF2	1,083 bp	3 copies
LR-05 type II secretion system protein	1,014 bp	4 copies
LR-06 ISA0963 putative transposase	963 bp	7 copies
LR-07 homologue of MJ0794	836 bp	3 copies
LR-08 conserved hypothetical protein	696 bp	2 copies
LR-09 conserved hypothetical protein	628 bp	2 copies

(median, 9.8) and *A. fulgidus* where the skew is toward low pI (median, 6.3).

Multigene families. In *A. fulgidus* 719 genes (30% of the total) belong to 242 families with two or more members (Table 1). Of these families, 157 contained genes with biological roles. Most of these families contain genes assigned to the 'energy metabolism', 'transport and binding proteins', and 'fatty acid and phospholipid metabolism' categories (Table 2). The superfamily of ATP-binding subunits of ABC transporters is the largest, containing 40 members. The importance of catabolic degradation and signal recognition systems is reflected by the presence of two large superfamilies: acyl-CoA ligases and signal-transducing histidine kinases. *A. fulgidus* does not contain a homologue of the large 16-member family found in *M. jannaschii*⁹.

Repetitive elements. Three regions of the *A. fulgidus* genome contain short (<40 bp) direct repeats (Table 1). Two regions (SR-1A and SR-1B) contain 48 and 60 copies, respectively, of an identical 30-bp repeat interspersed with unique sequences averaging 40 bp. The third region (SR-2) contains 42 copies of a 37-bp repeat similar in sequence to the SR-1 repeat and interspersed with unique sequence averaging 41 bp. These repeated sequences are similar to the short repeated sequences found in *M. jannaschii*.

Nine classes of long (>500 bp) repeated sequences with ≥95% sequence identity were found (LR1-LR9; Table 1). LR-3 is a novel element with 14-bp inverted repeats and two genes, one of which has weak similarity to a transposase from *Halobacterium salinarium*. One copy of LR-3 interrupts AF2090, a homologue of a large *M. jannaschii* gene encoding a protein of unknown function. LR-4 and LR-6 encode putative transposases not identified in *M. jannaschii* that may represent IS elements. The remaining LR elements are not similar to known IS elements.

Central intermediary and energy metabolism

Sulphur oxide reduction may be the dominant respiratory process in anaerobic marine and freshwater environments, and is an important aspect of the sulphur cycle in anaerobic ecosystems¹². In this pathway, sulphate (SO_4^{2-}) is first activated to adenylylsulphate (adenosine-5'-phosphosulphate; APS), then reduced to sulphite and subsequently to sulphide^{1,13} (Fig. 3). The most important enzyme in dissimilatory sulphate reduction, adenylylsulphate reductase, reduces the activated sulphate to sulphite, releasing AMP. In *A. fulgidus*, the APS reductase has a high degree of similarity and identical physiological properties to APS reductases in sulphate-reducing delta proteobacteria¹⁴. A desulphoviridin-type sulphite reductase then adds six electrons to sulphite to produce sulphide. As in the Eubacteria, three sulphite-reductase genes, *dsrABD*, constitute an operon. The genes for adenylylsulphate reductase and sulphate adenylyltransferase reside in a separate operon. In *A. fulgidus*, sulphate can be replaced as an electron acceptor by both thiosulphate ($\text{S}_2\text{O}_3^{2-}$) and sulphite (SO_3^{2-}), but not by elemental sulphur.

A. fulgidus VC-16 has been shown to use lactate, pyruvate, methanol, ethanol, 1-propanol and formate as carbon and energy sources². Glucose has been described as a carbon source¹, but neither an uptake-transporter nor a catabolic pathway could be identified. Although it has been reported that *A. fulgidus* is incapable of growth on acetate⁶, multiple genes for acetyl-CoA synthetase (which converts acetate to acetyl-CoA) were found. The organism may degrade a variety of hydrocarbons and organic acids because of the presence of 57 β -oxidation enzymes, at least one lipase, and a minimum of five types of ferredoxin-dependent oxidoreductases (Fig. 3). The predicted β -oxidation system is similar to those in Eubacteria and mitochondria, and has not previously been described in the Archaea. *Escherichia coli* requires both the *fadD* and *fadL* gene products to import long-chain fatty acids across the cell envelope into the cytosol¹⁵. *A. fulgidus* has 14 acyl-CoA ligases related to FadD, but as expected given that it has no outer membrane, no

FadL. In *E. coli*, FadB has several metabolic functions, but in *A. fulgidus* these functions seem to be distributed among separate enzymes. For example, AF0435 encodes an orthologue of enoyl-CoA hydratase and resembles the amino-terminal domain of FadB. This gene is immediately upstream of a gene encoding an orthologue of 3-hydroxyacyl-CoA dehydrogenase that resembles the carboxy-terminal domain of FadB.

Acetyl-CoA is degraded by *A. fulgidus* through a C_1 -pathway, not by the citric acid cycle or glyoxylate bypass^{6,16,17}. This degradation is catalysed through the carbon monoxide dehydrogenase (CODH) pathway that consists of a five-subunit acetyl-CoA decarboxylase/synthase complex (ACDS) and five enzymes that are typically involved in methanogenesis¹⁸. In *A. fulgidus*, however, reverse methanogenesis occurs, resulting in CO_2 production. All of the enzymes and cofactors of methanogenesis from formylmethanofuran to N^5 -methyltetrahydromethanopterin are used, but the absence of methyl-CoM reductase eliminates the possibility of methane production by conventional pathways. Production of trace amounts of methane ($<0.1 \mu\text{mol ml}^{-1}$)¹⁹ is probably a result of the reduction of N^5 -methyltetrahydromethanopterin to methane and tetrahydromethanopterin by carbon monoxide (CO) dehydrogenase.

A. fulgidus also contains genes suggesting it has a second CO dehydrogenase system, homologous to that which enables *Rhodospirillum rubrum* to grow without light using CO as its sole energy source. Genes were detected for the nickel-containing CO dehydrogenase (CooS), an iron-sulphur redox protein, and a protein associated with the incorporation of nickel in CooS. These represent elements of a system that could catalyse the conversion of CO and H_2O to CO_2 and H_2 .

In contrast to *M. jannaschii*, *A. fulgidus* contains genes representing multiple catabolic pathways. Systems include CoA-SH-dependent ferredoxin oxidoreductases specific for pyruvate, 2-ketoisovalerate, 2-ketoglutarate and indolepyruvate, as well as a 2-oxoacid with little substrate specificity^{20,21}. Four genes with similarity to the tungsten-containing aldehyde ferredoxin oxidoreductase were also found²².

Biochemical pathways characteristic of eubacterial metabolism, including the pentose-phosphate pathway, the Entner-Doudoroff pathway, glycolysis and gluconeogenesis, are either completely absent or only partly represented (Fig. 3). *A. fulgidus* does not have typical eubacterial polysaccharide biosynthesis machinery, yet it has been shown to produce a protein and carbohydrate-containing biofilm²³. Nitrogen is obtained by importing inorganic molecules or degrading amino acids (Fig. 3); neither a glutamate dehydrogenase nor a relevant *fix* or *nif* gene is present.

The F_{420}H_2 :quinone oxidoreductase complex²⁴ is recognized as

Figure 2 Linear representation of the *A. fulgidus* genome illustrating the location of each predicted protein-coding region, RNA gene, and repeat element in the genome. Symbols for the transporters are as follows: AsO, arsenite; COH, sugar; P_i, phosphate; aa2, dipeptide; NH₄⁺, ammonium; a/o, arginine/lysine/ornithine; s/p, spermidine/putrescine; glyc, glycerol; Cl⁻, chloride; Fe²⁺, iron(II); Fe³⁺, iron(III); I, L, V, branched-chain amino acids; P, proline; pan, pantothenate; rib, ribose; lac, lactate; Mg²⁺/Co²⁺, magnesium and cobalt; gln, glutamine; NO₃⁻, nitrate; ox/for, oxalate/formate; maln, malonic acid; Hg²⁺, mercury; phs, polysaccharide; SO₄²⁻, sulphate; OCN⁻, cyanate; hex, hexuronate; phs, polysialic acid; K⁺, potassium channel; H⁺/Na⁺, sodium/proton antiporter; Na⁺/Cl⁻, sodium- and chloride-dependent transporter; P/G, osmoprotection protein; Cu²⁺, copper-transporting ATPase; +?, cation-transporting ATPase; ?, ABC-transporter without known function. Triplets associated with tRNAs represent the anticodon sequence. Numbers associated with GES represent the number of membrane-spanning domains (MSDs) according to Goldman, Engelman and Steiz scale as determined by TopPred²⁹. Genes whose identification is based on genes in *M. jannaschii* are indicated by circles. Of the 236 proteins containing at least one MSD, 124 of these had two or more MSDs.

the main generator of proton-motive force. However, our analysis indicates the presence of heterodisulphide reductase and several molybdopterin-binding oxidoreductases, with polysulphide, nitrate, dimethyl sulphoxide, and thiosulphate as potential substrates, which might contribute to energizing the cell membrane. *A. fulgidus*

contains a large number of flavoproteins, iron-sulphur proteins and iron-binding proteins that contribute to the general intracellular flow of electrons (Fig. 3). Detoxification enzymes include a peroxidase/catalase, an alkyl-hydroperoxide reductase, arsenate reductase, and eight NADH oxidases, presumably catalysing the

Figure 3 An integrated view of metabolism and solute transport in *A. fulgidus*. Biochemical pathways for energy production, biosynthesis of organic compounds, and degradation of amino acids, aldehydes and acids are shown with the central components of *A. fulgidus* metabolism, sulphate, lactate and acetyl-CoA highlighted. Pathways or steps for which no enzymes were identified are represented by a red arrow. A question mark is attached to pathways that could not be completely elucidated. Macromolecular biosynthesis of RNA, DNA and other lipids have been omitted. Membrane-associated reactions that establish the proton-motive force (PMF) and generate ATP (electron transport chain and V_0V_1 -ATPase) are linked to cytosolic pathways for energy production. The oxalate-formate antiporters (*oxlT*) may also contribute to the PMF by mediating electrogenic anion exchange. Each gene product with a predicted function in ion or solute transport is illustrated. Proteins are grouped by substrate specificity with transporters for cations (green), anions (red), carbohydrates/organic alcohols/acids (yellow), and amino acids/peptides/amines (blue) depicted. Ion-coupled permeases are represented by ovals (*mae1*, *exuT*, *panF*, *lctP*, *arsB*, *cynX*, *napA*/*nhe2*, *amt*, *feoB*, *trkAH*, *cat* and *putP* encode transporters for malate, hexuronate, pantothenate, lactate, arsenite, cyanate, sodium, ammonium, iron (II), potassium, arginine/lysine and proline, respectively). ATP-binding cassette (ABC) transport systems are shown as composite figures of ovals, diamonds and circles (*proVWX*, *glnHPQ*, *dppABCD*, *potABCD*, *braCDEFG*, *hemUV*, *nrtBC*, *cysAT*, *pstABC*, *rbsAC*, *rfaAB* correspond to gene products for proline, glutamine, dipeptide,

spermidine/putrescine, branch-chain amino acids, iron (III), nitrate, sulphate, phosphate, ribose and polysialic acid transport, respectively). All other porters drawn as rectangles (*glpF*, glycerol uptake facilitator; *copB*, copper transporting ATPase; *corA*, magnesium and cobalt transporter). Export and import of solutes is designated by arrows. The number of paralogous genes encoding each protein is indicated in brackets for cytoplasmic enzymes, or within the figure for transporters. Abbreviations: *acs*, acetyl-CoA synthetase; *aor*, aldehyde ferredoxin oxidoreductase; *aprAB*, adenylylsulphate reductase; *aspBC*, aspartate aminotransferase; *cdh*, acetyl-CoA decarbonylase/synthase complex; *cysC*, adenylylsulphate 3-phosphotransferase; *dld*, D-lactate dehydrogenase; *dsrABD*, sulphite reductase; *eno*, enolase; *fadA/acaB*, 3-ketoacyl-CoA thiolase; *fadD*, long-chain-fatty-acid-CoA ligase; *fad*, enoyl-CoA hydratase; *fadE (acd)*, acyl-CoA dehydrogenase; *glpA*, glycerol-3-phosphate dehydrogenase; *glpK*, glycerol kinase; *gltB*, glutamate synthase; *hbd*, 3-hydroxyacyl-CoA dehydrogenase; *ilvE*, branched-chain amino acid aminotransferase; *iorAB*, indolepyruvate ferredoxin oxidoreductase; *korABDG*, 2-ketoglutarate ferredoxin oxidoreductase; *lldD*, L-lactate dehydrogenase; *mcmA*, methylmalonyl-CoA mutase; *mdhA*, L-malate dehydrogenase; *oadAB*, oxaloacetate decarboxylase; *orAB*, 2-oxoacid ferredoxin oxidoreductase; *pflD*, pyruvate formate lyase 2; *porABDG*, pyruvate ferredoxin oxidoreductase; *ppsA*, phosphoenolpyruvate synthase; *prsA*, ribose-phosphate pyrophosphokinase; *sucAB*, 2-ketoglutarate dehydrogenase; *sat*, sulphate adenylyltransferase; TCA, tricarboxylic acid cycle; *vorABDG*, 2-ketoisovalerate ferredoxin oxidoreductase.

four-electron reduction of molecular oxygen to water, with the concurrent regeneration of NAD.

Transporters

A. fulgidus may synthesize several transporters for the import of carbon-containing compounds, probably contributing to its ability to switch from autotrophic to heterotrophic growth⁵. Both *M. jannaschii* and *A. fulgidus* have branched-chain amino-acid ABC transport systems and a transporter for the uptake of arginine and lysine. *A. fulgidus* encodes proteins for dipeptide, spermidine/putrescine, proline/glycine-betaine and glutamine uptake, as well as transporters for sugars and acids, rather like the membrane systems described in eubacterial heterotrophs. These compounds provide the necessary substrates for numerous biosynthetic and degradative pathways (Fig. 3).

Many *A. fulgidus* redox proteins are predicted to require iron. Correspondingly, iron transporters have been identified for the import of both oxidized (Fe^{3+}) and reduced (Fe^{2+}) forms of iron. There are duplications in functional and regulatory genes in both systems. The uptake of Fe^{3+} may depend on haemin or a haemin-like compound because *A. fulgidus* has orthologues to the eubacterial hem transport system proteins, HemU and HemV. *A. fulgidus* may also use the regulatory protein Fur to modulate Fe^{3+} transport; this protein is not present in *M. jannaschii*. Fe^{2+} uptake occurs through a modified Feo system containing FeoB. This is the third example of an isolated *feoB* gene: *M. jannaschii* and *Helicobacter pylori* also appear to lack *feoA*, implying that FeoA is not essential for iron transport in these organisms.

A complex suite of proteins regulates ionic homeostasis. Ten distinct transporters facilitate the flux of the physiological ions K^+ , Na^+ , NH_4^+ , Mg^{2+} , Fe^{2+} , Fe^{3+} , NO_3^- , SO_4^{2-} and inorganic phosphate (P_i). Most of these transporters have homologues in *M. jannaschii* and are therefore likely to be critical for nutrient acquisition during autotrophic growth. *A. fulgidus* has additional ion transporters for the elimination of toxic compounds including copper, cyanate and arsenite. As in *M. jannaschii*, the *A. fulgidus* genome contains two paralogous operons of cobalamin biosynthesis-cobalt transporters, *cbiMQO*.

Sensory functions and regulation of gene expression

Consistent with its extensive energy-producing metabolism and versatile system for carbon utilization, *A. fulgidus* has complex sensory and regulatory networks. These networks contain over 55 proteins with presumed regulatory functions, including members of the ArsR, AsnC and Sir2 families, as well as several iron-dependent repressor proteins. There are at least 15 signal-transducing histidine kinases, but only nine response regulators; this difference suggests there is a high degree of cross-talk between kinases and regulators. Only four response regulators appear to be in operons with histidine kinases, including those in the methyl-directed chemotaxis system (Che), which lies adjacent to the flagellar biosynthesis operon. Although rich in regulatory proteins, *A. fulgidus* apparently lacks regulators for response to amino-acid and carbon starvation as well as to DNA damage. Finally, *A. fulgidus* contains a homologue of the mammalian mitochondrial benzodiazepine receptor, which functions as a sensor in signal-transduction pathways²⁵. These receptors have been previously identified only in Proteobacteria and Cyanobacteria²⁵.

Replication, repair and cell division

A. fulgidus possesses two family B DNA polymerases, both related to the catalytic subunit of the eukaryal delta polymerase, as previously observed in the Sulfolobales²⁶. It also has a homologue of the proofreading ϵ subunit of *E. coli* Pol III, not previously observed in the Archaea. The DNA repair system is more extensive than that found in *M. jannaschii*, including a homologue of the eukaryal Rad25, a 3-methyladenine DNA glycosylase, and exodeoxynuclease

III. As well as reverse gyrase, topoisomerase I (ref. 9), and topoisomerase VI (ref. 27), the genes for the first archaeal DNA gyrase were identified.

A. fulgidus lacks a recognizable type II restriction-modification system, but contains one type I system. In contrast, two type II and three type I systems were identified in *M. jannaschii*. No homologue of the *M. jannaschii* thermonuclease was identified.

The cell-division machinery is similar to that of *M. jannaschii*, with orthologues of eubacterial *fts* and eukaryal *cdc* genes. However, several *cdc* genes found in *M. jannaschii*, including homologues of *cdc23*, *cdc27*, *cdc47* and *cdc54*, appear to be absent in *A. fulgidus*.

Transcription and translation

A. fulgidus and *M. jannaschii* have transcriptional and translational systems distinct from their eubacterial and eukaryal counterparts. In both, the RNA polymerase contains the large universal subunits and five smaller subunits found in both Archaea and eukaryotes. Transcription initiation is a simplified version of the eukaryotic mechanism^{28,29}. However, *A. fulgidus* alone has a homologue of eukaryotic TBP-interacting protein 49 not seen in *M. jannaschii*, but apparently present in *Sulfolobus solfataricus*.

Translation in *A. fulgidus* parallels *M. jannaschii* with a few exceptions. The organism has only one rRNA operon with an Ala-tRNA gene in the spacer and lacks a contiguous 5S rRNA gene. Genes for 46 tRNAs were identified, five of which contain introns in the anticodon region that are presumably removed by the intron excision enzyme EndA. The gene for selenocysteine tRNA (SelC) was not found, nor were the genes for SelA, SelB and SelD. With the exception of Asp-tRNA^{GTC} and Val-tRNA^{CAC}, tRNA genes are not linked in the *A. fulgidus* genome. The RNA component of the tRNA maturation enzyme RNase P is present. Both *A. fulgidus* and *M. jannaschii* appear to possess an enzyme that inserts the tRNA-modified nucleoside archaeosine, but only *A. fulgidus* has the related enzyme that inserts the modified base queuine.

Both *A. fulgidus* and *M. jannaschii* lack glutamine synthetase and asparagine synthetase; the relevant tRNAs are presumably aminoacylated with glutamic and aspartic acids, respectively. An enzymatic *in situ* transamidation then converts the amino acid to its amide form, as seen in other Archaea and in Gram-positive Eubacteria³⁰. Indeed, genes for the three subunits of the Glu-tRNA amidotransferase (*gatABC*) have been identified in *A. fulgidus*. The Lys aminoacyl-tRNA synthetase in both organisms is a class I-type, not a class II-type³¹. *A. fulgidus* possesses a normal tRNA synthetase for both Cys and Ser, unlike *M. jannaschii* in which the former was not identifiable and the latter was unusual⁹.

M. jannaschii has a single gene belonging to the TCP-1 chaperonin family, whereas *A. fulgidus* has two that encode subunits α and β of the thermosome. Phylogenetic analysis of the archaeal TCP-1 family indicates that these *A. fulgidus* genes arose by a recent species-specific gene duplication, as is the case for the two subunits of the *Thermoplasma acidophilum* thermosome³² and the *Sulfolobus shibatae* rosettasome³³. As in *M. jannaschii*, no *dnaK* gene was identified.

Biosynthesis of essential components

Like most autotrophic microorganisms, *A. fulgidus* is able to synthesize many essential compounds, including amino acids, cofactors, carriers, purines and pyrimidines. Many of these biosynthetic pathways show a high degree of conservation between *A. fulgidus* and *M. jannaschii*. These two Archaea are similar in their biosynthetic pathways for siroheme, cobalamin, molybdopterin, riboflavin, thiamin and nicotinate, the role category with greatest conservation between these two organisms being amino-acid biosynthesis. Of 78 *A. fulgidus* genes assigned to amino-acid biosynthetic pathways, at least 73 (94%) have homologues in *M. jannaschii*. For both archaeal species, amino-acid biosynthetic pathways resemble those of *Bacillus subtilis* more closely than

those of *E. coli*. For example, in *A. fulgidus* and *M. jannaschii*, tryptophan biosynthesis is accomplished by seven enzymes, TrpA, B, C, D, E, F, G as in *B. subtilis*, rather than by five enzymes, TrpA, B, C, D, E (including the bifunctional TrpC and TrpD) as found in *E. coli*.

No biotin biosynthetic genes were identified, yet biotin can be detected in *A. fulgidus* cell extracts³⁴, and several genes encode a biotin-binding consensus sequence. Similarly, *A. fulgidus* lacks the genes for pyridoxine biosynthesis although pyridoxine can be found in cell extracts (albeit at lower levels than seen in *E. coli* and several Archaea³⁴). No gene encoding ferrochelatase, the terminal enzyme in haem biosynthesis, has been identified, although *A. fulgidus* is known to use cytochromes³⁴. These cofactors may be obtained by mechanisms that we have not recognized. Although all of the enzymes required for pyrimidine biosynthesis appear to be present, three enzymes in the purine pathway (GAR transformylase, AICAR formyltransferase and the ATPase subunit of AIR carboxylase) have not been identified, presumably because they exist as new isoforms.

The Archaea share a unique cell membrane composed of ether lipids containing a glycerophosphate backbone with a 2,3-*sn* stereochemistry³⁵ for which there are multiple biosynthetic pathways³⁶. In the case of *Halobacterium cutirubrum*, the backbone is apparently obtained by enantiomeric inversion of *sn*-glycerol-3-phosphate; in *Sulfolobus acidocaldarius* and *Methanobacterium thermoautotrophicum*, *sn*-glycerol-1-phosphate dehydrogenase builds the backbone from dihydroxyacetonephosphate. An orthologue of *sn*-glycerol-1-phosphate dehydrogenase has been identified in *A. fulgidus*, suggesting that the latter pathway is present.

Conclusions

Although *A. fulgidus* has been studied since its discovery ten years ago¹, the completed genome sequence provides a wealth of new information about how this unusual organism exploits its environment. For example, its ability to reduce sulphur oxides has been well characterized, but genome sequence data demonstrate that *A. fulgidus* has a great diversity of electron transport systems, some of unknown specificity. Similarly, *A. fulgidus* has been characterized as a scavenger with numerous potential carbon sources, and its gene complement reveals the extent of this capability. *A. fulgidus* appears to obtain carbon from fatty acids through β -oxidation, from degradation of amino acids, aldehydes and organic acids, and perhaps from CO.

A. fulgidus has extensive gene duplication in comparison with other fully sequenced prokaryotes. For example, in the fatty acid and phospholipid metabolism category, there are 10 copies of 3-hydroxyacyl-CoA dehydrogenase, 12 copies of 3-ketoacyl-CoA thiolase, and 12 of acyl-CoA dehydrogenase. The duplicated proteins are not identical, and their presence suggests considerable metabolic differentiation, particularly with respect to the pathways for decomposing and recycling carbon by scavenging fatty acids. Other categories show similar, albeit less dramatic, gene redundancy. For example, there are six copies of acetyl-CoA synthetase and four aldehyde ferredoxin oxidoreductases for fermentation, as well as four copies of aspartate aminotransferase for amino-acid biosynthesis. These observations, together with the large number of paralogous gene families, suggest that gene duplication has been an important evolutionary mechanism for increasing physiological diversity in the Archaeoglobales.

A comparison of two archaeal genomes is inadequate to assess the diversity of the entire domain. Given this caveat, it is nevertheless possible to draw some preliminary conclusions from the comparison of *M. jannaschii* and *A. fulgidus*. A comparison of the gene content of these Archaea reveals that gene conservation varies significantly between role categories, with genes involved in transcription, translation and replication highly conserved; approximately 80% of the *A. fulgidus* genes in these categories have homologues in *M. jannaschii*. Biosynthetic pathways are also

highly conserved, with approximately 80% of the *A. fulgidus* biosynthetic genes having homologues in *M. jannaschii*. In contrast, only 35% of the *A. fulgidus* central intermediary metabolism genes have homologues, reflecting their minimal metabolic overlap.

Over half of the *A. fulgidus* ORFs (1,290) have no assigned biological role. Of these, 639 have no database match. The remaining 651, designated 'conserved hypothetical proteins', have sequence similarity to hypothetical proteins in other organisms, two-thirds with apparent homologues in *M. jannaschii*. These shared hypothetical proteins will probably add to our understanding of the genetic repertoire of the Archaea. Analysis of the *A. fulgidus* and other archaeal and eubacterial genomes will provide the information necessary to begin to define a core set of archaeal genes, as well as to better understand prokaryotic diversity. □

Methods

Whole-genome random sequencing procedure. The type strain, *A. fulgidus* VC-16, was grown from a culture derived from a single cell isolated by optical tweezers³⁷ and provided by K. O. Stetter (University of Regensburg). Cloning, sequencing and assembly were essentially as described previously for genomes sequenced by TIGR^{38–40}. One small-insert and one medium-insert plasmid library were generated by random mechanical shearing of genomic DNA. One large-insert lambda (λ) library was generated by partial *Tsp509I* digestion and ligation to λ -DASHII/*EcoRI* vector (Stratagene). In the initial random sequencing phase, 6.7-fold sequence coverage was achieved with 27,150 sequences from plasmid clones (average read length 500 bases) and 1,850 sequences from λ -clones. Both plasmid and λ -sequences were jointly assembled using TIGR assembler⁴¹, resulting in 152 contigs separated by sequence gaps and five groups of contigs separated by physical gaps. Sequences from both ends of 560 λ -clones served as a genome scaffold, verifying the orientation, order and integrity and the contigs. Sequence gaps were closed by editing the ends of sequence traces and/or primer walking on plasmid or λ -clones clones spanning the respective gap. Physical gaps were closed by combinatorial polymerase chain reaction (PCR) followed by sequencing of the PCR product. At the end of gap closure, 90 regions representing 0.33% of the genome had only single-sequence coverage. These regions were confirmed with terminator reactions to ensure a minimum of 2-fold sequence coverage for the whole genome. The final genome sequence is based on 29,642 sequences, with a 6.8-fold sequence coverage. The linkage between the terminal sequences of 2,101 clones from the small-insert plasmid library (average size 1,419 bp) and 8,726 clones from the medium-insert plasmid library (average size 2,954 bp) supported the genome scaffold formed by the λ -clones (average size 16,381 bp), with 96.9% of the genome covered by λ -clones. The reported sequence differs in 20 positions from the 14,389 bp of DNA in a total of 11 previously published *A. fulgidus* genes.

ORF prediction and gene family identification. Coding regions (ORFs) were identified using a combination strategy based on two programs. Initial sets of ORFs were derived with GeneSmith (H.O.S., unpublished), a program that evaluates ORF length, separation and overlap between ORFs, and with CRITICA (J.H.B. & G.J.O., unpublished), a coding region identification tool using comparative analysis. The two largely overlapping sets of ORFs were merged into one joint set containing all members of both initial sets. ORFs were searched against a non-redundant protein database using BLASTX¹⁰ and those shorter than 30 codons 'coding' for proteins without a database match were eliminated. Frameshifts were detected and corrected where appropriate as described previously⁴⁰. Remaining frameshifts are considered authentic and corresponding regions were annotated as 'authentic frameshift'. In total, 527 hidden Markov models, based upon conserved protein families (PFAM version 2.0), were searched with HMMER to determine ORF membership in families and superfamilies⁴². Families of paralogous genes were constructed as described previously⁴⁰. TopPred⁴³ was used to identify membrane-spanning domains in proteins.

Received 9 September; accepted 4 November 1997.

1. Stetter, K. O., Lauerer, G., Thomm, M. & Neuner, A. Isolation of extremely thermophilic sulfate reducers: Evidence for a novel branch of archaeobacteria. *Science* **236**, 822–824 (1987).
2. Stetter, K. O., in *The Prokaryotes* (eds Balows, A., Trüper, H. G., Dworkin, M., Harder, W. & Schleifer, K. H.) 707–711 (Springer, Berlin, 1992).
3. Stetter, K. O. Microbial life in hyperthermal environments: Microorganisms from exotic environments continue to provide surprises about life's extremities. *ASM News* **61**, 285–290 (1995).

4. Stetter, K. O. *Archaeoglobus fulgidus* gen. nov., sp. nov.: a new taxon of extremely thermophilic archaeobacteria. *Syst. Appl. Microbiol.* **10**, 172–173 (1988).
5. Stetter, K. O. *et al.* Hyperthermophilic archaea are thriving in deep North Sea and Alaskan oil reservoirs. *Nature* **365**, 743–745 (1993).
6. Vorholt, J., Kunow, J., Stetter, K. O. & Thauer, R. K. Enzymes and coenzymes of the carbon monoxide dehydrogenase pathway for autotrophic CO₂ fixation in *Archaeoglobus lithotrophicus* and the lack of carbon monoxide dehydrogenase in the heterotrophic *A. profundus*. *Arch. Microbiol.* **163**, 112–118 (1995).
7. Woese, C. R. & Fox, G. E. Phylogenetic structure of the prokaryotic domain: The primary kingdoms. *Proc. Natl Acad. Sci. USA* **74**, 5088–5090 (1977).
8. Woese, C. R., Kandler, O. & Wheelis, M. L. Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. *Proc. Natl Acad. Sci. USA* **87**, 4576–4579 (1990).
9. Bult, C. J. *et al.* Complete genome sequence of the methanogenic archaeon *Methanococcus jannaschii*. *Science* **273**, 1058–1073 (1996).
10. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search tool. *J. Mol. Biol.* **215**, 403–410 (1990).
11. Riley, M. Functions of gene products of *Escherichia coli*. *Microbiol. Rev.* **57**, 862–952 (1993).
12. Cooling, F. B. III, Maloney, C. L., Nagel, E., Tabinowski, J. & Odom, J. M. Inhibition of sulfate respiration by 1,8-dihydroxyanthraquinone and other anthraquinone derivatives. *Appl. Environ. Microbiol.* **62**, 2999–3004 (1996).
13. Thauer, R. K. & Kunow, J. in *Sulfate Reducing Bacteria* (ed. Barton, L. L.) 33–48 (Plenum, New York, 1995).
14. Speich, D. *et al.* Adenylylsulfate reductase from the sulfate-reducing archaeon *Archaeoglobus fulgidus*: cloning and characterization of the genes and comparison of the enzyme with other iron-sulfur flavoproteins. *Microbiology* **140**, 1273–1284 (1994).
15. Clark, D. P. & Cronan, J. E. Jr in *Escherichia coli* and *Salmonella typhimurium: Cellular and Molecular biology* (ed. Neidhardt, F. C.) 343–357 (ASM Press, Washington DC, 1996).
16. Möller-zirkhan, D. & Thauer, R. K. Anaerobic lactate oxidation to 3 CO₂ by *Archaeoglobus fulgidus* via the carbon monoxide dehydrogenase pathway: demonstration of the acetyl-CoA carbon-carbon cleavage reaction in cell extracts. *Arch. Microbiol.* **153**, 215–218 (1990).
17. Schauder, R., Eikmanns, B., Thauer, R. K., Widdel, F. & Fuchs, G. Acetate oxidation to CO₂ in anaerobic-bacteria via a novel pathway not involving reactions of the citric-acid cycle. *Arch. Microbiol.* **145**, 162–172 (1986).
18. Dai, Y.-R. *et al.* Acetyl-CoA decarbonylase/synthase complex from *Archaeoglobus fulgidus*: purification, characterization, and properties. *Arch. Microbiol.* (submitted).
19. Gorris, L. G. M., Voet, A. C. W. A. & van der Drift, C. Structural characteristics of methanogenic cofactors in the non-methanogenic archaeobacterium *Archaeoglobus fulgidus*. *BioFactors* **3**, 29–35 (1991).
20. Zhang, Q., Iwasaki, T., Wakagi, T. & Oshima, T. 2-oxoacid:ferredoxin oxidoreductase from the thermoacidophilic archaeon, *Sulfolobus* sp. strain 7. *J. Biochem.* **120**, 587–599 (1996).
21. Tersteegen, A., Linder, D., Thauer, R. K. & Hedderich, R. Structures and functions of four anabolic 2-oxoacid oxidoreductases in *Methanobacterium thermoautotrophicum*. *Eur. J. Biochem.* **244**, 862–868 (1997).
22. Kletzin, A. & Adams, M. W. W. Molecular and phylogenetic characterization of pyruvate and 2-ketoisovalerate ferredoxin oxidoreductases from *Pyrococcus furiosus* and pyruvate ferredoxin oxidoreductase from *Thermotoga maritima*. *J. Bacteriol.* **178**, 248–257 (1996).
23. LaPaglia, C. & Hartzell, P. L. Stress-induced production of biofilm in the hyperthermophile *Archaeoglobus fulgidus*. *Appl. Environ. Microbiol.* **63**, 3158–3163 (1997).
24. Kunow, J., Linder, D., Stetter, K. O. & Thauer, R. K. F₄₂₀H₂: quinone oxidoreductase from *Archaeoglobus fulgidus*—characterization of a membrane-bound multisubunit complex containing FAD and iron–sulfur clusters. *Eur. J. Biochem.* **223**, 503–511 (1994).
25. Yeliseev, A. A., Krueger, K. E. & Kaplan, S. A mammalian mitochondrial drug receptor functions as a bacterial “oxygen” sensor. *Proc. Natl Acad. Sci. USA* **94**, 5101–5106 (1997).
26. Edgell, D. R., Klenk, H.-P. & Doolittle, W. F. Gene duplications in evolution of archaeal family B DNA polymerases. *J. Bacteriol.* **179**, 2632–2640 (1997).
27. Bergerat, A. *et al.* An atypical topoisomerase II from archaea with implications for meiotic recombination. *Nature* **386**, 414–417 (1997).
28. Marsh, T. L., Reich, C. I., Whitelock, R. B. & Olsen, G. J. Transcription factor IID in the Archaea: sequences in the *Thermococcus celer* genome would encode a product closely related to the TATA-binding protein of eukaryotes. *Proc. Natl Acad. Sci. USA* **91**, 4180–4184 (1994).
29. Kosa, P. E., Ghosh, G., DeDecker, B. S. & Sigler, P. B. The 2.1-Å crystal structure of an archaeal preinitiation complex: TATA-box-binding protein/transcription factor (II)B core/TATA-box. *Proc. Natl Acad. Sci. USA* **94**, 6042–6047 (1997).
30. Curnow, A. W. *et al.* Glu-tRNA^{Gln} amidotransferase: a novel heterotrimeric enzyme required for correct decoding of glutamine codons during translation. *Proc. Natl Acad. Sci. USA* **94**, 11819–11826 (1997).
31. Ibba, M., Bobo, J. L., Rosa, P. A. & Soll, D. Archaeal-type lysyl-tRNA synthetase in the Lyme disease spirochete *Borrelia burgdorferi*. *Proc. Natl Acad. Sci. USA* (submitted).
32. Waldmann, T., Lupas, A., Kellermann, J., Peters, J. & Baumeister, W. Primary structure of the thermosome from *Thermoplasma acidophilum*. *Hoppe-Seyler's Biol. Chem.* **376**, 119–126 (1995).
33. Kagawa, H. K. *et al.* The 60 kDa heat shock proteins in the hyperthermophilic archaeon *Sulfolobus shibatae*. *J. Mol. Biol.* **253**, 712–725 (1995).
34. Noll, K. M. & Barber, T. S. Vitamin contents of archaeobacteria. *J. Bacteriol.* **170**, 4315–4321 (1988).
35. Thornebene, T. G. & Langworthy, T. A. Diphytanyl and dibiphytanyl glycerol ether lipids of methanogenic archaeobacteria. *Science* **203**, 51–53 (1979).
36. Nishihara, M. & Koga, Y. sn-glycerol-1-phosphate dehydrogenase in *Methanobacterium thermoautotrophicum*: key enzyme in biosynthesis of the enantiomeric glycerophosphate backbone of ether phospholipids of archaeobacteria. *J. Biochem.* **117**, 933–935 (1995).
37. Huber, R. *et al.* Isolation of a hyperthermophilic archaeum predicted by *in situ* RNA analysis. *Nature* **376**, 57–58 (1995).
38. Fleischmann, R. D. *et al.* Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science* **269**, 496–511 (1995).
39. Fraser, C. M. *et al.* The minimal gene complement of *Mycoplasma genitalium*. *Science* **270**, 397–403 (1995).
40. Tomb, J.-F. *et al.* The complete genome sequence of the gastric pathogen *Helicobacter pylori*. *Nature* **388**, 539–547 (1997).
41. Sutton, G. G., White, O., Adams, M. D. & Kerlavage, A. R. TIGR Assembler: A new tool for assembling large shotgun sequencing projects. *Genome Sequence Technol.* **1**, 9–19 (1995).
42. Sonnhammer, E. L., Eddy, S. R. & Durbin, R. Pfam: A comprehensive database of protein families based on seed alignments. *Proteins* **28**, 405–420 (1997).
43. Claros, M. G. & von Heijne, G. TopPred II: an improved software for membrane protein structure predictions. *Comput. Appl. Biosci.* **10**, 685–686 (1994).

Acknowledgements. We thank M. Heaney, J. Scott and R. Shirley for software and database support; V. Sapiro, B. Vincent, J. Meehan and D. Maas for computer system support; B. Cameron and D. J. Doyle for editorial assistance; and K. O. Stetter for providing *A. fulgidus* VC-16. This work was supported by the US Department of Energy.

Correspondence and requests for materials should be addressed to J.C.V. (e-mail: gaf@tigr.org). The annotated genome sequence and the gene family alignments are available on the World-Wide Web at <http://www.tigr.org/db/mdb/afdb/afdb.html>. The sequence has been deposited in GenBank with accession number AE000782.

Table 2 . List of *A. fulgidus* genes with putative identification. Gene numbers correspond to those in Fig. 2. Percentages represent per cent identities.

AMINO ACID BIOSYNTHESIS			CELLULAR PROCESSES		
<i>General</i>			<i>General</i>		
AF0906	hydantoin utilization protein A (hyaA)	27.4%	AF0722	cobalamin biosynthesis precorrin-6Y methylase (cbiE)	32.4%
<i>Aromatic amino acid family</i>			AF0732	cobalamin biosynthesis precorrin-8W decarboxylase (cbiT)	30.8%
AF0228	3-dehydroquinate dehydratase (aroD)	36.8%	AF1336	cobalamin biosynthesis protein (cbiB)	38.4%
AF1497	5-enolpyruvylshikimate 3-phosphate synthase (aroA)	41.5%	AF0723	cobalamin biosynthesis protein (cbiD)	36.3%
AF1603	anthranilate synthase component I (trpE)	43.7%	AF0728	cobalamin biosynthesis protein (cbiM-1)	51.4%
AF1604	anthranilate synthase component II (trpD)	43.8%	AF1843	cobalamin biosynthesis protein (cbiM-2)	41.2%
AF1602	anthranilate synthase component II (trpG)	50.0%	AF0731	cobalt transport ATP-binding protein (cbiO-1)	47.2%
AF0227	chorismate mutase/prephenate dehydratase (pheA)	32.2%	AF1841	cobalt transport ATP-binding protein (cbiO-2)	41.1%
AF0670	chorismate synthase (aroC)	55.3%	AF0729	cobalt transport protein (cbiN)	56.0%
AF1601	phosphoribosyl anthranilate isomerase (trpF)	37.1%	AF0730	cobalt transport protein (cbiQ-1)	32.6%
AF2327	shikimate 5-dehydrogenase (aroC)	43.1%	AF1842	cobalt transport protein (cbiQ-2)	30.3%
AF0343	tryptophan repressor binding protein (wrbA)	46.6%	AF1338	cobryric acid synthase (cbp)	44.5%
AF1599	tryptophan synthase, subunit alpha (trpA)	39.5%	AF2229	cobrynic acid α -clamidase synthase (cblA)	42.3%
AF1240	tryptophan synthase, subunit beta (trpB-1)	39.4%	AF1241	glutamate 1-semialdehyde aminotransferase (hemL)	54.3%
AF1600	tryptophan synthase, subunit beta (trpB-2)	64.1%	AF1975	glutamyl-tRNA reductase (hemA)	42.7%
<i>Aspartate family</i>			AF1594	heme biosynthesis protein (nirH)	25.2%
AF2112	5-methyltetrahydropteroyltryglutamate-homocysteine methyltransferase (metE)	28.1%	AF1125	heme biosynthesis protein (nir-1)	38.7%
AF0882	asparaginase (asnA)	45.9%	AF2009	heme biosynthesis protein (nir-2)	31.8%
AF1439	asparagine synthetase (asnB)	36.9%	AF1593	heme d1 biosynthesis protein (nirD)	29.4%
AF2366	aspartate aminotransferase (aspB-1)	42.3%	AF1311	oxygen-independent coporphyrinogen III oxidase, putative	27.1%
AF2129	aspartate aminotransferase (aspB-2)	45.4%	AF1242	porphobilinogen deaminase (hemC)	46.3%
AF1623	aspartate aminotransferase (aspB-3)	39.4%	AF1974	porphobilinogen synthase (hemB)	60.4%
AF0409	aspartate aminotransferase (aspB-4)	45.2%	AF1784	protoporphyrinogen oxidase (hemK)	33.5%
AF1417	aspartate aminotransferase (aspC)	46.2%	AF0422	uroporphyrin-III C-methyltransferase (cysG-1)	41.7%
AF0700	aspartate kinase (lysC)	48.1%	AF1243	uroporphyrin-III C-methyltransferase (cysG-2)	52.5%
AF1422	aspartate racemase	48.0%	AF0116	uroporphyrinogen III synthase (hemD)	27.4%
AF1506	aspartate-semialdehyde dehydrogenase (asd)	60.9%	<i>Menaquinone and ubiquinone</i>		
AF0800	diaminopimelate decarboxylase (lysA)	45.6%	AF2176	4-hydroxybenzoate octaprenyltransferase (ubiA)	41.6%
AF0747	diaminopimelate epimerase (dapF)	45.8%	AF0404	4-hydroxybenzoate octaprenyltransferase, putative	30.6%
AF0909	dihydrodipicolinate reductase (dapB)	48.6%	AF2413	coenzyme PQQ synthesis protein (pqeE)	30.5%
AF0910	dihydrodipicolinate synthase (dapA)	51.0%	AF1191	dihydroxynaphthoic acid synthase (menB)	54.6%
AF0935	homoserine dehydrogenase (hom)	47.9%	AF1551	octaprenyl-diphosphate synthase (ispB)	33.2%
AF0886	S-adenosylhomocysteine hydrolase (ahcY-1)	31.7%	AF0140	ubiquinone/menaquinone biosynthesis methyltransferase (ubiE)	31.0%
AF2000	S-adenosylhomocysteine hydrolase (ahcY-2)	67.3%	<i>Molybdopterin</i>		
AF0051	succinyl-diaminopimelate desuccinylase (dapE-1)	30.5%	AF2006	molybdenum cofactor biosynthesis protein (moaA)	47.8%
AF0904	succinyl-diaminopimelate desuccinylase (dapE-2)	43.8%	AF0265	molybdenum cofactor biosynthesis protein (moaB)	44.4%
AF0551	threonine synthase (thrC-1)	40.5%	AF2150	molybdenum cofactor biosynthesis protein (moaC)	62.0%
AF1316	threonine synthase (thrC-2)	61.0%	AF0931	molybdenum cofactor biosynthesis protein (moaA-1)	50.8%
<i>Glutamate family</i>			AF0930	molybdenum cofactor biosynthesis protein (moaA-2)	44.8%
AF1280	acetylglutamate kinase (argB)	56.1%	AF0161	molybdenum cofactor biosynthesis protein (moaA-3)	30.0%
AF2286	acetylglutamate kinase, putative	23.0%	AF0531	molybdenum cofactor biosynthesis protein (moaB)	44.0%
AF0080	acetylornithine aminotransferase (argD-1)	48.3%	AF1022	molybdenum-pterin-binding protein (moaB)	39.3%
AF1815	acetylornithine aminotransferase (argD-2)	36.2%	AF1624	molybdopterin converting factor, subunit 1 (moaD)	36.6%
AF0522	acetylornithine deacetylase (argE)	29.4%	AF2179	molybdopterin converting factor, subunit 2 (moaE)	33.3%
AF0883	argininosuccinate lyase (argH)	42.2%	AF2005	molybdopterin-guanine dinucleotide biosynthesis protein A (moaA)	33.2%
AF2252	argininosuccinate synthetase (argG)	62.0%	AF2253	molybdopterin-guanine dinucleotide biosynthesis protein B (moaB)	40.0%
AF1147	glutamate N-acetyltransferase (argI)	47.8%	<i>Pantothenate</i>		
AF0953	glutamate synthase (gltB)	57.9%	AF1645	pantothenate metabolism flavoprotein (dhp)	42.4%
AF0949	glutamine synthetase (glnA)	43.3%	<i>Riboflavin</i>		
AF2071	N-acetyl-gamma-glutamyl-phosphate reductase (argC)	53.3%	AF0494	GTP cyclohydrolase II (ribA-1)	44.5%
AF1255	ornithine carbamoyltransferase (argF)	51.7%	AF2107	GTP cyclohydrolase II (ribA-2)	47.1%
<i>Pyruvate family</i>			AF1416	riboflavin synthase (ribC)	53.3%
AF0957	2-isopropylmalate synthase (leuA-1)	53.5%	AF2128	riboflavin synthase, subunit beta (ribE)	75.9%
AF0219	2-isopropylmalate synthase (leuA-2)	53.9%	AF2007	riboflavin-specific deaminase (ribG)	43.7%
AF2199	3-isopropylmalate dehydratase, large subunit (leuC)	49.3%	<i>Thiamine</i>		
AF0629	3-isopropylmalate dehydratase, small subunit (leuD-1)	56.4%	AF2075	hydroxyethylthiazole kinase (thiM)	33.6%
AF1761	3-isopropylmalate dehydratase, small subunit (leuD-2)	57.1%	AF2208	hydroxymethylpyrimidine phosphate kinase (thiD)	35.5%
AF0628	3-isopropylmalate dehydrogenase (leuB)	59.2%	AF1695	thiamine biosynthesis protein (apbA)	36.9%
AF1720	acetoacetylase synthase, large subunit (ilvB-1)	57.5%	AF2412	thiamine biosynthesis protein (thiC)	60.2%
AF1780	acetoacetylase synthase, large subunit (ilvB-2)	32.1%	AF0553	thiamine biosynthesis protein (thiF)	38.1%
AF2015	acetoacetylase synthase, large subunit (ilvB-3)	34.1%	AF0088	thiamine biosynthesis protein, putative	28.2%
AF2100	acetoacetylase synthase, large subunit (ilvB-4)	38.4%	AF0702	thiamine biosynthetic enzyme (thiI)	50.0%
AF1719	acetoacetylase synthase, small subunit (ilvN)	60.4%	AF0733	thiamine monophosphate kinase (thiL)	30.4%
AF1672	acetoacetylase synthase, small subunit, putative	29.7%	AF2074	thiamine phosphate pyrophosphorylase (thiE)	45.5%
AF0933	branched-chain amino acid aminotransferase (ilvE)	59.0%	<i>Pyridine nucleotides</i>		
AF1014	dihydroxy-acid dehydratase (ilvD)	54.5%	AF1839	NH(3)-dependent NAD ⁺ synthetase (nadE)	52.0%
AF1985	ketol-acid reductoisomerase (ilvC)	61.8%	AF1837	nicotinate-nucleotide pyrophosphorylase (nadC)	43.2%
<i>Serine family</i>			AF1837	quinolinate synthetase (nadA), authentic frameshift	53.9%
AF0813	phosphoglycerate dehydrogenase (serA)	48.8%	CELL ENVELOPE		
AF2138	phosphoserine phosphatase (serB)	50.7%	<i>Membranes, lipoproteins, and porins</i>		
AF0273	sarcosine oxidase, subunit alpha (soxA)	31.1%	AF1420	membrane protein	51.9%
AF0274	sarcosine oxidase, subunit beta (soxB)	26.5%	AF1354	membrane protein, putative	32.8%
AF0852	serine hydroxymethyltransferase (glyA)	56.1%	<i>Surface polysaccharides, lipopolysaccharides and antigens</i>		
<i>Histidine family</i>			AF0324	dTDP-glucose 4,6-dehydratase (rfbB)	50.0%
AF0690	ATP phosphoribosyltransferase (hisG)	31.9%	AF0043	first mannoolyl transferase (wbaZ-1)	30.0%
AF0212	histidinol dehydrogenase (hisD)	51.9%	AF0806	first mannoolyl transferase (wbaZ-2)	29.0%
AF2002	histidinol-phosphate aminotransferase (hisC-1)	39.8%	AF1728	galactosyltransferase	26.9%
AF2024	histidinol-phosphate aminotransferase (hisC-2)	36.8%	AF0044	GDP-D-mannose dehydratase (gmd-1), authentic frameshift	40.7%
AF0985	imidazoleglycerol-phosphate synthase (hisB)	42.2%	AF1142	glucose-1-phosphate cytidylyltransferase (rfbF)	38.6%
AF0819	imidazoleglycerol-phosphate synthase, cyclase subunit (hisF)	67.0%	AF0242	glucose-1-phosphate thymidyltransferase (graD-1)	27.7%
AF2285	imidazoleglycerol-phosphate synthase, subunit H (hisH)	44.4%	AF0325	glucose-1-phosphate thymidyltransferase (graD-2)	45.2%
AF0509	imidazoleglycerol-phosphate synthase, subunit H, putative	43.2%	AF0321	glycosyl transferase	30.7%
AF1950	phosphoribosyl-AMP cyclohydrolase/phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0387	glycosyltransferase, putative	33.8%
AF0713	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0467	immunogenic protein (bcsp31-1)	34.7%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0635	immunogenic protein (bcsp31-2)	44.3%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0988	immunogenic protein (bcsp31-3)	28.3%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0602	LPS biosynthesis protein, putative	29.6%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0617	LPS biosynthesis protein, putative	29.0%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0607	LPS glycosyltransferase, putative	29.7%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0326	mannose-1-phosphate guanylyltransferase (rfbM), authentic frameshift	42.4%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF1097	mannose-6-phosphate isomerase/mannose-1-phosphate guanylyl transferase (manC)	43.1%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0035	mannosephosphate isomerase, putative	31.3%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0045	mannosyltransferase A (mtfA)	38.7%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0311	O-antigen biosynthesis protein (rfbC), authentic frameshift	30.6%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0458	phosphomannomutase (pmn)	39.5%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0595	polysaccharide biosynthesis protein, putative	24.1%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0322	ramnosyl transferase (rfbQ)	27.5%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0323	spore coat polysaccharide biosynthesis protein (spk-2), authentic frameshift	36.3%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0620	succinoglycan biosynthesis protein (exoM)	24.8%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0361	UDP-glucose 4-epimerase (galE-1)	38.6%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF2016	UDP-glucose 4-epimerase (galE-2)	30.0%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0302	UDP-glucose dehydrogenase (ugd-1)	43.8%
AF0986	phosphoribosyl-ATP pyrophosphorylase (hisE)	59.6%	AF0696	UDP-glucose dehydrogenase (ugd-2)	44.1%
<i>BIOSYNTHESIS OF COFACTORS, PROSTHETIC GROUPS, AND CARRIERS</i>			<i>Surface structures</i>		
<i>General</i>			AF1054	flagellin (flaB-1)	30.0%
AF1855	2,3-dihydroxybenzoate-AMP ligase (entE)	27.2%	AF1055	flagellin (flaB-2)	31.1%
AF1070	coenzyme F390 synthetase (ftsA-1)	30.3%	AF0275	surface layer protein B (slgB-1)	30.8%
AF1671	coenzyme F390 synthetase (ftsA-2)	31.9%	AF1413	surface layer protein B (slgB-2)	29.9%
AF2013	coenzyme F390 synthetase (ftsA-3)	30.4%	<i>ENERGY METABOLISM</i>		
AF2151	isochorismatase (entB)	31.2%	<i>Amino acids and amines</i>		
AF1414	dihydropterolate synthase	40.8%	AF1958	2-hydroxyglutaryl-CoA dehydratase, subunit alpha (hgdA)	30.5%
<i>Heme and porphyrin</i>			<i>Chaperones</i>		
AF1648	bacteriochlorophyll synthase, 33 kDa subunit	27.9%	AF1296	small heat shock protein (hsp20-1)	52.3%
AF0464	bacteriochlorophyll synthase, 43 kDa subunit (chp-1)	25.7%	AF1971	small heat shock protein (hsp20-2)	38.1%
AF1023	bacteriochlorophyll synthase, 43 kDa subunit (chp-2)	31.2%	AF2238	thermosome, subunit alpha (thsA)	70.6%
AF1637	bacteriochlorophyll synthase, 43 kDa subunit (chp-3)	27.0%	AF1451	thermosome, subunit beta (thsB)	68.2%
AF0037	cobalamin (5'-phosphate) synthase (cobS-1)	33.9%	<i>Chromosome-associated protein</i>		
AF2323	cobalamin (5'-phosphate) synthase (cobS-2)	34.4%	AF0337	archaeal histone A1 (hpyA1-1)	64.6%
AF0725	cobalamin biosynthesis precorrin methylase (cbiG)	30.7%	AF1493	archaeal histone A1 (hpyA1-2)	69.7%
AF0727	cobalamin biosynthesis precorrin-2 methyltransferase (cbiL)	31.5%	<i>Detoxification</i>		
AF0726	cobalamin biosynthesis precorrin-3 methylase (cbiF)	49.2%	AF2171	2-nitropropane dioxygenase (ncd2)	39.7%
AF0724	cobalamin biosynthesis precorrin-3 methylase (cbiH)	49.0%	AF0270	alkyl hydroperoxide reductase	73.5%
			AF1361	arsenate reductase (arsC)	40.5%
			AF0550	N-ethylmaleimide chlorohydrolase (trza-1)	45.9%
			AF0987	N-ethylmaleimide chlorohydrolase (trza-2)	44.5%
			AF0254	NADH oxidase (noxA-1)	35.1%
			AF0395	NADH oxidase (noxA-2)	35.5%
			AF0400	NADH oxidase (noxA-3)	40.8%
			AF0951	NADH oxidase (noxA-4)	36.7%
			AF1858	NADH oxidase (noxA-5)	34.0%
			AF0485	NADH oxidase (noxB-1)	43.3%
			AF1282	NADH oxidase (noxB-2)	42.9%
			AF0226	NADH oxidase (noxC)	38.4%
			AF0515	NADH oxidase, putative	25.5%
			AF2233	peroxidase / catalase (perA)	62.9%
			<i>Protein and peptide secretion</i>		
			AF1902	protein translocase, subunit SEC61 alpha (secY)	50.0%
			AF0536	protein translocase, subunit SEC61 gamma (secE)	25.0%
			AF2062	signal recognition particle receptor (dps)	54.8%
			AF1258	signal recognition particle, subunit SRP19 (srp19)	36.6%
			AF0622	signal recognition particle, subunit SRP54 (srp54)	51.2%
			AF1791	signal sequence peptidase (sec11)	36.3%
			AF1657	signal sequence peptidase (spc21)	47.0%
			AF1655	signal sequence peptidase, putative	34.5%
			AF0338	type II secretion system protein (gspE-1)	38.5%
			AF0659	type II secretion system protein (gspE-2)	38.2%
			AF0996	type II secretion system protein (gspE-3)	41.7%
			AF1049	type II secretion system protein (gspE-4)	46.5%
<i>CENTRAL INTERMEDIARY METABOLISM</i>			<i>Degradation of polysaccharides</i>		
			AF1207	2-deoxy-D-glucuronate 3-dehydrogenase (kduD)	45.3%
			AF1795	endoglucanase (celM)	55.4%
<i>Phosphorus compounds</i>			<i>Phosphorus compounds</i>		
			AF0756	exopolysphatase (ppx1)	55.1%
<i>Polyamine biosynthesis</i>			<i>Polyamine biosynthesis</i>		
			AF0646	agmatinase (speB)	33.3%
			AF2334	spermidine synthase (speE)	37.1%
<i>Polysaccharides - (cytoplasmic)</i>			<i>Polysaccharides - (cytoplasmic)</i>		
			AF0599	dolichol phosphate mannose synthase, putative	32.1%
<i>Sulfur metabolism</i>			<i>Sulfur metabolism</i>		
			AF		

AF1957	2-hydroxyglutaryl-CoA dehydratase, subunit beta (hgdB)	24.4%	AF0499	molycoprotein oxidoreductase, iron-sulfur binding subunit	41.5%	TCa cycle				
AF0130	acetylpolymine aminohydrolase (aphA)	38.7%	AF0500	molycoprotein oxidoreductase, membrane subunit	27.9%	AF1963	aconitase (acn)	57.1%		
AF2290	acetylpolymine aminohydrolase, putative	33.3%	AF1202	molycoprotein oxidoreductase, iron-sulfur binding subunit	35.5%	AF1340	citrate synthase (citZ)	50.3%		
AF0991	glutaryl-CoA dehydrogenase (gcdH)	48.7%				AF1098	fumarase (fum-1)	49.1%		
AF1323	group II decarboxylase	28.0%	AF1203	molycoprotein oxidoreductase, molycoprotein binding subunit	30.1%	AF1099	fumarase (fum-2)	53.4%		
AF2004	group II decarboxylase	46.1%				AF0647	isocitrate dehydrogenase, NADP (icd)	57.2%		
AF2295	group II decarboxylase	30.5%	AF2384	molycoprotein oxidoreductase, molycoprotein binding subunit	34.6%	AF1727	malate oxidoreductase (mae)	52.3%		
AF1665	ornithine cyclodeaminase (arcB)	35.3%				AF0681	succinate dehydrogenase, flavoprotein subunit A (sdhA)	48.2%		
Anaerobic										
AF1145	4-hydroxybutyrate CoA transferase (cat2-1)	46.5%	AF2385	molycoprotein oxidoreductase, iron-sulfur binding subunit	46.9%	AF0682	succinate dehydrogenase, iron-sulfur subunit B (sdhB)	51.3%		
AF1854	4-hydroxybutyrate CoA transferase (cat2-2)	47.5%	AF2386	molycoprotein oxidoreductase, membrane subunit	30.3%	AF0683	succinate dehydrogenase, subunit C (sdhC)	36.6%		
AF0886	glycerol kinase (gpkK)	33.8%	AF0159	molycoprotein oxidoreductase, molycoprotein binding subunit, putative	30.9%	AF1539	succinyl-CoA synthetase, alpha subunit (sucD-1)	56.9%		
AF1328	glycerol-3-phosphate dehydrogenase (gfpA)	27.8%	AF2267	NAD(P)H-flavin oxidoreductase	31.4%	AF2185	succinyl-CoA synthetase, alpha subunit (sucD-2)	63.5%		
AF0871	glycerol-3-phosphate dehydrogenase (NAD(P)+) (gpsA)	36.3%	AF0131	NAD(P)H-flavin oxidoreductase, putative	28.2%	AF1540	succinyl-CoA synthetase, beta subunit (sucC-1)	51.3%		
AF0020	L-carnitine dehydratase (caib-1)	33.3%	AF2352	NADH dehydrogenase, subunit 1, putative	28.9%	AF2186	succinyl-CoA synthetase, beta subunit (sucC-2)	49.6%		
AF0990	L-carnitine dehydratase (caib-2)	31.2%	AF1828	NADH dehydrogenase, subunit 3	24.3%	FATTY ACID AND PHOSPHOLIPID METABOLISM				
ATP-proton motive force interconversion										
AF1158	ATP synthase, subunit E, putative	47.1%	AF0248	NADH-dependent flavin oxidoreductase	36.7%	General				
AF1165	H+-transporting ATP synthase, subunit A (atpA)	67.0%	AF0342	nigerythrin, putative	33.3%	AF1736	3-hydroxy-3-methylglutaryl-coenzyme A reductase (mvsa)	57.1%		
AF1167	H+-transporting ATP synthase, subunit B (atpB)	72.6%	AF0546	nitrate reductase, gamma subunit (narI)	30.1%	AF0017	3-hydroxyacyl-CoA dehydrogenase (hbd-1)	41.1%		
AF1164	H+-transporting ATP synthase, subunit C (atpC)	37.5%	AF0501	nitrate reductase, gamma subunit, putative	29.3%	AF0285	3-hydroxyacyl-CoA dehydrogenase (hbd-2)	55.8%		
AF1168	H+-transporting ATP synthase, subunit D (atpD)	47.1%	AF1126	P450 cytochrome, putative	30.5%	AF0434	3-hydroxyacyl-CoA dehydrogenase (hbd-3)	40.7%		
AF1163	H+-transporting ATP synthase, subunit E (atpE)	36.3%	AF0463	polyferredoxin (mvhB), authentic frameshift	32.2%	AF1025	3-hydroxyacyl-CoA dehydrogenase (hbd-4)	45.6%		
AF1165	H+-transporting ATP synthase, subunit F (atpF)	45.0%	AF1379	quinone-reactive Ni/Fe-hydrogenase B-type cytochrome subunit (hycD)	29.0%	AF1122	3-hydroxyacyl-CoA dehydrogenase (hbd-5)	45.2%		
AF1159	H+-transporting ATP synthase, subunit I (atpI)	30.1%	AF0173	reductase, assembly protein	30.0%	AF1177	3-hydroxyacyl-CoA dehydrogenase (hbd-6)	35.8%		
AF1160	H+-transporting ATP synthase, subunit K (atpK-1)	46.3%	AF0547	reductase, iron-sulfur binding subunit	28.3%	AF1190	3-hydroxyacyl-CoA dehydrogenase (hbd-7)	46.5%		
AF1162	H+-transporting ATP synthase, subunit K (atpK-2)	46.3%	AF0867	reductase, putative	33.3%	AF1206	3-hydroxyacyl-CoA dehydrogenase (hbd-8)	36.3%		
Electron transport										
AF2036	cytochrome C oxidase folding protein (coxD)	33.3%	AF0880	rubredoxin (rd-1)	69.2%	AF2017	3-hydroxyacyl-CoA dehydrogenase (hbd-9)	35.4%		
AF0144	cytochrome C oxidase, subunit II (cbaB)	34.2%	AF1349	rubredoxin (rd-2)	67.9%	AF2273	3-hydroxyacyl-CoA dehydrogenase (hbd-10)	39.4%		
AF0142	cytochrome C oxidase, subunit II, putative	38.0%	AF0832	rubrythrin (rr1)	45.7%	AF0018	3-ketoacyl-CoA thiolase (acaB-1)	41.0%		
AF0190	cytochrome C oxidase, subunit II, putative	31.7%	AF0831	rubrythrin (rr2)	63.7%	AF0034	3-ketoacyl-CoA thiolase (acaB-2)	38.3%		
AF1057	cytochrome C-type biogenesis protein (coaA)	30.7%	AF1640	rubrythrin (rr3)	37.8%	AF0133	3-ketoacyl-CoA thiolase (acaB-3)	32.3%		
AF2192	cytochrome C-type biogenesis protein (nirE)	36.1%	AF2312	rubrythrin (rr4)	41.4%	AF0134	3-ketoacyl-CoA thiolase (acaB-4)	32.5%		
AF2286	cytochrome oxidase, subunit I (cydA-1)	22.9%	AF0711	thioredoxin (trx-1)	28.4%	AF0201	3-ketoacyl-CoA thiolase (acaB-5)	26.9%		
AF2297	cytochrome oxidase, subunit I (cydA-2)	31.5%	AF0769	thioredoxin (trx-2)	38.5%	AF0202	3-ketoacyl-CoA thiolase (acaB-6)	33.5%		
AF2046	cytochrome oxidase, subunit I, putative	25.1%	AF1284	thioredoxin (trx-3)	52.9%	AF0283	3-ketoacyl-CoA thiolase (acaB-7)	42.0%		
AF0528	cytochrome-c3 hydrogenase, subunit gamma	39.3%	AF2144	thioredoxin (trx-4)	48.9%	AF0438	3-ketoacyl-CoA thiolase (acaB-8)	42.4%		
AF0833	desulfoferredoxin (dfr)	63.0%	AF1339	ubiquinol-cytochrome C reductase complex, subunit VI requiring protein	60.9%	AF0967	3-ketoacyl-CoA thiolase (acaB-9)	33.7%		
AF0344	desulfoferredoxin, putative	47.3%	Fermentation					AF0968	3-ketoacyl-CoA thiolase (acaB-10)	28.0%
AF0287	electron transfer flavoprotein, subunit alpha (etfA)	39.7%	AF1779	2-hydroxyacyl dehydrogenase, putative	37.6%	AF1291	3-ketoacyl-CoA thiolase (acaB-11)	40.1%		
AF0286	electron transfer flavoprotein, subunit beta (etfB)	38.8%	AF0469	2-ketoglutarate ferredoxin oxidoreductase, subunit alpha (korA)	52.3%	AF2416	3-ketoacyl-CoA thiolase (acaB-12)	49.9%		
AF1380	F420-nonreducing hydrogenase (vhtA)	34.8%	AF0468	2-ketoglutarate ferredoxin oxidoreductase, subunit beta (korB)	51.2%	AF1028	3-ketoacyl-CoA thiolase (fadA-1)	38.8%		
AF1371	F420-nonreducing hydrogenase (vhtD-1)	30.9%	AF0470	2-ketoglutarate ferredoxin oxidoreductase, subunit delta (korD)	47.2%	AF1197	3-ketoacyl-CoA thiolase (fadA-2)	47.2%		
AF1378	F420-nonreducing hydrogenase (vhtD-2)	33.1%	AF0471	2-ketoglutarate ferredoxin oxidoreductase, subunit gamma (korG)	40.0%	AF2243	3-ketoacyl-CoA thiolase (fadA-3)	40.3%		
AF1824	F420H2-quinone oxidoreductase, 11.2 kDa subunit, putative	24.1%	AF2053	2-ketoglutarate ferredoxin oxidoreductase, subunit alpha (vorA)	41.2%	AF0033	acyl carrier protein synthase (acaA-1)	28.6%		
AF1823	F420H2-quinone oxidoreductase, 16.5 kDa subunit, putative	25.7%	AF2052	2-ketoglutarate ferredoxin oxidoreductase, subunit beta (vorB)	42.7%	AF2415	acyl carrier protein synthase (acaA-2)	58.7%		
AF1832	F420H2-quinone oxidoreductase, 32 kDa subunit (nuoI)	95.5%	AF2054	2-ketoglutarate ferredoxin oxidoreductase, subunit delta (vorD)	51.5%	AF0199	acyl-CoA dehydrogenase (acd-1)	35.9%		
AF1833	F420H2-quinone oxidoreductase, 39 kDa subunit, putative	33.6%	AF2055	2-ketoglutarate ferredoxin oxidoreductase, subunit gamma (vorG)	45.2%	AF0436	acyl-CoA dehydrogenase (acd-2)	44.1%		
AF1829	F420H2-quinone oxidoreductase, 39.7 kDa subunit, putative	43.8%	AF0749	2-oxoacid ferredoxin oxidoreductase, subunit alpha (ora)	33.7%	AF0498	acyl-CoA dehydrogenase (acd-3)	22.9%		
AF1831	F420H2-quinone oxidoreductase, 41.2 kDa subunit, putative	34.8%	AF0750	2-oxoacid ferredoxin oxidoreductase, subunit beta (orb)	49.2%	AF0671	acyl-CoA dehydrogenase (acd-4)	37.9%		
AF1827	F420H2-quinone oxidoreductase, 43.2 kDa subunit, putative	26.9%	AF1286	acetyl-CoA synthetase (acs-1)	35.1%	AF0845	acyl-CoA dehydrogenase (acd-5)	44.8%		
AF1830	F420H2-quinone oxidoreductase, 45 kDa subunit (nuoD)	80.0%	AF0197	acetyl-CoA synthetase (acs-2)	27.1%	AF0984	acyl-CoA dehydrogenase (acd-6)	35.8%		
AF1825	F420H2-quinone oxidoreductase, 53.9 kDa subunit (nuoM)	32.1%	AF0366	acetyl-CoA synthetase (acs-3)	47.3%	AF1026	acyl-CoA dehydrogenase (acd-7)	42.8%		
AF1826	F420H2-quinone oxidoreductase, 72.4 kDa subunit (nuoL)	33.2%	AF0677	acetyl-CoA synthetase (acs-4)	40.9%	AF1141	acyl-CoA dehydrogenase (acd-8)	43.2%		
AF0156	ferredoxin (fdx-1)	45.3%	AF0975	acetyl-CoA synthetase (acs-5)	36.2%	AF1293	acyl-CoA dehydrogenase (acd-9)	45.8%		
AF0166	ferredoxin (fdx-2)	49.2%	AF0976	acetyl-CoA synthetase (acs-6)	34.3%	AF2057	acyl-CoA dehydrogenase (acd-10)	44.6%		
AF0365	ferredoxin (fdx-3)	53.2%	AF1287	acetyl-CoA synthetase (acs-6)	34.3%	AF2244	acyl-CoA dehydrogenase (acd-11)	42.6%		
AF0427	ferredoxin (fdx-4)	56.1%	AF0024	alcohol dehydrogenase, iron-containing	36.2%	AF2275	acyl-CoA dehydrogenase (acd-12)	38.9%		
AF0923	ferredoxin (fdx-5)	56.9%	AF0339	alcohol dehydrogenase, iron-containing	37.4%	AF1175	acyl-CoA dehydrogenase, short chain-specific (acdS)	30.1%		
AF1010	ferredoxin (fdx-6)	44.4%	AF2019	alcohol dehydrogenase, iron-containing	35.7%	AF0818	acylphosphatase (acyP)	36.8%		
AF1239	ferredoxin (fdx-7)	29.0%	AF2389-C	acetyl-CoA synthetase, putative	64.8%	AF0868	alkylidihydroxyacetonephosphate synthase	33.6%		
AF2142	ferredoxin (fdx-8)	38.0%	AF2389-N	acetyl-CoA synthetase, putative	59.3%	AF2286	bifunctional short chain isoprenyl diphosphate synthase (idsA)	42.7%		
AF0164	ferredoxin-nitrite reductase (nirA)	29.7%	AF2101	alcohol dehydrogenase, zinc-dependent	34.8%	AF0220	biotin carboxylase (acc)	52.1%		
AF2332	flavodoxin, putative	30.3%	AF0023	aldehyde ferredoxin oxidoreductase (aor-1)	41.1%	AF0865	carboxylesterase (est-1)	27.1%		
AF0167	flavoprotein (fprA-1)	33.2%	AF0077	aldehyde ferredoxin oxidoreductase (aor-2)	32.6%	AF1537	carboxylesterase (est-2)	29.0%		
AF1520	flavoprotein (fprA-2)	47.2%	AF0340	aldehyde ferredoxin oxidoreductase (aor-3)	38.4%	AF2336	carboxylesterase (est-3)	30.4%		
AF0557	flavoprotein reductase	26.2%	AF2281	aldehyde ferredoxin oxidoreductase (aor-4)	53.0%	AF1716	carboxylesterase (estA)	40.4%		
AF1463	fumarate reductase, flavoprotein subunit (fdrA)	27.0%	AF0006	corrinoid methyltransferase protein (mtaC-1)	30.7%	AF1744	CDP-diacylglycerol-glycerol-3-phosphate 3-phosphatidyltransferase (pgsA-2)	26.7%		
AF1536	glutaredoxin (grx-1)	34.3%	AF0001	corrinoid methyltransferase protein (mtaC-2)	29.5%	AF1143	CDP-diacylglycerol-glycerol-3-phosphate-3-phosphatidyltransferase (pgsA-1)	27.0%		
AF2145	glutaredoxin (grx-2)	38.8%	AF0394	D-formate dehydrogenase, cytochrome-type (dld)	31.9%	AF2044	CDP-diacylglycerol-serine O-phosphatidyltransferase (pssa)	36.6%		
AF0663	heterodisulfide reductase, subunit A (hdrA-1)	42.2%	AF0560	formate dehydrogenase (fdhH1), authentic frameshift	32.9%	AF0435	enoyl-CoA hydratase (fad-1)	47.6%		
AF1377	heterodisulfide reductase, subunit A (hdrA-2)	46.8%	AF1199	glutamate CoA-transferase, subunit A (gctA)	31.9%	AF0885	enoyl-CoA hydratase (fad-2)	59.1%		
AF0662	heterodisulfide reductase, subunit A/		AF1198	glutamate CoA-transferase, subunit B (gctB)	37.0%	AF1641	enoyl-CoA hydratase (fad-4)	41.7%		
AF1238	heterodisulfide reductase, subunit A/methylviologen reducing hydrogenase, subunit delta	34.2%	AF1489	indolepyruvate ferredoxin oxidoreductase, subunit alpha (iorA)	48.1%	AF2429	enoyl-CoA hydratase (fad-5)	34.7%		
	heterodisulfide reductase, subunit delta		AF2030	indolepyruvate ferredoxin oxidoreductase, subunit beta (iorB)	41.1%	AF1763	lipase, putative	33.5%		
AF1375	heterodisulfide reductase, subunit B (hdrB)	36.0%	AF0807	L-lactate dehydrogenase, cytochrome-type (lldD)	39.4%	AF0089	long-chain-fatty-acyl-CoA ligase (fadD-1)	31.9%		
AF0271	heterodisulfide reductase, subunit B, putative	35.3%	AF0885	L-malate dehydrogenase, NAD+-dependent (mdhA)	40.1%	AF0200	long-chain-fatty-acyl-CoA ligase (fadD-2)	34.8%		
AF1376	heterodisulfide reductase, subunit C (hdrC)	33.3%	AF2085	oxaloacetate decarboxylase, biotin carboxyl carrier subunit, putative	38.7%	AF0687	long-chain-fatty-acyl-CoA ligase (fadD-3)	31.1%		
AF0502	heterodisulfide reductase, subunit D, putative	33.8%	AF2084	oxaloacetate decarboxylase, sodium ion pump subunit (oadB)	59.8%	AF0840	long-chain-fatty-acyl-CoA ligase (fadD-4)	38.1%		
AF0809	heterodisulfide reductase, subunit D, putative	100.0%	AF1252	oxaloacetate decarboxylase, subunit alpha (oadA)	63.3%	AF1029	long-chain-fatty-acyl-CoA ligase (fadD-5)	37.8%		
AF0661	heterodisulfide reductase, subunit E, putative	23.8%	AF1701	pyruvate ferredoxin oxidoreductase, subunit alpha (porA)	50.3%	AF1772	long-chain-fatty-acyl-CoA ligase (fadD-6)	36.0%		
AF0755	heterodisulfide reductase, subunits E and D, putative	31.8%				AF1932	long-chain-fatty-acyl-CoA ligase (fadD-7)	31.0%		
AF0506	iron-sulfur binding reductase	38.5%				AF2368	long-chain-fatty-acyl-CoA ligase (fadD-8)	38.7%		
AF1773	iron-sulfur binding reductase	33.3%				AF1753	lysophospholipase	33.5%		
AF1998	iron-sulfur binding reductase	29.6%				AF0196	medium-chain acyl-CoA ligase (alkK-1)	34.6%		
AF0627	iron-sulfur cluster binding protein	45.5%				AF0262	medium-chain acyl-CoA ligase (alkK-2)	38.6%		
AF0688	iron-sulfur cluster binding protein	44.8%				AF0672	medium-chain acyl-CoA ligase (alkK-3)	31.0%		
AF1153	iron-sulfur cluster binding protein	27.9%				AF1261	medium-chain acyl-CoA ligase (alkK-4)	42.7%		
AF1185	iron-sulfur cluster binding protein	36.7%				AF2033	medium-chain acyl-CoA ligase (alkK-5)	33.5%		
AF1263	iron-sulfur cluster binding protein	42.1%				AF2289	mevalonate kinase (mvk)	40.8%		
AF2380	iron-sulfur cluster binding protein	36.3%				AF1784	myo-inositol 1-phosphate synthase (ino1)	32.2%		
AF2381	iron-sulfur cluster binding protein	34.4%				AF2045				

AF1935	N5,N10-methylenetetrahydromethanopterin cyclohydrolase (mch)	97.3%	AF0004	RNase I inhibitor	54.5%	AF0633	isoleucyl-tRNA synthetase (ileS)	48.9%		
AF0714	N5,N10-methylenetetrahydromethanopterin dehydrogenase (mtd)	61.8%	AF0021	signal-transducing histidine kinase	26.1%	AF2421	leucyl-tRNA synthetase (leuS)	49.7%		
AF1066	N5,N10-methylenetetrahydromethanopterin reductase (mer-1)	59.1%	AF0208	signal-transducing histidine kinase	27.9%	AF1216	lysyl-tRNA synthetase (lysS)	43.6%		
AF1196	N5,N10-methylenetetrahydromethanopterin reductase (mer-2)	37.4%	AF0450	signal-transducing histidine kinase	32.4%	AF1453	methionyl-tRNA synthetase (metS)	45.2%		
AF0009	N5-methyltetrahydromethanopterin:coenzyme M methyltransferase (mtm)	42.1%	AF0770	signal-transducing histidine kinase	26.9%	AF1955	phenylalanyl-tRNA synthetase, subunit alpha (pheS)	44.4%		
AF1587	ribulose biphosphate carboxylase, large subunit (rbcL-1)	40.6%	AF0893	signal-transducing histidine kinase	28.7%	AF1424	phenylalanyl-tRNA synthetase, subunit beta (pheT)	42.6%		
AF1638	ribulose biphosphate carboxylase, large subunit (rbcL-2)	44.9%	AF1184	signal-transducing histidine kinase	29.8%	AF1609	prolyl-tRNA synthetase (proS)	56.8%		
AF1930	tungsten formylmethanofuran dehydrogenase, subunit A (fwdA)	48.9%	AF1452	signal-transducing histidine kinase	28.5%	AF2035	seryl-tRNA synthetase (serS)	45.4%		
AF1650	tungsten formylmethanofuran dehydrogenase, subunit B (fwdB-1)	37.0%	AF1467	signal-transducing histidine kinase	37.4%	AF0548	threonyl-tRNA synthetase (thrS)	46.3%		
AF1929	tungsten formylmethanofuran dehydrogenase, subunit B (fwdB-2)	49.4%	AF1472	signal-transducing histidine kinase	30.4%	AF1694	tyrosyl-tRNA synthetase (tyrS)	52.4%		
AF1931	tungsten formylmethanofuran dehydrogenase, subunit C (fwdC)	44.1%	AF1515	signal-transducing histidine kinase	32.0%	AF0776	tyrosyl-tRNA synthetase (tyrS)	57.6%		
AF1651	tungsten formylmethanofuran dehydrogenase, subunit D (fwdD-1)	32.6%	AF1639	signal-transducing histidine kinase	29.9%	AF2224	valyl-tRNA synthetase (valS)	54.5%		
AF1928	tungsten formylmethanofuran dehydrogenase, subunit D (fwdD-2)	52.6%	AF1721	signal-transducing histidine kinase	34.5%	Degradation of proteins, peptides, and glycopeptides				
AF0177	tungsten formylmethanofuran dehydrogenase, subunit E (fwdE)	29.7%	AF2109	signal-transducing histidine kinase	31.6%	AF1976	26S protease regulatory subunit 4	66.0%		
AF1644	tungsten formylmethanofuran dehydrogenase, subunit F (fwdF)	38.2%	AF0881	signal-transducing histidine kinase, authentic frameshift	26.5%	AF1653	alkaline serine protease (aprM)	44.5%		
AF1649	tungsten formylmethanofuran dehydrogenase, subunit G (fwdG)	45.6%	AF0277	signal-transducing histidine kinase, putative	29.8%	AF0578	aminopeptidase, putative	27.9%		
PURINES, PYRIMIDINES, NUCLEOSIDES, AND NUCLEOTIDES					AF0410	signal-transducing histidine kinase, putative	27.1%	AF0364	ATP-dependent protease La (lon)	36.6%
2'-Deoxyribonucleotide metabolism					AF0448	signal-transducing histidine kinase, putative	26.1%	AF1946	cysteine proteinase, putative	36.2%
AF1108	deoxycytidine triphosphate deaminase, putative	38.1%	AF1482	signal-transducing histidine kinase, putative	25.2%	AF1281	intracellular protease (ppl)	56.0%		
AF1664	ribonucleotide reductase (nrd)	59.7%	AF0202	signal-transducing histidine kinase, putative	22.5%	AF1112	O-sialoglycoprotein endopeptidase (gcp)	57.6%		
AF1554	thioredoxin reductase (trxB)	45.2%	AF2420	signal-transducing histidine kinase, putative	28.4%	AF0665	O-sialoglycoprotein endopeptidase, putative	35.8%		
AF0247	thymidylate synthase, putative	33.1%	AF0442	succinoyl-glycan biosynthesis regulator (exsB)	37.2%	AF2086	protease inhibitor, putative	37.0%		
Nucleotide and nucleoside interconversions					AF1516	sugar fermentation stimulation protein (sfsA)	31.0%	AF0490	proteasome, subunit alpha (psmA)	60.8%
AF0876	5'-nucleotidase (ntf)	30.9%	AF1270	transcriptional regulatory protein, ArsR family	35.4%	AF0481	proteasome, subunit beta (psmB)	58.3%		
AF0676	adenylate kinase (adk)	56.1%	AF1544	transcriptional regulatory protein, ArsR family	32.3%	AF2034	X-pro aminopeptidase (pepQ)	34.6%		
AF1900	cytidylate kinase (cmk)	48.6%	AF1853	transcriptional regulatory protein, ArsR family	34.9%	Protein modification				
AF0767	nucleoside diphosphate kinase (ndk)	56.4%	AF2136	transcriptional regulatory protein, ArsR family	39.8%	AF0656	antibiotic maturation protein (pmbA)	32.7%		
AF0061	thymidylate kinase (tmk)	34.9%	AF0439	transcriptional regulatory protein, AsnC family	29.8%	AF0378	CODH nickel-insertion accessory protein (cocC-1)	35.7%		
AF1308	thymidylate kinase, putative	26.3%	AF0474	transcriptional regulatory protein, AsnC family	51.0%	AF1685	CODH nickel-insertion accessory protein (cocC-2)	47.4%		
AF2042	uridylate kinase (pyrH)	53.6%	AF0584	transcriptional regulatory protein, AsnC family	35.3%	AF1615	cofactor modifying protein (cmo)	27.2%		
Purine ribonucleotide biosynthesis					AF1121	transcriptional regulatory protein, AsnC family	35.8%	AF2195	deoxyhypusine synthase (dys1-1)	32.6%
AF2242	adenylosuccinate lyase (purB)	52.3%	AF1404	transcriptional regulatory protein, AsnC family	32.6%	AF2200	deoxyhypusine synthase (dys1-2)	34.9%		
AF0841	adenylosuccinate synthetase (purA)	70.8%	AF1448	transcriptional regulatory protein, AsnC family	30.6%	AF0381	diphthine synthase (dphS)	40.8%		
AF0873	amidophosphoribosyltransferase (purF)	55.8%	AF1223	transcriptional regulatory protein, AsnC family	46.4%	AF2324	fmu and frmv protein	40.0%		
AF0253	GMP synthase (guaA-1)	59.8%	AF1743	transcriptional regulatory protein, AsnC family	34.9%	AF1367	hydrogenase expression/formation protein (hvpA)	40.4%		
AF1320	GMP synthase (guaA-2)	49.4%	AF2127	transcriptional regulatory protein, LysR family	30.8%	AF1368	hydrogenase expression/formation protein (hvpB)	54.4%		
AF1811	inosine monophosphate cyclohydrolase	38.3%	AF0114	transcriptional regulatory protein, putative	35.6%	AF1369	hydrogenase expression/formation protein (hvpC)	40.5%		
AF0847	inosine monophosphate dehydrogenase (guaB-1)	41.6%	AF1998	transcriptional regulatory protein, Rok family	32.9%	AF1370	hydrogenase expression/formation protein (hvpD)	46.0%		
AF2118	inosine monophosphate dehydrogenase (guaB-2)	31.9%	AF1112	transcriptional regulatory protein, Sir2 family	38.9%	AF1365	hydrogenase expression/formation protein (hvpE)	51.5%		
AF1259	inosine monophosphate dehydrogenase, putative	51.6%	AF1676	transcriptional regulatory protein, Sir2 family	40.6%	AF1366	hydrogenase expression/formation regulatory protein (hvpF)	45.1%		
AF1157	phosphoribosylamino-glycine lyase (purD)	40.9%	AF1817	transcriptional regulatory protein, TetR family	24.5%	AF0036	L-isocaparyl protein carboxyl methyltransferase (pcm-1)	60.7%		
AF1271	phosphoribosylaminoimidazole carboxylase (purE)	42.8%	AF0363	transcriptional repressor (cinR)	27.5%	AF2322	L-isocaparyl protein carboxyl methyltransferase (pcm-2)	59.3%		
AF1272	phosphoribosylaminoimidazolesuccinocarboxamide synthase (purC)	34.7%	REPLICATION					AF1840	methionyl aminopeptidase (map)	48.6%
AF1693	phosphoribosylformylglycinamide cyclo-lyase (purM)	53.8%	AF2117	3-methyladenine DNA glycosylase (alkA)	30.0%	AF1989	peptidyl-prolyl cis-trans isomerase (slyD)	34.4%		
AF1280	phosphoribosylformylglycinamide synthase I (purQ)	40.9%	AF2060	activator 1, replication factor C, 35 KDa subunit	66.3%	AF0853	proliferating-cell nuclear antigen P120, putative	35.7%		
AF1940	phosphoribosylformylglycinamide synthase II (purL)	41.5%	AF1195	activator 1, replication factor C, 53 KDa subunit	43.7%	AF2039	proliferating-cell nuclear antigen P120, putative	44.2%		
AF0589	ribose-phosphate pyrophosphokinase (prsA-1)	35.0%	AF0465	DNA gyrase, subunit A (gyrA)	48.4%	AF1449	pyruvate formate-lyase 2 (pflD)	37.8%		
AF1419	ribose-phosphate pyrophosphokinase (prsA-2)	41.1%	AF0530	DNA gyrase, subunit B (gyrB)	58.4%	AF1450	pyruvate formate-lyase 2 activating enzyme (pflC)	38.8%		
Pyrimidine ribonucleotide biosynthesis					AF1388	DNA helicase, putative	46.8%	AF0117	pyruvate formate-lyase activating enzyme (act-1)	25.5%
AF0106	aspartate carbamoyltransferase, catalytic subunit (pyrB)	60.7%	AF1960	DNA helicase, putative	32.7%	AF0918	pyruvate formate-lyase activating enzyme (act-2)	42.3%		
AF0107	aspartate carbamoyltransferase, regulatory subunit (pyrI)	48.2%	AF0623	DNA ligase (lig)	44.4%	AF1330	pyruvate formate-lyase activating enzyme (act-3)	45.8%		
AF1274	carbamoyl-phosphate synthase, large subunit (carB)	65.1%	AF1725	DNA ligase, putative	32.7%	AF2278	pyruvate formate-lyase activating enzyme (act-4)	42.5%		
AF1273	carbamoyl-phosphate synthase, small subunit (carA)	55.2%	AF0497	DNA polymerase B1 (polB)	45.1%	AF1961	pyruvate formate-lyase activating enzyme (pflX)	50.2%		
AF0252	CTP synthase (pyrG)	58.3%	AF0693	DNA polymerase B2 (boxA), authentic frameshift	32.3%	AF0380	transmembrane oligosaccharyl transferase, putative	27.8%		
AF2250	dihydroorotase (pyrC)	37.2%	AF0972	DNA polymerase III, subunit epsilon (dnaQ)	31.9%	AF0329	transmembrane oligosaccharyl transferase, putative	29.3%		
AF0745	dihydroorotase dehydrogenase (pyrD)	44.8%	AF2277	DNA polymerase, bacteriophage-type	36.9%	Ribosomal proteins: synthesis and modification				
AF1741	orate phosphoribosyl transferase (pyrE)	49.0%	AF0742	DNA primase, putative	26.8%	AF1490	LSU ribosomal protein L1P (rpL1P)	48.6%		
AF0386	orate phosphoribosyl transferase, putative	39.0%	AF0264	DNA repair protein RAD2 (rad2)	44.4%	AF1922	LSU ribosomal protein L2P (rpL2P)	60.4%		
Salvage of nucleosides and nucleotides					AF0358	DNA repair protein RAD25	32.5%	AF1925	LSU ribosomal protein L3P (rpL3P)	56.5%
AF0240	adenine deaminase (adeC)	39.5%	AF1031	DNA repair protein RAD32 (rad32)	37.6%	AF1924	LSU ribosomal protein L4P (rpL4P)	56.4%		
AF1764	dCMP deaminase, putative	39.0%	AF0363	DNA repair protein RAD51 (radA)	59.3%	AF1912	LSU ribosomal protein L5P (rpL5P)	51.7%		
AF1788	methylothioadenosine phosphorylase (mtaP)	40.0%	AF2118	DNA repair protein, putative	28.9%	AF1909	LSU ribosomal protein L6P (rpL6P)	53.7%		
AF1341	thymidine phosphorylase (deoA-1)	46.7%	AF1806	DNA topoisomerase I (topA)	36.2%	AF0764	LSU ribosomal protein L7AE (rpL7AE)	60.7%		
AF1342	thymidine phosphorylase (deoA-2)	40.7%	AF0940	DNA topoisomerase VI, subunit A (top6A)	39.8%	AF1491	LSU ribosomal protein L10E (rpL10E)	45.6%		
AF0239	xanthine-guanine phosphoribosyltransferase (gptA-1)	25.7%	AF0652	DNA topoisomerase VI, subunit B (top6B)	43.9%	AF0538	LSU ribosomal protein L11P (rpL11P)	67.8%		
AF1789	xanthine-guanine phosphoribosyltransferase (gptA-2)	28.2%	AF1692	endonuclease III (ntf)	44.3%	AF1492	LSU ribosomal protein L12A (rpL12A)	76.0%		
REGULATORY FUNCTIONS					AF0580	exodeoxyribonuclease III (xthA)	41.3%	AF1258	LSU ribosomal protein L13P (rpL13P)	47.4%
AF1959	(R)-hydroxylglutaryl-CoA dehydratase activator (hgdC)	51.2%	AF2314	methylated-DNA-protein-cysteine methylation transferase (ogt)	55.3%	AF1915	LSU ribosomal protein L14P (rpL14P)	66.7%		
AF0168	arsenical resistance operon repressor, putative	36.7%	AF1409	modification methylase, type III R/M system	31.4%	AF2319	LSU ribosomal protein L15E (rpL15E)	70.3%		
AF2204	arylsulfatase regulatory protein, putative	29.9%	AF1234	mutator protein MutT (mutT)	63.6%	AF1903	LSU ribosomal protein L15P (rpL15P)	53.8%		
AF0074	biotin operon repressor/biotin-[acetyl CoA carboxylase] ligase (birA)	36.6%	AF2200	mutator protein MutT, putative	42.0%	AF1127	LSU ribosomal protein L18E (rpL18E)	53.8%		
AF1724	dinitrogenase reductase activating glycohydrolase (draG)	37.9%	AF0335	proliferating-cell nuclear antigen (pou30)	33.7%	AF1906	LSU ribosomal protein L18P (rpL18P)	57.8%		
AF2232	ferric uptake regulation protein (fur)	25.8%	AF0694	replication control protein A, putative	30.2%	AF1907	LSU ribosomal protein L19E (rpL19E)	55.5%		
AF1785	iron-dependent repressor	42.0%	AF1024	reverse gyrase (top-RG)	40.7%	AF1529	LSU ribosomal protein L21E (rpL21E)	53.2%		
AF2385	iron-dependent repressor	40.0%	AF0621	ribonuclease HII (mhb)	39.3%	AF1920	LSU ribosomal protein L22P (rpL22P)	55.2%		
AF0245	iron-dependent repressor (desR)	28.2%	AF1715	type I restriction-modification enzyme, M subunit, authentic frameshift	63.0%	AF1923	LSU ribosomal protein L23P (rpL23P)	55.6%		
AF1984	iron-dependent repressor (troR)	28.3%	AF1708	type I restriction-modification enzyme, R subunit	38.2%	AF0537	LSU ribosomal protein L24A (rpL24A)	51.4%		
AF2430	lacZ expression regulatory protein (lcc)	29.6%	AF1710	type I restriction-modification enzyme, S subunit	33.0%	AF0766	LSU ribosomal protein L24E (rpL24E)	61.1%		
AF1622	leucine responsive regulatory protein (lrp)	29.1%	TRANSCRIPTION					AF1914	LSU ribosomal protein L24P (rpL24P)	57.8%
AF0673	mercuric resistance operon regulatory protein (merR)	37.6%	DNA-dependent RNA polymerase					AF1918	LSU ribosomal protein L29P (rpL29P)	44.6%
AF2425	methanol dehydrogenase regulatory protein (mxoR)	48.3%	AF1888	DNA-directed RNA polymerase, subunit A' (rhoA1)	63.6%	AF1890	LSU ribosomal protein L30E (rpL30E)	41.7%		
AF1475	mitochondrial benzodiazepine receptor/sensory transduction protein	38.4%	AF1889	DNA-directed RNA polymerase, subunit A'' (rhoA2)	55.7%	AF1904	LSU ribosomal protein L30P (rpL30P)	55.9%		
AF0198	monoamine oxidase regulatory protein, putative	31.7%	AF1887	DNA-directed RNA polymerase, subunit B' (rhoB1)	65.3%	AF2066	LSU ribosomal protein L31E (rpL31E)	50.6%		
AF1933	monoamine oxidase regulatory protein, putative	46.9%	AF1886	DNA-directed RNA polymerase, subunit B'' (rhoB2)	57.1%	AF1908	LSU ribosomal protein L32E (rpL32E)	51.2%		
AF0978	nitrogen regulatory protein P-II (glnB-1)	61.7%	AF2282	DNA-directed RNA polymerase, subunit D (rhoD)	34.6%	AF0057	LSU ribosomal protein L37AE (rpL37AE)	47.8%		
AF1747	nitrogen regulatory protein P-II (glnB-2)	58.0%	AF1117	DNA-directed RNA polymerase, subunit E' (rhoE1)	48.4%	AF0874	LSU ribosomal protein L37E (rpL37E)	57.9%		
AF1750	nitrogen regulatory protein P-II (glnB-3)	60.7%	AF1166	DNA-directed RNA polymerase, subunit E'' (rhoE2)	40.0%	AF2067	LSU ribosomal protein L39E (rpL39E)	56.9%		
AF0331	pheromone shutdown protein (traB)	40.5%	AF1885	DNA-directed RNA polymerase, subunit H (rhoH)	59.5%	AF1430	LSU ribosomal protein L40E (rpL40E)	73.3%		
AF1797	phosphate regulatory protein, putative	30.7%	AF1131	DNA-directed RNA polymerase, subunit K (rhoK)	61.5%	AF1938	LSU ribosomal protein L44E (rpL44E)	46.8%		
AF0521	protease synthase and sporulation regulator Pai1, putative	52.4%	AF0207	DNA-directed RNA polymerase, subunit L (rhoL)	42.0%	AF2064	LSU ribosomal protein LXA (rpLXA)	53.8%		
AF1627	repressor protein	59.1%	AF1130	DNA-directed RNA polymerase, subunit N (rhoN)	56.8%	AF0739	ribosomal protein S18 alanine acetyltransferase	38.5%		
AF1793	repressor protein	54.5%	Transcription factors					AF2303	ribosomal protein S6 modification protein (rimK)	32.2%
AF0449	response regulator	38.1%	AF1813	TBP-interacting protein TIP49	45.7%	AF1133	SSU ribosomal protein S2P (rpS2P)	58.3%		
AF1063	response regulator	36.3%	AF1299	transcription initiation factor IIB	60.4%	AF1919	SSU ribosomal protein S3P (rpS3P)	50.0%		
AF1256	response regulator	42.5%	AF0							

AF2328	Glu-tRNA amidotransferase, subunit C (gatC)	35.1%	AF1768	protein (dppA)	33.1%	AF2258	multidrug resistance protein	31.3%
AF0815	N ₂ ,N ₂ -dimethylguanosine tRNA methyltransferase (trm1)	38.2%	AF1769	dipeptide ABC transporter, permease protein (dppB)	39.3%	OTHER CATEGORIES		
AF1730	pseudouridylylase synthase I (truA)	37.4%	AF0680	glutamine ABC transporter, permease protein (dppC)	40.8%	<i>Adaptations and atypical conditions</i>		
AF1485	queuine tRNA-ribosyltransferase (tgtB)	44.1%	AF0231	glutamine ABC transporter, periplasmic glutamine-binding protein (glnH)	38.0%	AF0608	ethylene-inducible protein	74.5%
AF0493	ribonuclease PH (rph)	30.8%	AF0232	glutamine ABC transporter, permease protein (glnP)	39.3%	AF0235	heat shock protein (htpK)	32.9%
AF0900	tRNA intron endonuclease (endA)	41.8%	AF0981	osmoprotection protein (proV-1)	39.0%	AF0942	surE stationary-phase survival protein (surE)	50.2%
AF2156	tRNA nucleotidyltransferase (cca)	43.9%	AF0979	osmoprotection protein (proW-1)	32.8%	AF1996	virulence associated protein C (vapC-1)	50.0%
<i>Translation factors</i>			AF0980	osmoprotection protein (proW-2)	36.8%	AF1690	virulence associated protein C (vapC-2)	30.0%
AF2350	ATP-dependent RNA helicase HepA, putative	31.5%	AF0882	osmoprotection protein (proX)	28.7%	<i>Drug and analog sensitivity</i>		
AF2254	ATP-dependent RNA helicase, DEAD-family (deaD)	52.2%	AF0015	proline permease (putP-1)	26.2%	AF1884	daunorubicin resistance ATP-binding protein (drrA)	47.1%
AF0071	ATP-dependent RNA helicase, putative	29.6%	AF0969	proline permease (putP-2)	27.4%	AF1883	daunorubicin resistance membrane protein (drrB)	27.0%
AF1458	ATP-dependent RNA helicase, putative	48.1%	AF1222	proline permease (putP-3)	27.0%	AF0487	penicillin G acylase	31.7%
AF2406	ATP-dependent RNA helicase, putative	35.2%	AF1608	spermidine/putrescine ABC transporter, ATP-binding protein (potA)	50.2%	AF1214	phenylacrylic acid decarboxylase (pad1)	43.2%
AF1149	large helicase-related protein (lhr-1)	34.5%	AF1605	spermidine/putrescine ABC transporter, periplasmic spermidine/putrescine-binding protein (potD), authentic frameshift	31.0%	AF2194	rRNA (adenine-N ⁶)-methyltransferase, putative	23.2%
AF2177	large helicase-related protein (lhr-2), authentic frameshift	56.0%	AF1607	spermidine/putrescine ABC transporter, permease protein (potB)	38.0%	AF1696	small multidrug export protein (qacE)	39.0%
AF1220	peptide chain release factor eRF, subunit 1	51.2%	AF1606	spermidine/putrescine ABC transporter, permease protein (potC)	38.7%	<i>Transposon-related functions</i>		
AF2445	SK12-family helicase, authentic frameshift	45.7%	<i>Anions</i>			AF0120	insertion sequence ISH S1, authentic frameshift	34.5%
AF0937	translation elongation factor EF-1, subunit alpha (tuf)	74.4%	AF2308	arsenite transport protein (arsB)	27.3%	AF0193	ISA0963-1, putative transposase, authentic frameshift	34.3%
AF0574	translation elongation factor EF-1, subunit beta	31.3%	AF1415	chloride channel, putative	27.3%	AF0309	ISA0963-2, putative transposase	33.5%
AF1894	translation elongation factor EF-2 (fus)	62.5%	AF0025	cyanoate transport protein (cynX)	24.5%	AF1310	ISA0963-3, putative transposase	33.5%
AF0777	translation initiation factor eIF-1A (eif1A)	57.5%	AF0087	nitrate ABC transporter, ATP-binding protein (nrtC-1)	47.4%	AF1383	ISA0963-4, putative transposase	33.5%
AF0527	translation initiation factor eIF-2, subunit alpha (eif2A)	51.1%	AF0638	nitrate ABC transporter, ATP-binding protein (nrtC-2)	55.5%	AF1410	ISA0963-5, putative transposase	33.5%
AF2326	translation initiation factor eIF-2, subunit beta, putative	45.5%	AF0640	nitrate ABC transporter, ATP-binding protein, putative	32.5%	AF1706	ISA0963-6, putative transposase	33.5%
AF0592	translation initiation factor eIF-2, subunit gamma (eif2G)	64.4%	AF0086	nitrate ABC transporter, permease protein (nrtB-1)	35.4%	AF1836	ISA0963-7, putative transposase, authentic frameshift	23.2%
AF0370	translation initiation factor eIF-2B, subunit delta (eif2BD)	53.3%	AF0639	nitrate ABC transporter, permease protein (nrtB-2)	37.4%	AF0678	ISA1083-1, ISORF2	33.6%
AF2037	translation initiation factor eIF-2B, subunit delta (eif2BD)	57.9%	AF1359	phosphate ABC transporter, ATP-binding protein (pstB)	66.0%	AF0679	ISA1083-1, putative transposase	37.2%
AF0645	translation initiation factor eIF-5A (eif5A)	50.4%	AF1356	phosphate ABC transporter, periplasmic phosphate-binding protein (phoX)	25.1%	AF1351	ISA1083-2, ISORF2	30.8%
AF0768	translation initiation factor IF-2 (infB)	52.2%	AF1358	phosphate ABC transporter, permease protein (pstA)	34.1%	AF1352	ISA1083-2, putative transposase	31.5%
TRANSPORT AND BINDING PROTEINS			AF1357	phosphate ABC transporter, permease protein (pstC)	33.7%	AF2140	ISA1083-3, ISORF2	30.8%
<i>General</i>			AF1360	phosphate ABC transporter, regulatory protein (phoU)	26.9%	AF2139	ISA1083-3, putative transposase	31.5%
AF0393	ABC transporter, ATP-binding protein	34.5%	AF0791	phosphate permease, putative	31.1%	AF0278	ISA1214-1, ISORF2	27.7%
AF0394	ABC transporter, ATP-binding protein	35.2%	AF1798	phosphate permease, putative	52.9%	AF0279	ISA1214-1, putative transposase	33.3%
AF1006	ABC transporter, ATP-binding protein	35.1%	AF0092	sulfate ABC transporter, ATP-binding protein (cysA)	54.2%	AF0305	ISA1214-2, ISORF2	27.7%
AF1018	ABC transporter, ATP-binding protein	57.7%	AF0093	sulfate ABC transporter, permease protein (cysT)	44.1%	AF0306	ISA1214-2, putative transposase	33.3%
AF1021	ABC transporter, ATP-binding protein	37.8%	<i>Carbohydrates, organic alcohols, and acids</i>			AF0642	ISA1214-3, ISORF2	26.5%
AF1136	ABC transporter, ATP-binding protein	39.3%	AF0347	C4-dicarboxylate transporter (mae1)	24.5%	AF0857	ISA1214-4, ISORF2	27.7%
AF1139	ABC transporter, ATP-binding protein	38.2%	AF1426	glycerol uptake facilitator, MIP channel (glpF)	36.2%	AF0858	ISA1214-4, putative transposase	33.3%
AF1300	ABC transporter, ATP-binding protein	34.1%	AF0013	hexuronate transporter (exuT)	25.1%	AF2091	ISA1214-5, ISORF2	26.5%
AF1469	ABC transporter, ATP-binding protein	43.5%	AF0806	L-lactate permease (lcpP)	31.7%	AF2092	ISA1214-5, putative transposase	33.3%
AF1819	ABC transporter, ATP-binding protein	51.1%	AF0008	oxalate/formate antiporter (oxfT-1)	25.7%	AF2223	ISA1214-6, ISORF2	26.5%
AF1982	ABC transporter, ATP-binding protein	41.3%	AF0367	oxalate/formate antiporter (oxfT-2)	33.2%	AF2222	ISA1214-6, putative transposase	25.6%
AF2364	ABC transporter, ATP-binding protein	53.5%	AF1069	pantothenate permease (panF-1)	28.9%	AF0138	transposase IS240-A	43.3%
AF1005	ABC transporter, ATP-binding protein, putative	28.7%	AF1205	pantothenate permease (panF-2)	24.8%	AF0895	transposase IS240-A	46.2%
AF1064	ABC transporter, ATP-binding protein, putative	36.0%	AF0237	pantothenate permease (panF-3)	25.1%	AF2390	transposase, authentic frameshift	24.0%
AF1983	ABC transporter, periplasmic binding protein	25.4%	AF0041	polysaccharide ABC transporter, ATP-binding protein (rtbB-1)	42.5%	AF0137	transposase, putative	23.6%
AF1981	ABC transporter, permease protein	29.9%	AF0290	polysaccharide ABC transporter, ATP-binding protein (rtbB-2)	43.9%	UNKNOWN		
AF1995	sodium- and chloride-dependent transporter	52.5%	AF0042	polysaccharide ABC transporter, permease protein (rtbA-1)	27.5%	AF0477	AAA superfamily ATPase	35.0%
<i>Amino acids, peptides and amines</i>			AF0289	polysaccharide ABC transporter, permease protein (rtbA-2)	28.5%	AF0513	allene oxide synthase, putative	39.5%
AF1766	amino acid ABC transporter, periplasmic binding protein/protein kinase	27.4%	AF0887	ribose ABC transporter, ATP-binding protein (rbsA-1)	33.3%	AF0478	ATP-binding protein PhnP (phnP)	30.9%
AF0222	branched-chain amino acid ABC transporter, ATP-binding protein (braF-1)	42.7%	AF1170	ribose ABC transporter, ATP-binding protein (rbsA-2)	27.9%	AF1775	atrazine chlorohydrolase, putative	34.4%
AF0822	branched-chain amino acid ABC transporter, ATP-binding protein (braF-2)	44.7%	AF0888	ribose ABC transporter, permease protein (rbsC-1)	24.1%	AF0973	bile acid-inducible operon protein F (baIF-1)	30.8%
AF0959	branched-chain amino acid ABC transporter, ATP-binding protein (braF-3)	37.6%	AF0889	ribose ABC transporter, permease protein (rbsC-2)	31.2%	AF0974	bile acid-inducible operon protein F (baIF-2)	29.9%
AF1390	branched-chain amino acid ABC transporter, ATP-binding protein (braF-4)	58.7%	AF2014	sugar transporter, putative	26.0%	AF1315	bile acid-inducible operon protein F (baIF-3)	31.3%
AF0221	branched-chain amino acid ABC transporter, ATP-binding protein (braG-1)	48.2%	<i>Cations</i>			AF2063	c-myc binding protein, putative	21.7%
AF0823	branched-chain amino acid ABC transporter, ATP-binding protein (braG-2)	42.9%	AF3977	ammonium transporter (amt-1)	44.3%	AF1992	calcium-binding protein, putative	31.2%
AF0958	branched-chain amino acid ABC transporter, ATP-binding protein (braG-3)	34.1%	AF1746	ammonium transporter (amt-2)	40.0%	AF2287	carotenoid biosynthetic gene ERWORTS, putative	49.4%
AF1389	branched-chain amino acid ABC transporter, ATP-binding protein (braG-4)	64.6%	AF0473	cation-transporting ATPase, P-type (pacS)	44.0%	AF0512	chloroplast inner envelope membrane protein	42.5%
AF0223	branched-chain amino acid ABC transporter, periplasmic binding protein (braC-1)	34.3%	AF0152	copper-transporting ATPase, P-type (copB)	44.5%	AF2251	competence-damage protein, putative	28.0%
AF0827	branched-chain amino acid ABC transporter, periplasmic binding protein (braC-2)	26.8%	AF2394	iron (II) transporter (feoB-1)	33.3%	AF0090	dehydrogenase	34.1%
AF0962	branched-chain amino acid ABC transporter, periplasmic binding protein (braC-3)	25.6%	AF0561	iron (II) transporter (feoB-2)	48.0%	AF1498	dehydratase, putative	29.4%
AF1391	branched-chain amino acid ABC transporter, periplasmic binding protein (braC-4)	50.1%	AF0430	iron (III) ABC transporter, ATP-binding protein (hemV-1)	50.4%	AF1518	DNA/pantothenate metabolism flavoprotein, putative	51.4%
AF0224	branched-chain amino acid ABC transporter, permease protein (braD-1)	25.4%	AF1401	iron (III) ABC transporter, ATP-binding protein (hemV-2)	58.7%	AF0039	dolichol-P-glucose synthetase, putative	33.7%
AF0825	branched-chain amino acid ABC transporter, permease protein (braD-2)	30.8%	AF1397	iron (III) ABC transporter, ATP-binding protein (hemV-3)	35.2%	AF0328	dolichol-P-glucose synthetase, putative	39.0%
AF0961	branched-chain amino acid ABC transporter, permease protein (braD-3)	23.9%	AF0431	iron (III) ABC transporter, permease protein (hemU-1)	28.2%	AF0581	dolichol-P-glucose synthetase, putative	27.5%
AF1392	branched-chain amino acid ABC transporter, permease protein (braD-4)	65.4%	AF1402	iron (III) ABC transporter, permease protein (hemU-2)	36.2%	AF0569	DR-beta chain MHC class II	37.7%
AF0225	branched-chain amino acid ABC transporter, permease protein (braE-1)	28.7%	AF0786	magnesium and cobalt transporter (corA)	40.1%	AF0383	endonuclease III, putative	47.1%
AF0824	branched-chain amino acid ABC transporter, permease protein (braE-2)	31.3%	AF0217	Na ⁺ /H ⁺ antiporter (napA-1)	28.2%	AF1150	erpK protein, putative	54.9%
AF0960	branched-chain amino acid ABC transporter, permease protein (braE-3)	30.1%	AF1245	Na ⁺ /H ⁺ antiporter (napA-2)	28.4%	AF2372	extragenic suppressor (subB)	37.0%
AF1393	branched-chain amino acid ABC transporter, permease protein (braE-4)	60.5%	AF0846	Na ⁺ /H ⁺ antiporter (rhe2)	33.1%	AF1418	glycerol-3-phosphate cytidyltransferase (taqD)	56.6%
AF1612	cationic amino acid transporter (cat-1)	29.5%	AF0715	potassium channel, putative	39.5%	AF0744	GTP-binding protein	33.4%
AF1774	cationic amino acid transporter (cat-2)	38.0%	AF1673	potassium channel, putative	36.3%	AF1181	GTP-binding protein	36.3%
AF1770	dipeptide ABC transporter, ATP-binding protein (dppD)	47.8%	AF2197	potassium uptake system protein (trkA-1)	24.6%	AF1364	GTP-binding protein	57.5%
AF1771	dipeptide ABC transporter, ATP-binding protein (dppF)	43.1%	AF0218	TRK potassium uptake system protein (trkA-2)	30.2%	AF2146	GTP-binding protein	65.9%
AF1767	dipeptide ABC transporter, dipeptide-binding		AF0838	TRK potassium uptake system protein (trkH)	39.8%	AF0428	GTP-binding protein, GTP1/OBG-family	43.9%
			<i>Other</i>			AF2237	HAM1 protein	31.4%
			AF0834	ferritin, putative	39.8%	AF2211	HIT family protein (hit)	29.6%
			AF1980	heme exporter protein C (helC)	29.0%	AF0216	L-isospartyl protein carboxyl methyltransferase	
			AF1144	multidrug resistance protein	29.2%		PimT, putative	35.5%
			AF1325	multidrug resistance protein	29.9%	AF2313	maoC protein (maoC)	43.0%
						AF0429	methyltransferase	43.8%
						AF0186	nifS protein, class-V aminotransferase (nifS-1)	46.1%
						AF0564	nifS protein, class-V aminotransferase (nifS-2)	46.1%
						AF0185	nifU protein (nifU-1)	55.6%
						AF0565	nifU protein (nifU-2)	55.6%
						AF0632	nifU protein (nifU-3)	47.4%
						AF1781	nodulation protein NteD (nfeD)	33.4%
						AF2269	nucleotide-binding protein	48.7%
						AF2382	nucleotide-binding protein	49.1%
						AF0374	p-nitrophenyl phosphatase (pho2)	31.7%
						AF1978	periplasmic divalent cation tolerance protein (cutA)	31.3%
						AF1652	prepro-subtilisin sendai, putative	31.6%
						AF2021	rod shape-determining protein (mreB)	26.6%
						AF1778	stage V sporulation protein (spoVG)	43.9%
						AF1970	TPR domain-containing protein	29.0%
						AF2202	tryptophan-specific permease, putative	25.2%
						AF0816	vfp-therm, putative	42.1%
						AF1679	vfp-therm, putative	45.1%