sample, should fit the sample better than the transition probability model.

Whether the G_1 rate model provides a better description of cell-cycle kinetics than the transition probability model is partly a matter of personal preference. Certainly, it can match the experimental data. However, the G_1 rate model requires six arbitrary parameters, the values of which are selected by trial and error using a computer. With this degree of flexibility, it would be surprising if the model could not be made to fit. In contrast, the two-transition modification of the transition probability model² describes observed distributions of cycle times using only three parameters, the values of which are not flexible but derived uniquely from measured cell-cycle statistics. It also explains the observed correlation between sibling generation times, the lag which follows mitogenic stimulation, and the similarity of this lag to the minimum generation time of rapidly proliferating cells—unlike any other model we know of.

We also find the G_1 rate model to be biologically implausible. Because the overall G_1 rate for a cell is not simply the average of the rates at different 'points' in G_1 , an inverse normal distribution of G_1 times implies one of two things: either that each cell must maintain an invariant rate of progress through G_1 , the individual rates being normally distributed, or, if G_1 is considered to consist of several 'steps', that the rates of each must be perfectly correlated with the rate of the initial step, this being normally distributed in the population (that is, non-independent variance between steps). In either case, the G_1 rate must effectively be predetermined at birth. Furthermore, for those cases where there is no mother/daughter correlation, the G_1 rate must be reset at random for each generation. The problem then is to explain how the G_1 rate can vary between cells while at the same time being predetermined at the start of G_1 . We cannot think of any mechanism that could achieve this. In particular, we cannot see how such things as RNA content or protein synthetic capacity could be responsible (as suggested by Castor¹) as they do not remain constant during G_1 .

Regardless of mechanism, the basic contention of the G_1 rate model is that variation in cycle times arises from an underlying Gaussian distribution, that is, the summation of many random variables. The 'attraction' of this is that it allows cycle variability to be ignored, or at least relegated as being unimportant. In contrast, the modified transition probability model asserts that variation is inherent to the cell cycle because two critical events occur with low probability². Whether the latter viewpoint is eventually vindicated depends on whether the hypothetical transitions can be ascribed to real biological events, and this will not be possible by kinetics alone. However, the

parallels between the two-transition model and the biogenesis of mitotic centres may prove helpful².

R. F. BROOKS
J. A. SMITH
D. C. BENNETT
K. M. V. RICHMOND

Cell Proliferation Laboratory,
Imperial Cancer Research Fund,
PO Box 123,
Lincoln's Inn Fields,
London WC2A 3PX, UK

1. Castor, L. N. *Nature* **287**, 857–859 (1980).
2. Brooks, R. F., Bennett, D. C. & Smith, J. A. *Cell* **19**, 493–504 (1980).
3. Shields, R. in *Control Mechanisms in Animal Cells* (eds Jimenez de Asua, L., Levi-Montalcini, R., Shields, R. & Iacobelli, S.) 157–164 (Raven, New York, 1980).
4. Miyamoto, H., Zeuthen, E. & Rasmussen, L. J. *Cell Sci.* **13**, 879–888 (1973).
5. Shields, R. & Smith, J. A. *J. cell. Physiol.* **91**, 345–356 (1977).
6. Dawson, K. B., Madoc-Jones, H. & Field, E. O. *Expl Cell Res.* **38**, 75–84 (1965).

CASTOR REPLIES—In considering a mechanism for the G_1 rate model¹, we can speculate that a cell requires a fixed quantity of a set of proteins before becoming committed to DNA synthesis, and that the rate of synthesis of these proteins is normally distributed. The random resetting in each generation of some proportion of the synthetic capacity may result from the disaggregation and reaggregation of polyribosomes at mitosis, or from different activities of nucleolar organizing regions in successive generations^{2,3}.

Mother/daughter correlation coefficients found in several cell populations^{4,5} are too high to be explained by experimental artefacts such as clonal heterogeneity or local inhomogeneities in the culture environment. A more likely explanation is that protein synthetic capacity is inherited to some extent from mother to daughter. Positive mother/daughter correlation can be obtained in computer simulations by combining cycle time data having different means. This procedure does not give any increase in β curvature. On the contrary, there is a slight decrease when data generated by the G_1 rate model are combined. With regard to the 1-h frame interval of the data of Miyamoto *et al.*⁴, my simulations show that it yields the same β curvature as a frame interval of 4 min, provided the time of the former is taken as the middle of the interval. This was done for both the experimental data and the simulation of curve *b* in ref. 1.

The method used to determine the statistical significance of the curvatures gives a quantitative measure of what can otherwise be appreciated only from subjective comparisons of many successive simulations: in relatively few cases (200 of 10,000 for data set *a* and 600 of 10,000 for set *b*) does the transition probability model produce β curves that deviate as far from linearity as do some experimental data. Changing the midpoint of the calculation would affect these numbers

slightly, but it would not invalidate the conclusion. In contrast to the linearity of β curves generated by the transition probability model, the G_1 rate model is consistent with both the slight downward curvature of sets *a* and *b* and the linearity (within sampling error) of sets *c* and *d*.

The 'flexibility' of the G_1 rate model arises because the means, variances and measures of sibling correlation in the G_1 phase and in the remainder of the cycle are potentially independent entities. Fewer parameters would be a convenience for computation, but the number of parameters does not determine whether a model realistically describes the mechanism of cycle control. A generalization such as transition probability might eventually simplify the problem, but all reasonable alternatives must be excluded before it can be accepted. It is more likely that future models will be even more complex.

The derivation of the parameters of the transition probability model from the means, variances and correlation coefficients of the cell-cycle distributions, even using several ostensibly independent calculations^{6,7}, does not validate the model. To argue that it does is to imply that a given distribution could have been derived from only one set of conditions. In formal logic this is the fallacy of affirming the consequent⁸. The G_1 rate model generates differences in the cycle times of siblings that conform to an exponential distribution as accurately as do populations of cells. It leads to the same means, variances and correlation coefficients as those measured for the cells, and therefore to the same mathematical relationships, without any random transition.

These considerations do not rule out a random transition as a possible explanation for cell-cycle control. The inability of the transition probability model to produce β curvature could be remedied by increasing its complexity by adding one or more parameters. Both models would then probably match all the kinetic data. As both models cannot possibly provide the correct explanation for cycle control, it is essential that further efforts be made to exclude one or both of them.

LARON N. CASTOR

Institute for Cancer Research,
Fox Chase Cancer Center,
7701 Burholme Avenue,
Philadelphia, Pennsylvania 19111, USA

1. Castor, L. N. *Nature* **287**, 857–859 (1980).
2. Mazia, D. in *The Cell* Vol. 3 (eds Brachet, J. & Mirsky, A. E.) 308–309, 372–374 (Academic, New York, 1961).
3. Goodpasture, C. & Bloom, S. E. *Chromosoma* **53**, 37–50 (1975).
4. Miyamoto, H., Zeuthen, E. & Rasmussen, L. J. *Cell Sci.* **13**, 879–888 (1973).
5. Collyn-d'hooghe, M., Valleron, A.-j. & Malaise, E. P. *Expl Cell Res.* **106**, 405–407 (1977).
6. Brooks, R. F., Bennett, D. C. & Smith, J. A. *Cell* **19**, 493–504 (1980).
7. Shields, R. in *Control Mechanisms in Animal Cells* (eds Jimenez de Asua, L., Levi-Montalcini, R., Shields, R. & Iacobelli, S.) 157–164 (Raven, New York, 1980).
8. Salmon, W. C. *Logic*, 27 (Prentice-Hall, Englewood Cliffs, 1963).