

In our article¹, we did not intend to misinterpret Nicholson and Flohn⁹ when citing their hypothesis of distinct latitudinal shifts of the subtropical climatic belt from interglacial to glacial times. Accidental over-simplification of the alternatives may be one of the reasons for the apparent wrong citation. On the other hand, we feel rather corroborated in our statement by the comments of Nicholson discussed above.

G. TETZLAFF
K. WOLTER

*Institut für Meteorologie
und Klimatologie,
Universität D-3000
Hannover 21, FRG*

M. SARNTHEIN

*Geologisch-Paläontologisches Institut,
Universität D-2300 Kiel 1, FRG*

1. Sarnthein, M., Tetzlaff, G., Koopmann, B., Wolter, K. & Pfäumann, U. *Nature* **293**, 193–196 (1981).
2. Street-Perrott, F. A. & Roberts, N. in *Variations in the Global Water Budget* (eds Street-Perrott, F. A., Beran, M. & Ratcliffe, R. A. S.) (Reidel, Dordrecht, in the press).
3. Sarnthein, M. et al. in *The Geology of the Northwest African Continental Margin* (eds von Rad et al.) 544–604 (Springer, Heidelberg, 1982).
4. Dubief, J. *Le Climat du Sahara* Vols 1, 2 (1963).
5. Wiin-Nielsen, A. & Krishnamurti, T. N. (eds) *Compendium of Meteorology* Vol. 2, Pt 4 (WMO no. 364, 1979).
6. Jaenicke, R. & Schuetz, L. *J. geophys. Res.* **83**, 3585–3599 (1978).
7. Pedgley, D. & Krishnamurti, T. N. *Mon. Weath. Rev.* **104**, 149–167 (1976).
8. Houghton, D. in *Rep. U.S. GATE Central Program Workshop*, Boulder (1977).
9. Nicholson, S. E. & Flohn, H. *Climat. Change* **2**, 313–348 (1980).

Western diets and faecal nitrosamines

THE paper by Suzuki and Mitsuoka¹, while apparently demonstrating clear differences in nitrosamine production in individuals on different diets, makes unwarranted claims for the significance of these findings.

Japan has a low incidence rate of large bowel cancer compared with the United States² or Australia³. International correlation studies^{4,5}, studies of subcultures within societies⁶, and case-control and cohort studies of individuals^{7–10} have implicated dietary factors in colon carcinogenesis. Current hypotheses involve fat^{4,9,11}, fibre^{9,11}, alcohol¹¹, female sex hormones¹², vitamin A¹³, and cruciferous vegetables⁸ either as promoting colon cancer, via endogenous and bacterial bile acid metabolism, or as protective agents.

To test the hypothesis that a balanced Western diet constitutes a greater risk of exposure to nitrosamines (which act as initiating agents in some cancers) it is necessary to design an experiment which actually uses a balanced Western diet. The

diet used by Suzuki and Mitsuoka bears no resemblance to a normal Western diet. Specifically, the energy content of a typical Western diet^{14,15} is 9.7–10.5 MJ per day, compared with 11.8 MJ per day in the study of Suzuki and Mitsuoka. As a percentage of energy, protein intake is normally 12–15% (as opposed to 25%), fat ~40% (compared with 55%), carbohydrate ~45% (as opposed to 20%). In some European countries, certain vegetables may be consumed for breakfast but nowhere are salad vegetables eaten for breakfast. Indeed these vegetables were deliberately added to the 'typical' Western breakfast to increase nitrate intake. This ensures a good test of an hypothesis but not the one we are led to believe is being tested by the authors.

Although Suzuki and Mitsuoka have established that a very high protein and fat diet with a large intake of nitrate increases the faecal output of nitrosamines, they have not demonstrated that this has anything to do with differences existing in the real world.

J. D. POTTER

*Division of Human Nutrition,
CSIRO, Kintore Avenue,
Adelaide, South Australia 5000*

1. Suzuki, K. & Mitsuoka, T. *Nature* **294**, 453–456 (1981).
2. Waterhouse, J., Muir, C., Correa, P. & Powell, J. *Cancer Incidence in Five Continents* Vol. 3 (IARC Scient. Publ. no. 15, Lyon, 1976).
3. Bonett, A. *Cancer in South Australia* (South Australian Government, 1980).
4. Armstrong, B. & Doll, R. *Int. J. Cancer* **15**, 617–631 (1975).
5. Drasar, B. S. & Irving, D. *Br. J. Cancer* **27**, 167–172 (1973).
6. Phillips, R. *Cancer Res.* **35**, 3513–3522 (1975).
7. Hirayama, T. *Nutr. Cancer* **1**, 67–81 (1979).
8. Graham, S., Dayal, H., Swanson, M., Wittelman, A. & Wilkinson, G. *J. natn. Cancer Inst.* **61**, 709–714 (1978).
9. Dales, L. G., Friedman, G. D., Ury, H. K., Grossman, S. & Williams, S. R. *Am. J. Epidemiol.* **109**, 132–144 (1979).
10. Jain, M. et al. *Int. J. Cancer* **26**, 757–768 (1980).
11. McMichael, A. J., Potter, J. D. & Hetzel, B. S. *Int. J. Epidemiol.* **8**, 295–303 (1979).
12. McMichael, A. J. & Potter, J. D. *J. natn. Cancer Inst.* **65**, 1201–1207 (1980).
13. Peto, R., Doll, R., Buckley, J. D. & Sporn, M. B. *Nature* **299**, 201–208 (1981).
14. Baghurst, K. I. & Baghurst, P. A. in *The Assessment of Nutritional Status* (eds Hetzel, B. S. & Baghurst, K. I.) 139–160 (Menzies Foundation, Melbourne, 1981).
15. U.S. Senate Select Committee of Nutrition and Human Needs, *Dietary Goals for America* (US Government Printing Office, Washington D.C., 1977).

SUZUKI AND MITSUOKA REPLY—We agree with Potter that the diet used in our study does not exactly resemble a typical Western diet. To determine that nitrosamine formation in the intestine depends on diet, we used in our experiments a diet which contained a large

amount of nitrosamine precursors. Our study demonstrated that the faecal output of nitrosamines was increased by this kind of Western-style diet. However, we cannot implicate these diets directly in the aetiology of colon cancer. Further investigations of nitrosamine formation in the intestine in populations of high and low risk for colon cancer are necessary.

With respect to the p.p.b. analysis of nitrosamines in biological specimens, there are several problems associated with artefacts and contamination during the analysis. Nitrosamines are generally extracted from the specimens in alkaline conditions, because this inhibits nitrosation. We have noticed recently, however, that artificial formation of nitrosamines occurs in these conditions. This reaction seems to resemble the 'alkali effect' reported by Tozawa¹. We have used 1 mM of morpholine to monitor artificial nitrosamine formation during the analysis, as morpholine is nitrosated 500 times more readily than piperidine². However, we have recently found that morpholine is more difficult to nitrosate than piperidine in alkaline conditions, which renders morpholine unsuitable for use as a monitor of possible artificial formation of nitrosamines. We are re-investigating the faecal output of nitrosamines, thus our data³ may require correction.

KUNIO SUZUKI
TOMOTARI MITSUOKA

*Animal Physiology Laboratory,
The Institute of Physical
and Chemical Research,
Wako-shi, Saitama 351, Japan*

1. Tozawa, H. *Bull. Japan Soc. scient. Fish.* **44**, 797–802 (1978); **45**, 1585 (1979); in *N-nitroso Compounds: Occurrence and Biological Effect* (eds Bartsch, H., O'Neill, I. K., Castegnaro, M. & Okada, M.) (IARC Publ. no. 41, Lyon, in the press).
2. Mirvish, S. S. *Tox. appl. Pharma.* **31**, 325–351 (1975).
3. Suzuki, K. & Mitsuoka, T. *Nature* **294**, 453–456 (1981).

Matters Arising

Matters Arising is meant as a vehicle for comment and discussion about papers that appear in *Nature*. The originator of a Matters Arising contribution should initially send his manuscript to the author of the original paper and both parties should, wherever possible, agree on what is to be submitted. Neither contribution nor reply (if one is necessary) should be longer than 500 words and the briefest of replies, to the effect that a point is taken, should be considered.