Plant biochemistry

Mobile genes of chloroplasts and the promiscuity of DNA

from John Ellis

A KEY problem in evolutionary biology is to account for the origins and properties of the genetic systems that occur outside the nucleus in all eukaryotic cells. It seems a high cost to the plant cell to specify at least 200 extra polypeptides involved in replication, transcription and translation by mitochondria and chloroplasts in order to make a rather small number of polypeptides inside these organelles instead of outside them. In the case of chloroplasts, the explanation may lie in their proposed origin in free-living photosynthetic cyanobacteria through endosymbiosis^{1,2}.

Extant cyanobacteria which carry out oxygenic photosynthesis contain the CO₂-fixing enzyme ribulose biphosphate carboxylase/oxygenase, which consists of catalytic large subunits and regulatory small subunits. In these cyanobacteria, both types of subunit are encoded in the same genome along with all the other proteins necessary for a phototrophic prokaryote.

A similar enzyme is found inside the chloroplasts of photosynthetic eukaryotes, but in these organisms the two types of subunit are encoded and synthesized within different genetic compartments³. The large subunit is a product of the chloroplast genome, while the small subunit is synthesized as a higher-molecular-weight precursor by cytoplasmic ribosomes translating a nuclear-encoded mRNA. Post-translational transport of this precursor into the chloroplast is accompanied by cleavage to the mature size by a soluble chloroplast protease which removes the amino-terminal extension⁴. The small subunit is representative of the majority of chloroplast polypeptides, since these are also encoded in nuclear genes and synthesized as precursors by cytoplasmic ribosomes⁵.

Thus if the endosymbiont hypothesis is correct, a massive transfer of genes from symbiont to nucleus must have occurred during evolution; and one might expect that intermediate stages may have persisted. In this issue of *Nature*⁶, Heinhorst and Shively report that in the organism *Cyanophora paradoxa*, the carboxylase

small subunit is encoded not in the nuclear genome but in the genome of the photosynthetic organelle.

The protist *C. paradoxa* contains two to four intracellular structures termed cyanelles, each surrounded by a vacuolar membrane; the cyanelles contain unstacked thylakoids and photosynthetic pigments typical of free-living cyanobacteria, and are invested by a thin peptidoglycan cell wall⁷. Cyanelles were first regarded as ingested cyanobacteria, but their genome resembles that of chloroplasts both in structure (a circular molecule containing an inverted repeat encoding rRNA) and in size (molecular weight about 10⁸). This genome is too small to encode enough proteins for a free-living organism⁸. Should cyanelles then be regarded as chloroplasts? But in this event, it would be expected that the carboxylase small subunit present in cyanelles would be encoded in the nuclear genome of the protist, and Heinhorst and Shively⁶ have shown that this is not the case. Use of a cloned cDNA hybridization probe prepared from the small-subunit mRNA of *Pisum sativum* demonstrates that the gene for the small subunit in *Cyanophora* resides near that for the large subunit in the cyanelle genome. (The hybridization was performed at low stringency to allow for the variability of the small subunit and the evolutionary distance between *Cyanophora* and *Pisum*.) Similar results were reported for *Cyanophora* by H.J. Bohnert, E.J. Crouse, W. Löffelhardt and H. Mücke at a recent meeting in Tübingen*.

These observations support the view that cyanelles 'represent a bridge between cyanobacteria and chloroplasts'⁹, and suggest that the location of the small-subunit gene should be determined in a wider range of eukaryotes, especially among the more primitive algae where transfer of the gene to the nucleus may not have occurred. Evidence was presented at the Tübingen

colloquium by K. Zetsche, K. Steinmüller and M. Kaling that in some red algae, the carboxylase small-subunit mRNA is present in the same poly(A)⁻ RNA fraction that contains the large-subunit mRNA; in other eukaryotes, such as *Pisum*, the small-subunit mRNA contains poly(A) while the large-subunit mRNA does not. This suggests that in these red algae the small subunit is synthesized inside the chloroplast, although direct evidence for this conclusion was not presented. Of special interest is the finding that the small subunit synthesized by incubating red algal poly(A)⁻ RNA with a reticulocyte lysate is made not as a precursor but at its mature size. This observation raises the question as to the evolutionary origin of the amino-terminal extension characteristic of cytoplasmically synthesized chloroplast proteins.

It is striking that in the four species so far examined (soybean, pea, wheat and *Chlamydomonas*), the amino terminus of the mature small subunit is occupied by methionine, even though this is not the first amino acid to be polymerized¹⁰. The presence of methionine at this position may be a coincidence, but a more interesting possibility is that it reflects the evolutionary origin of the precursor. Fusion of a prokaryotic gene for mature-sized small subunit possessing methionine as its initiating amino acid with a DNA sequence specifying transport across the chloroplast envelope would produce the observed amino acid sequence. The origin of the transport DNA sequence is a mystery.

The unusual location of the carboxylase small-subunit gene in *Cyanophora* supports the inference of the endosymbiont hypothesis that DNA leaves the symbiont as it evolves into an organelle. Presumably duplicate copies of the exported sequences are retained within the organelle until the nuclear sequences become fully functional by addition of the information for the amino-terminal extension of the polypeptides. Is there evidence for shared nucleotide sequences between chloroplast and nucleus in extant species?

In an earlier article in *News and Views*, I proposed the term 'promiscuous DNA' to describe nucleotide sequences found in more than one of the three membrane-bound organellar genetic compartments of eukaryotes¹¹. Examples of promiscuous DNA are mitochondrial sequences present in the nucleus of yeast¹² and chloroplast sequences present in the mitochondria of maize¹³. A further example is reported by Timmis and Scott¹⁴ who find that the nuclear DNA of spinach contains in-

*The Second International Colloquium on Endocytobiology was held in Tübingen on 11-15 April 1983.

ERRATA

In the *News and Views* article 'New light on climate from old isotope ratios' by Pieter M. Grootes (*Nature* 303, 753-754; 1983), there was a mistake in the third paragraph. The first sentence of that paragraph should read:

"The d term, introduced by Dansgaard², is the constant in a simple empirical relationship between the hydrogen and oxygen isotopic composition of worldwide freshwater sources ($d = \delta D - 8\delta^{18}O$ where δ is the relative difference in

isotopic composition between the sample and a standard, expressed in units per mill)."

In the article by David Blow in last week's *News and Views* on 'Computer cues to combat hypertension' (*Nature* 304, 213-214; 1983), the name of the third author of the paper to which the *News and Views* article referred and which appeared in the same issue (p.273-275) was erroneously missed out. Thus Laurence Pearl should have been acknowledged in the first paragraph and in reference 1.