Fig. 4 Changes in the *HpaII*/*Bam*HI digestion pattern of the δ -crystallin genes in different tissues and in lens ectoderm during development. **A**, *HpaII*/*Bam*HI digestion of 13-day headless embryo (Emb.) and lens DHA (Lens) with the enzymes indicated at the bottom of the figure. **B**, *HpaII*/*Bam*HI digests of lens ectoderm DNA from 40–144-h embryos staged as described for Fig. 3. **C**, *HpaII*/*Bam*HI digests of DNA from 40-h body ectoderm and 65-h headless embryos. The procedures in these experiments were identical to those described for Fig. 2, except that simian virus 40 DNA was used as an internal standard to monitor complete cleavage during restriction enzyme digestions.

While the appearance of the 5-kb fragment may be related to gene activation, we cannot be certain that this event precedes transcription because of limitations in staging embryos and because there may be low levels of transcription before this stage. Our assay would not have detected a possible earlier event in the hypomethylation process, the formation of hemimethylated DNA at this site, since *HpaII* does not cleave hemimethylated DNA¹⁰. This could be a functionally important step affecting transcription. As *HpaII* analysis only monitors a small fraction of the possible methylated sites (and in this study only those *HpaII* sites in the 3' region of the genes), there could also be other methylation changes which occur even earlier than that observed at 52 h.

Several studies argue that hypomethylation does not occur when transcription begins^{11,12}, or that it is not sufficient for transcription¹³; however, our results indicate how important it may be to assay many methylation changes during gene activation to observe a few sites which are tightly coupled to the onset of transcription. Only by a developmental profile can one identify transient products, such as the 5-kb fragment we observed. Unfortunately, this is not a simple procedure in most embryonic systems. The only other study of this kind¹⁴ showed that in the red blood cell lineage there is no hypomethylation of the globin genes 12–15 h before globin is synthesized. Here, however, because it was impossible to isolate a pure population of red blood cell precursors during the 12 h before globin gene activation, the exact timing of hypomethylation could not be determined.

As most of the methylation changes we see follow the onset of transcription, they cannot have a causal role in gene activation; it is possible that they affect transcriptional efficiency or stabilize the transcriptionally active state. If these sites were hemimethylated earlier, they might have a role in gene activation; however, this seems unlikely as the 48-h gap between first δ -crystallin synthesis and the appearance of these hypomethylated sites far exceeds the one cell cycle (9 h¹⁵) expected to yield hypomethylated DNA from hemimethylated DNA.

Several of the methylation changes which follow δ -crystallin gene activation (Fig. 4B) are also seen in non-lens tissues from older embryos (Fig. 4A; ref. 16), further suggesting that they are not involved in transcriptional regulation. Although the δ -crystallin genes are transcribed at extremely low levels in some non-lens tissues¹⁶, other control mechanisms, possibly including other methylation changes, must also be important in modulating the level of δ -crystallin transcripts in these tissues.

We thank Dr Joram Piatigorsky for the cDNA clone used in this study. This work was supported by NIH grant HD-07098 to the Marine Biological Laboratory in support of training in Embryology and by a NIH Research Career Development Award (HD-00289) and grants from the American Cancer Society (VC-306), NSF (PCM-82-08266) and the Thomas F. and Kate Miller Jeffress Memorial Trust (J-24) to R.M.G.

Received 10 June; accepted 18 August 1983.

1. Razin, A. & Riggs, A. D. *Science* **210**, 604–610 (1980).
2. Doerfler, W. *J. gen. Virol.* **57**, 1–20 (1981).
3. Taylor, S. M. & Jones, P. A. *Cell* **17**, 771–779 (1979).
4. Waalwijk, C. & Flavell, R. A. *Nucleic Acids Res.* **5**, 4631–4641 (1978).
5. Jones, R. E., DeFeo, D. & Piatigorsky, J. *J. biol. Chem.* **256**, 8172–8176 (1981).
6. Zwaan, J. & Ikeda, A. *Expl Eye Res.* **7**, 301–311 (1968).
7. Shinohara, T. & Piatigorsky, J. *Proc. natn. Acad. Sci. U.S.A.* **73**, 2808–2812 (1976).
8. Karkinen-Jääskeläinen, M. *J. Embryol. exp. Morph.* **44**, 167–179 (1978).
9. Zwaan, J. *Wilhelm Roux's Arch. dev. Biol.* **175**, 13–25 (1974).
10. Bird, A. P. *J. molec. Biol.* **118**, 49–60 (1978).
11. McKeon, C., Ohkubo, H., Pastan, I. & de Crombrughe, B. *Cell* **29**, 203–210 (1982).
12. Wilks, A. F., Cozens, P. J., Mattaj, I. W. & Jost, J.-P. *Proc. natn. Acad. Sci. U.S.A.* **79**, 4252–4255 (1982).
13. Ott, M.-O. *et al. Cell* **30**, 825–833 (1982).
14. Groudine, M. & Weintraub, H. *Cell* **24**, 393–401 (1981).
15. Zwaan, J. & Pearce, T. L. *Dev. Biol.* **25**, 96–118 (1971).
16. Bower, D. J., Errington, L. H., Cooper, D. N., Morris, S. & Clayton, R. M. *Nucleic Acids Res.* **11**, 2513–2527 (1983).
17. Southern, E. M. *J. molec. Biol.* **98**, 503–517 (1975).
18. Maniatis, T. *et al. Cell* **15**, 687–701 (1978).
19. Maniatis, T., Fritsch, E. F. & Sambrook, J. *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory, New York, 1982).
20. O'Rahilly, R. & Meyer, D. B. *Acta anat.* **36**, 20–58 (1959).
21. Hamburger, V. & Hamilton, H. L. *J. Morph.* **88**, 49–92 (1951).
22. Weintraub, H., Palter, K. & Van Lente, F. *Cell* **6**, 85–110 (1975).
23. Oakley, B. R., Kirsch, D. P. & Morris, N. R. *Analyt. Biochem.* **105**, 361–363 (1980).

Errata

IN the letter 'Viability of λ phages carrying a perfect palindrome in the absence of recombination nucleases' by D. R. Leach and F. W. Stahl, *Nature* **305**, 448–451 (1983), the two figures were transposed. Figure 2 should show the micrograph of Red⁺ DNA (B) and gels of restriction digests (A).

IN the letter 'Evidence for opiate receptors on pituitocytes' by S. L. Lightman *et al.*, *Nature* **305**, 235–237 (1983), the plates for Figs 1 and 2 on page 236 have been transposed.

IN the letter 'Discovery of a 6.1-ms binary pulsar PSR1953+29' by V. Boriakoff *et al.*, *Nature* **304**, 417–419 (1983), in Table 1 $\langle \eta_c \rangle$ should read $\langle n_c \rangle$ and the top line in the mass function in Table 1 should read $m_c^3 \sin^3(i)$.