



Altered contractile properties of the quadriceps muscle in people with spinal cord injury following functional electrical stimulated cycle training

HL Gerrits^{*1,2}, A de Haan^{1,3}, AJ Sargeant^{1,3}, A Dallmeijer¹ and MTE Hopman²

¹Institute for Fundamental and Clinical Human Movement Sciences, Vrije University Amsterdam, Amsterdam, The Netherlands; ²Institute for Fundamental and Clinical Human Movement Sciences, University of Nijmegen, Nijmegen, The Netherlands; ³Neuromuscular Biology Group, Manchester Metropolitan University, Alsager, UK

Study design: A longitudinal training study.

Objectives: To assess if contractile speed and fatigability of paralysed quadriceps muscles in individuals with spinal cord injury (SCI) can be altered by functional electrical stimulation leg cycle ergometry (FES-LCE) training.

Settings: The Sint Maartenskliniek rehabilitation centre and the University of Nijmegen, Nijmegen, the Netherlands.

Methods: Contractile properties of the quadriceps muscle were studied in seven people with motor-complete SCI who participated in a FES-LCE training program. Subjects trained for 30 min, three times per week for 6 weeks. Contractile speed and fatigue characteristics of electrically stimulated isometric contractions were compared before and after 6 weeks of FES-LCE.

Results: Fatigue resistance improved following FES-LCE training as indicated by the higher forces maintained in response to repetitive electrical stimulation. In contrast with an improved fatigue resistance, the maximal rate of force rise was unaffected, the speed of relaxation increased and the fusion of a 10 Hz force signal decreased. Furthermore, the force-frequency relationship shifted to the right at low stimulation frequencies, indicated by a decline in the ratio of 1 and 100 Hz force responses as well as the ratio of 10 and 100 Hz force responses.

Conclusion: FES-LCE training can change the physiological properties of the quadriceps muscle in people with SCI. Even after a short period of training, the stimulated muscles become more resistant to fatigue. Furthermore, the increased speed of relaxation and associated decreased fusion and altered force-frequency relationship following training may be related to adaptations in the calcium handling processes, which reflect an early response of long-term disused muscles.

Spinal Cord (2000) **38**, 214–223

Keywords: cycle ergometer; muscle characteristics; fatigue; paraplegia

Introduction

Spinal cord injury (SCI) results in loss of motor control, sensation and autonomic function below the level of the spinal lesion. This traumatic event often leads to significant central and peripheral physiological adaptations including impaired cardiovascular fitness,¹ muscle atrophy^{2,3} with concomitant greater prevalence of fast type II muscle fibres,^{4–6} osteoporosis^{7,8} and reduced blood circulation of the lower limbs.^{9–11}

For almost two decades now, much attention has been focused on improving health and fitness of individuals with SCI using a functional electrical stimulation leg cycle ergometer (FES-LCE). This ergometer provides

computerised electrical stimulation to hamstring, gluteal and quadriceps muscles in a co-ordinated sequence, which enables active leg cycling.¹² Several studies that assessed effects of FES-LCE training have demonstrated an improved cycling performance,^{3,13–17} and improved cardiovascular fitness.¹⁵ Musculoskeletal adaptations in the lower limbs following FES-LCE included reduced loss of bone mineral density,¹⁸ increased muscle mass^{3,14,19} and proportional increases in capillary number.¹⁹ Furthermore, it has recently been demonstrated that FES-LCE training can significantly alter the expression of myosin heavy chain in the quadriceps muscle of people with SCI. The cycle training induced a change from a predominance of muscle fibres containing IIB or a mixture of IIA/IIB isoforms to a predominance of fibres containing only the IIA isoform.^{3,20}

*Correspondence: HL Gerrits, Institute for Fundamental and Clinical Human Movement Sciences, Vrije University Amsterdam, v/d Boechorststraat 9, 1081 BT Amsterdam, The Netherlands

From animal studies, it is well known that chronically increased contractile activity can lead to adaptations in the muscle fibre composition and associated physiological properties of the muscle, such as an increased resistance to fatigue and altered contractile characteristics.^{21,22} In agreement with these findings, studies that applied different modes of electrical stimulation protocols on paralysed muscles of people with SCI, reported a decreased number of glycolytic type IIB and an increased number of oxidative type IIA^{2,23,24} or type I² fibres. Furthermore, in the tibialis anterior muscle, these structural, molecular and metabolic changes were accompanied by an increased fatigue resistance and slowing of contractions.²⁵

Recently, we have shown that the quadriceps muscles of individuals with SCI are characterised by faster contraction and relaxation rates and increased fatigability compared with muscles of able-bodied individuals.²⁶ Although these muscles have been studied extensively with respect to structural, morphological properties, to our knowledge no systemic data are available in the literature, which quantify alterations in the contractile properties of the stimulated quadriceps muscles after FES-LCE training. It may be expected, however, that these physiological properties undergo significant adaptations that are consistent with alterations in fibre type expression.^{3,20} Therefore, it is hypothesised that the characteristics of the quadriceps muscles in people with SCI shift towards that of more fatigue resistant slower muscles following FES-LCE training. To test this hypothesis we assessed the effect of a 6-week FES-LCE training period on the contractile properties of the paralysed quadriceps muscle of individuals with SCI.

Methods

Subjects

This study included seven male subjects with motor-complete spinal lesions. Table 1 represents the characteristics of the subjects with respect to the lesion and body composition. All subjects participated in a pre-training screening procedure. Screening

included a general physical examination, performed by the rehabilitation physician; evaluation of joint range of motion, spinal reflexes and spasticity performed by the local physiotherapist. Subjects who met any of the following exclusion criteria were excluded from participation: cardiac arrhythmia, high blood pressure (systolic blood pressure >180 mmHg, diastolic blood pressure >110 mmHg), presence of a pacemaker, pressure ulcers, metal implants in the area of stimulation, severely reduced mobility in hip or knee joints, absence of spinal reflexes, and previous bone fractures. All subjects were carefully informed about training and testing procedures and involved risks and gave their written informed consent. The Medical Ethical Committee of the University of Nijmegen approved this study.

FES-LCE training

The cycle training was performed using a computer controlled FES leg cycle ergometer (Ergys 2, Therapeutic Alliances Inc., Ohio, USA). This device provides electrical stimulation to 12 self-adhesive 50 × 89 mm surface electrodes (Bioflex, PE 3590, Danica, Leusden, the Netherlands), placed over hamstring, gluteal and quadriceps muscles (two electrodes for each muscle). Electrical stimulation (monophasic square wave pulses of 450 μs duration, 30 Hz, synchronised for each pair of electrodes) is applied in a co-ordinated sequence allowing cyclic patterns of muscle contractions resulting in leg cycling. The device is programmed to increase the stimulation current, with a maximum of 140 mA, in order to achieve a target-peddalling rate of 50 rpm. The external pedal resistance of the ergometer can be varied with 1/8 kp increments (corresponding with ~6.1 W at 50 rpm). When the pedalling rate drops below 45 rpm, the resistance is reduced and when the pedalling rate drops below 35 rpm, the stimulation is stopped.

Training protocol

All subjects trained for 6 weeks, three times a week. A training session consisted of 30 min FES-LCE exercise. Each exercise session started with 1 min of passive

Table 1 Characteristics of SCI subjects with respect to body composition and the level, classification and duration of the lesion

Subject	Age (years)	Height (m)	Weight (kg)	Lesion level	ASIA class*	Lesion duration (years)
1	44	1.91	95	C5/C6	B	14
2	36	1.72	62	C5	A	18
3	30	1.78	68	T4	A	2
4	61	1.80	82	T7/T8	A	2
5	40	1.78	78	T8	A	5
6	44	1.85	88	C5	B	27
7	28	1.89	72	T4	A	1

*ASIS (American Spinal Injury Association⁴⁹) score is used to classify the completeness of the lesion: A, sensory and motor complete; B, sensory incomplete but motor complete)

warming-up (without stimulation) as a research assistant turned the crankshaft manually. After the warming-up, stimulation of the leg muscles started, allowing active FES-LCE training until the subject could no longer maintain the pedalling rate above 35 rpm (represents one 'run'). When fatigue occurred before the end of the 30-min training session this run was followed by a 5-min rest period. This procedure was repeated with a maximum of five runs available to complete 30 min of active cycling. Each bout of exercise was followed by a 2-min passive cooling down, during which the pedals were turned by the assistant. When a subject was able to complete 30 min of FES-LCE exercise in one run with at least 90% of the programmed pedal resistance, the resistance was increased with 1/8 kp in the next training session up to a maximum of 7/8 kp.

Thigh girth

Thigh circumference was measured at 5 cm above the superior aspect of the patella and at mid-thigh (~20 cm above the patella) before and after the training period. In order to optimise the reproducibility of these measurements the average of three measurements was taken at each location.

Measurements of contractile properties

Isometric contractions of the knee extensors were obtained while subjects were seated upright with the pelvis and upper thigh securely fixed to the seat and the knee angle fixed at 100°. Electrically stimulated quadriceps contractions were generated using a high voltage constant current stimulator (model DS7A, Digitimer Limited, Hertfordshire, UK), delivering electrical pulses of 0.2 ms width to surface electrodes (10 × 13 cm, Electro-Medical Supplies, Greenham Limited, UK) placed over the anterior thigh. The muscles were activated with a stimulation intensity of 121 ± 30 mA. Although we were not able to determine exactly what proportion of the muscle was being stimulated, it was established from previous experiments that approximately 25% to 30% of the muscle can be activated with stimulation intensities between 120–160 mA.²⁶

Some of the subjects developed muscle spasms during stimulation with intensities that exceeded 120 mA. In these cases, lower stimulation intensities were chosen. Binder-Macleod *et al*²⁷ showed little systematic change in contractile properties with variable stimulation intensities. We used the same stimulation current in the pre- and post-training experiments for each individual and therefore it can be assumed that the intensity of the stimulation current did not materially affect the results of the present study.

Knee extension forces were recorded using a force transducer connected to a non-extensible strap placed around the lower leg. Electrical trains of 1 s duration

were applied to the muscle at stimulation frequencies of 1, 10, 20, 50 and 100 Hz with a rest period of 2 min between the stimulation trains.

The resistance to fatigue was assessed by activating the quadriceps muscle repetitively for 2 min using 30-Hz stimulation trains of 1 s duration every 2 s (ie on: off time=1 s: 1 s). To assess the recovery of the muscle, 1-s stimulation trains of 100 Hz were applied at 0, 2 and 5 min post-exercise.

In order to minimise possible effects of muscle temperature before and after training, room temperature was kept constant at 25°C during the testing




Figure 1 Typical example of a 10-Hz force response before (thin line) and after (bold line) 6 weeks FES-LCE training. Forces are normalised for peak force. dF Represents the average force of three consecutive force oscillations just after the peak force is reached and F_{mean} represents the force during this time




Figure 2 Progression in work output during the FES-LCE training program for seven individuals with SCI. During three training sessions, exercise test protocols were performed on the ergometer, which differed from the original training protocol. Work output values of these sessions are therefore not presented in this graph

procedure and subjects were in the room at least 45 min before the actual protocol started.

Data analysis

Off-line analysis of force records was performed using custom-made software programs. Several indices of contractile speed were obtained. Normalised maximal rate of force rise (MFR) was calculated from the positive filtered (30 Hz filter frequency) force differential and expressed as a percentage of peak force according to the method described by Buller *et al.*²⁸ Half relaxation time ($\frac{1}{2}$ Rt), which is related to cross-

bridge dissociation and calcium re-uptake,²⁹ was defined as the time taken for force to decline from 50% to 25% of the peak 100-Hz force. The degree of fusion as a result of 10 Hz stimulation was determined by calculating the force oscillation amplitudes (FOA) relative to the mean force according to the method described by Gerrits *et al.*²⁶ This was established by taking the average of the three consecutive oscillation amplitudes in the 10-Hz tetanus after force reached its peak value, divided by the mean force during that time (Figure 1).

In order to compare force-frequency relationships, which can distinguish fast-twitch from slow-twitch muscles,³⁰ ratios of 1 and 100-Hz force responses and of 10 and 100-Hz force responses were calculated before and after training.

Force responses resulting from 30-Hz stimulation were continuously recorded during the fatigue proto-




Figure 3 Force (a) and half relaxation time ($\frac{1}{2}$ Rt) (b) expressed as percent of the pre-fatigue value during 2 min of repeated 30-Hz stimulation. Force responses before (open circles, dashed line) and after (closed circles, solid line) 6 weeks of FES-LCE training are plotted every 10 s. $\frac{1}{2}$ Rt values before (black bars) and after (grey bars) training are given of the first (pre-fatigue) and the last (post-fatigue) contraction. Values are mean \pm SD, $n=7$ (a), $n=6$ (b). Missing values are due to stimulus artefacts. *Refers to significantly different from pre-fatigue value, $P<0.05$. †Refers to significantly different from pre-training value, $P<0.05$




Figure 4 Force (a) and half relaxation time ($\frac{1}{2}$ Rt) (b) of 100-Hz responses during recovery from the fatigue protocol before (black bars) and after (grey bars) 6 weeks FES-LCE training. Values at 0 min, 2 min and 5 min post-exercise are expressed as per cent of the pre-fatigue values. Values are mean \pm SD ($n=7$ (a), $n=6$ (b)). Missing values are due to stimulus artefacts. Recovery of force or $\frac{1}{2}$ Rt was not significantly different after training




Figure 5 Contractile properties before (black bars) and after (grey bars) training. Individual data of (a) half relaxation time ($\frac{1}{2}$ Rt), (b) maximal rate of force rise (MFR), (c) relative force oscillation amplitudes (FOA), (d) ratio of 1 and 100-Hz responses (1/100 ratio), and (e) ratio of 10 and 100-Hz responses (10/100 ratio) from subjects 1 to 7. Mean values \pm SD are also shown. Missing values are due to stimulus artefacts. *Refers to significantly different from pre-training values, $P < 0.05$

col. To compare differences in fatigue induced force decline and slowing of relaxation, force and $\frac{1}{2}$ Rt of the 30-Hz responses were calculated at the start and at 2 min of the fatigue protocol. In order to characterise the recovery of the muscle, force and $\frac{1}{2}$ Rt from 100-Hz responses were calculated at 2 min and 5 min after the fatigue protocol.

Statistical analysis

Comparisons of contractile properties before and after FES-LCE training were made using paired *t*-tests. A repeated measures analysis of variance was used to compare the recovery of force and $\frac{1}{2}$ Rt, before and after FES-LCE training. All values are described as mean \pm SD, unless otherwise stated and levels of significance for two-tailed tests were set at 0.05.

Results

FES-LCE exercise training

All subjects completed the total number of 18 training sessions. The majority of the subjects accomplished these sessions within 6 weeks. Two subjects completed the training in longer periods: subject no. 6 in \sim 9 weeks and subject no. 7 in \sim 7.5 weeks. At the start of the training period, most subjects had a very low exercise tolerance to FES-LCE: the mean work output, averaged over the first three training sessions, was 4 ± 5 kJ. Most subjects showed an increase in work output as training progressed, although a large variability between subjects was found. After 6 weeks of training, work output, averaged over the last three training sessions, had significantly improved to 14 ± 13 kJ, respectively ($P=0.019$). The changes in




Figure 6 Force-frequency relationship before (open circles, dotted line) and after (closed circles, solid line) 6 weeks of FES-LCE training. Force responses to different stimulation frequencies are normalised for peak isometric 100-Hz force. Values are mean \pm SD, $n=7$. *Refers to significantly different from pre-training values, $P<0.05$

total work output (kJ) for each subject achieved during the training period is shown in Figure 2. Work output in subject 2 and 6 did not increase during the 6 weeks of FES-LCE. Their total exercise time however, improved from \sim 9 and 7 min, respectively, before training (average of the first three training sessions) to \sim 21 and 13 min, respectively, after training (average of the last three training sessions).

Fatigue and recovery

Expressed as a percentage of the pre-fatigue value, force responses generated during repeated 30-Hz stimulation declined progressively during the fatigue protocol before and after training (Figure 3a). There was a large variability in fatigue resistance between subjects and this was negatively correlated with the time since injury (Pearson correlation coefficient: -0.87 , $P=0.01$). However, quadriceps force declined significantly less ($P=0.012$) following 6 weeks of FES-LCE. At the end of the 2-min stimulation protocol, force declined to $32 \pm 10\%$ of the pre-fatigue value before training and to $42 \pm 10\%$ after training.

Following 2 min of repeated stimulation, $\frac{1}{2}$ Rt increased to $353 \pm 115\%$ of the pre-fatigue value before the FES-LCE training. After the training, the slowing of relaxation was significantly reduced ($200 \pm 24\%$ of the pre-fatigue value, $P=0.015$) (Figure 3b). Figure 4 presents the recovery of force (a) and $\frac{1}{2}$ Rt (b) resulting from 100-Hz responses after the 2-min fatigue protocol.

The relative changes in both force and $\frac{1}{2}$ Rt during recovery were similar before and after 6 weeks of FES-LCE training ($P=0.266$ and 0.545 , respectively).

Contractile properties: contractile speed and low frequency fusion

Figure 5 presents the contractile characteristics before and after 6 weeks FES-LCE training. MFR remained unchanged after training ($1.28 \pm 0.33\% \cdot \text{ms}^{-1}$ before and $1.27 \pm 0.17\% \cdot \text{ms}^{-1}$ after the training program, respectively). However, mean $\frac{1}{2}$ Rt decreased significantly ($P=0.014$) from 39 ± 8 ms before the FES-LCE training program to 33 ± 8 ms after 6 weeks of FES-LCE training. Furthermore, the degree of fusion in a 10-Hz force response diminished, which was represented by a significant increase in FOA from $61 \pm 20\%$ before to $85 \pm 30\%$ after FES-LCE training ($P=0.006$).

Contractile properties: frequency response

The normalised peak force responses, generated at each frequency before and after 6 weeks of FES-LCE training are presented in Figure 6. As can be clearly seen the force-frequency curve shifted to the right at low stimulation frequencies following training. This shift is indicated by a significant decline in the ratio of 1 and 100-Hz responses from 0.27 ± 0.08 to 0.16 ± 0.03 ($P=0.018$), and a significant decline in the ratio of 10

and 100-Hz responses from 0.48 ± 0.05 to 0.35 ± 0.10 ($P = 0.038$) after the training procedure (see also Figure 5d and e, respectively).

Thigh circumference

Six weeks of FES-LCE training did not significantly alter thigh circumference. Thigh girth measured 5 cm proximally of the upper aspect of the patella was 40.2 ± 3.2 cm before, and 40.4 ± 2.5 cm after the training period. Mid-thigh circumference was 47.8 ± 2.9 cm before, and 48.0 ± 3.4 cm after the training.

Discussion

This study was undertaken to quantify changes in the functional characteristics of the quadriceps muscles in individuals with SCI as a result of a 6-week FES-cycling program. This is the first study to demonstrate profound alterations in selected isometric contractile speed properties of the paralysed quadriceps muscle following FES-LCE training. Training increased the fatigue resistance of the quadriceps muscles but to our surprise, contractile speed characteristics adapted towards those of faster muscles.

At the beginning of the training period, most SCI subjects had a very low exercise tolerance during FES-LCE and demonstrated a low resistance to fatigue during 2 min of repeated stimulation. The close negative correlation found between the per cent of pre-fatigue force that remained after the 2-min fatigue protocol and the time since injury occurred, indicates that the fatigue resistance of these muscles decreased with the duration of disuse, which may explain the variation in muscle endurance between subjects. This study clearly provided evidence that fatigue resistance can be improved after only 6 weeks of FES-LCE training since all subjects showed a reduction in the rate at which force declined and a reduction in the slowing of relaxation in response to 2 min of repeated stimulation after the training period. Despite the heterogeneity of the study population, all subjects showed the same changes, indicating that FES-LCE training can improve muscle endurance capacity in a broad population with a large variety of characteristics. Recovery of force and relaxation rate after the 2-min fatigue protocol seemed not affected by training, but the lack of more frequent force measurements during the recovery period limits the interpretation of these results. Martin *et al*² reported higher activities of oxidative enzymes after prolonged electrical stimulation of the tibialis anterior muscle in people with SCI. Furthermore, Chilibeck *et al*¹⁹ found an increase in the capillary number in the quadriceps muscle of subjects with SCI after 8 weeks of FES-LCE. These results indicate that electrical stimulation training programs can increase the oxidative capacity and improve the blood supply to the exercising muscle, which would improve the delivery of oxygen and energy to the

exercising muscles and diminish accumulation of metabolic products. These adaptations are known to be the earliest changes after chronic electrical stimulation³¹ and most likely underlie the improved endurance capacity of the quadriceps muscle in the present study.

In addition, the improved cycle performance, which was also found in previous studies^{3,13–17} is in agreement with the increased endurance capacity of these muscles. Cycle performance is probably best represented by an increased work output, which is determined by the level of pedal resistance multiplied by the time that resistance could be maintained. Two subjects were unable to increase their pedal resistance above 0 kp during the 6-week training program, and thus work output had not improved in these people. However, the total time these individuals were able to cycle doubled after FES-LCE, indicating a substantial improvement in exercise performance.

It is possible that an increased muscle mass as a result of FES-LCE training contributes to the improved work capacity in some of our subjects. We were not able to demonstrate significant increases in thigh girth, which may have been due to the low sensitivity of these measurements, short exercise period or changes in the relative amount of fat tissue. Other investigators however, using computer tomography, MRI, or muscle biopsies, reported an increased whole muscle^{3,14} or fibre cross sectional area¹⁹ after a prolonged period,³ but also after only 8 weeks^{14,19} of FES-LCE training.

The most striking result of the present study is that 6 weeks of FES-LCE increases the speed of relaxation, consequently leading to a decreased fusion of a 10-Hz force response. As relaxation rate increases, force at low frequencies becomes less fused and tetanic fusion would require higher stimulation frequencies resulting in the right-shift of the force-frequency relationship observed in the present study. The increased relaxation rate appears to be inconsistent with our observation of an increased fatigue resistance and in conflict with findings in numerous previous studies. Three decades ago Salmons and Vrbova³² already demonstrated that a transformation of fast- into slow-twitch muscle can be brought about by the application of chronic electrical stimulation in animals and this was confirmed extensively by others.^{21,22} Furthermore, studies that applied electrical stimulation programs on the muscles of humans with SCI, have reported the expected slowing of contractile properties^{25,33} and associated changes in muscle fibre composition towards more oxidative type IIA or type I muscle fibres.^{2,3,20,23,24,33}

The speed of relaxation is closely related to the cross-bridge dissociation and the re-uptake of calcium by the sarcoplasmic reticulum.²⁹ In animals, chronic low frequency electrical stimulation programs reduce the calcium uptake capacity by decreasing the calcium ATPase activity and these changes would be associated

with increases in time-to-peak of isometric tension and half relaxation times.³⁴ The faster relaxation after training found in the present study is in contrast with these findings, and suggests that FES-LCE increased the calcium uptake in the stimulated quadriceps muscle. To our knowledge, it is unknown how long-term SCI in humans affect the mechanisms involving calcium handling. Hunter *et al*³⁵ however, reported that the rate of calcium uptake by the sarcoplasmic reticulum was lower in elderly women compared with young women, and that resistance training increased this rate of calcium uptake in elderly women, but not in young women. It is possible therefore, that although normally, fast-twitch muscles have higher rates of calcium uptake than slow-twitch muscles,³⁶ these processes may be impaired in the fast-twitch paralysed muscles of people with SCI, owing to long periods of inactivity. Previous studies support this speculation by reporting evidence that implies a reduced rate of calcium uptake by the sarcoplasmic reticulum following denervation,³⁷⁻³⁹ although others have found conflicting results.^{40,41} Thus, we would suggest that following prolonged disuse leading to an impairment of calcium handling mechanisms, training initially leads to an increase in the calcium uptake capacity as reflected by the faster relaxation.

In contrast with the changes in the relaxation phase of the muscle, the rate of force development in a 100-Hz force response remained unchanged, suggesting that 6 weeks of FES-LCE training did not affect the net rate of actomyosin cross-bridge attachment.⁴² It should however be noted that higher stimulation frequencies are needed to generate maximal rate of force rise than to produce maximal tetanic tension.^{28,43} Therefore, stimulation frequencies in excess of 100 Hz may have been required to obtain the maximal rate of force development. However, the use of higher stimulation frequencies may increase the risk for damage to the weak paralysed muscles and tendons, and we therefore decided not to use stimulation frequencies above 100 Hz. Moreover, it may be expected that differences in the rate of force development as a result of an increased net cross-bridge attachment would be detected already at sub-maximal levels of activation. A more likely explanation for the unchanged rate of force rise is that although adaptations in metabolic properties, capillary blood flow and calcium ATPase activity occur soon after the start of electrical stimulation programs, alterations in myosin ATPase and associated increased rate of tension development develop at a later stage.^{31,34}

The increased fatigue resistance, faster speed of relaxation and unchanged rate of force development demonstrated in the current study, probably reflect the first adaptations of long-term disused muscles to electrically induced increases in contractile activity.

The quadriceps muscle in our study was stimulated for 30 min, three times a week, for 6 weeks. Other studies however, who actually reported significant adaptations in structural, molecular and physiological properties towards characteristics of slower, more oxidative muscles used considerably longer or more intensive training programs. Andersen *et al*²⁰ and Mohr *et al*³ used a similar FES-LCE training program, but continued the training for 6 to 12 months. Others applied daily electrical stimulation for 1 h a day during approximately 13 weeks²³ or even up to 8 h a day during 6³³ or 24 weeks.² Rochester *et al*^{24,25} used a 4-week training program, which was less than in our study, but again stimulation was applied daily for approximately 2.5 h a day. Thus, it may be speculated that longer or more intensive training periods are required to induce muscular adaptations, which would lead to a detectable slowing of contractile speed characteristics in paralysed muscle of people with SCI.

In addition to altered physiological characteristics of the contractile elements of the muscle, possible mechanical adaptations in the non-contractile tissue of the muscle and/or tendon as a result of training may affect the functional properties of the muscle. Immobilisation leads to an increased muscle stiffness in animals,^{44,45} which may be attributed to both quantitative and qualitative changes in the connective tissue.⁴⁴ Furthermore, contractile activity seems to be important for the maintenance of normal muscle compliance⁴⁶ and training can decrease muscle stiffness.⁴⁷ In contrast, Douglas *et al*⁴⁸ who studied muscle tone in humans, reported a decreased muscle stiffness in people with paraplegia compared with non-injured controls. Furthermore, the muscle stiffness of these paralysed muscles was increased by electrically stimulated shortening contractions and decreased by a period of passive movements.

It remains unclear therefore, whether the FES-LCE training in the present study induced an increase or decrease in muscle stiffness and to what extent these possible mechanical adaptations in muscle and tendon may have affected the results of the present study.

The present study clearly demonstrates that 6 weeks of FES-LCE training can improve the physiological properties of the quadriceps muscle in individuals with SCI with respect to fatigue resistance. Furthermore, an increase in the speed of relaxation was found which might reflect an early response in the calcium handling mechanisms of the muscles consequent upon training of long-term disused muscles. The possible impairment of calcium handling following long-term disuse in humans, and the recovery from that impairment is clearly an issue which deserves further investigation if monitoring of training programs is used to inform the development of effective interventions in this group of SCI individuals. It is possible that longer training programs are required to induce significant adaptations towards characteristics of slower contracting muscles.

References

- 1 Davis GM. Exercise capacity of individuals with paraplegia. *Med Sci Sports Exerc* 1993; **25**: 423–432.
- 2 Martin TP, Stein RB, Hoepfner PH, Reid DC. Influence of electrical stimulation on the morphological and metabolic properties of paralyzed muscle. *J Appl Physiol* 1992; **72**: 1401–1406.
- 3 Mohr T et al. Long-term adaptation to electrically induced cycle training in severe spinal cord injured individuals [published erratum appears in *Spinal Cord* 1997 Apr;35(4):262]. *Spinal Cord* 1997; **35**: 1–16.
- 4 Burnham R et al. Skeletal muscle fibre type transformation following spinal cord injury. *Spinal Cord* 1997; **35**: 86–91.
- 5 Grimby G, Broberg C, Krotkiewska I, Krotkiewski M. Muscle fiber composition in patients with traumatic cord lesion. *Scand J Rehabil Med* 1976; **8**: 37–42.
- 6 Lotta S et al. Morphometric and neurophysiological analysis of skeletal muscle in paraplegic patients with traumatic cord lesion. *Paraplegia* 1991; **29**: 247–252.
- 7 Biering-Sorensen F, Bohr H, Schaadt O. Bone mineral content of the lumbar spine and lower extremities years after spinal cord lesion. *Paraplegia* 1988; **26**: 293–301.
- 8 Ragnarsson KT, Sell GH. Lower extremity fractures after spinal cord injury: a retrospective study. *Arch Phys Med Rehabil* 1981; **62**: 418–423.
- 9 Hopman MT, van Asten WN, Oeseburg B. Changes in blood flow in the common femoral artery related to inactivity and muscle atrophy in individuals with long-standing paraplegia. *Adv Exp Med Biol* 1996; **388**: 379–383.
- 10 Nash MS, Montalvo BM, Applegate B. Lower extremity blood flow and responses to occlusion ischemia differ in exercise-trained and sedentary tetraplegic persons. *Arch Phys Med Rehabil* 1996; **77**: 1260–1265.
- 11 Taylor PN et al. Limb blood flow, cardiac output and quadriceps muscle bulk following spinal cord injury and the effect of training for the Odstock functional electrical stimulation standing system. *Paraplegia* 1993; **31**: 303–310.
- 12 Petrofsky JS, Heaton HHD, Phillips CA. Leg exerciser for training of paralyzed muscle by closed-loop control. *Med Biol Eng Comput* 1984; **22**: 298–303.
- 13 Bremner LA et al. A clinical exercise system for paraplegics using functional electrical stimulation. *Paraplegia* 1992; **30**: 647–655.
- 14 Hjeltnes N et al. Improved body composition after 8 wk of electrically stimulated leg cycling in tetraplegic patients. *Am J Physiol* 1997; **273**: R1072–1079.
- 15 Faghri PD, Glaser RM, Ficoni SF. Functional electrical stimulation leg cycle ergometer exercise: training effects on cardiorespiratory responses of spinal cord injured subjects at rest and during submaximal exercise. *Arch Phys Med Rehabil* 1992; **73**: 1085–1093.
- 16 Ragnarsson KT. Physiologic effects of functional electrical stimulation-induced exercises in spinal cord-injured individuals. *Clin Orthopaed Relat Res* 1988; **233**: 53–63.
- 17 Sloan KE et al. Musculoskeletal effects of an electrical stimulation induced cycling programme in the spinal injured. *Paraplegia* 1994; **32**: 407–415.
- 18 Hangartner TN, Rodgers MM, Glaser RM, Barre PS. Tibial bone density loss in spinal cord injured patients: effects of FES exercise. *J Rehabil Res Dev* 1994; **31**: 50–61.
- 19 Chilibeck PD et al. Histochemical changes in muscles of individuals with spinal cord injury following functional electrical stimulated exercise training. *Spinal Cord* 1999; **37**: 264–268.
- 20 Andersen JL et al. Myosin heavy chain isoform transformation in single fibres from m. vastus lateralis in spinal cord injured individuals: effects of long-term functional electrical stimulation (FES). *Pflugers Archiv Eur J Physiol* 1996; **431**: 513–518.
- 21 Pette D, Vrbova G. Adaptation of mammalian skeletal muscle fibers to chronic electrical stimulation. *Rev Physiol Biochem Pharmacol* 1992; **120**: 115–202.
- 22 Salmons S. Exercise, stimulation and type transformation of skeletal muscle. *Int J Sports Med* 1994; **15**: 136–141.
- 23 Greve JM et al. Functional electrical stimulation (FES): muscle histochemical analysis. *Paraplegia* 1993; **31**: 764–770.
- 24 Rochester L et al. Influence of electrical stimulation of the tibialis anterior muscle in paraplegic subjects. 2. Morphological and histochemical properties. *Paraplegia* 1995; **33**: 514–522.
- 25 Rochester L et al. Influence of electrical stimulation of the tibialis anterior muscle in paraplegic subjects. 1. Contractile properties. *Paraplegia* 1995; **33**: 437–449.
- 26 Gerrits HL et al. Contractile properties of the quadriceps muscle in individuals with spinal cord injury. *Muscle Nerve* 1999; **22**: 1249–1256.
- 27 Binder-Macleod SA, Halden EE, Jungles KA. Effects of stimulation intensity on the physiological responses of human motor units. *Med Sci Sports Exerc* 1995; **27**: 556–565.
- 28 Buller AJ, Lewis DM. The rate of tension development in isometric tetanic contractions of mammalian fast and slow skeletal muscle. *J Physiol* 1965; **176**: 337–354.
- 29 Edwards RH, Hill DK, Jones DA. Metabolic changes associated with the slowing of relaxation in fatigued mouse muscle. *J Physiol* 1975; **251**: 287–301.
- 30 Cooper S, Eccles JC. The isometric responses of mammalian muscles. *J Physiol* 1930; **69**: 377–385.
- 31 Salmons S, Henriksson J. The adaptive response of skeletal muscle to increased use. *Muscle Nerve* 1981; **4**: 94–105.
- 32 Salmons S, Vrbova G. The influence of activity on some contractile characteristics of mammalian fast and slow muscles. *J Physiol* 1969; **201**: 535–549.
- 33 Stein RB et al. Optimal stimulation of paralyzed muscle after human spinal cord injury. *J Appl Physiol* 1992; **72**: 1393–1400.
- 34 Pette D. JB Wolffe memorial lecture. Activity-induced fast to slow transitions in mammalian muscle. *Med Sci Sports Exerc* 1984; **16**: 517–528.
- 35 Hunter SK et al. Human skeletal sarcoplasmic reticulum Ca²⁺ uptake and muscle function with aging and strength training. *J Appl Physiol* 1999; **86**: 1858–1865.
- 36 Pette D, Staron RS. Cellular and molecular diversities of mammalian skeletal muscle fibers. *Rev Physiol Biochem Pharmacol* 1990; **116**: 1–76.
- 37 Howell S, Zhan WZ, Sieck GC. Diaphragm disuse reduces Ca²⁺ uptake capacity of sarcoplasmic reticulum. *J Appl Physiol* 1997; **82**: 164–171.
- 38 Lehotsky J, Bezakova G, Kaplan P, Raeymaekers L. Distribution of Ca(2+)-modulating proteins in sarcoplasmic reticulum membranes after denervation. *Gen Physiol Biophys* 1993; **12**: 339–348.
- 39 Schulte L et al. Sarcoplasmic reticulum Ca²⁺ pump expression in denervated skeletal muscle. *Am J Physiol* 1994; **267**: C617–622.
- 40 Sreter FA. Effect of denervation on fragmented sarcoplasmic reticulum of white and red muscle. *Exp Neurol* 1970; **29**: 52–64.
- 41 Takekura H, Kasuga N, Kitada K, Yoshioka T. Morphological changes in the triads and sarcoplasmic reticulum of rat slow and fast muscle fibres following denervation and immobilization. *J Muscle Res Cell Motil* 1996; **17**: 391–400.
- 42 Ranatunga KW. Temperature-dependence of shortening velocity and rate of isometric tension development in rat skeletal muscle. *J Physiol* 1982; **329**: 465–483.
- 43 de Haan A. The influence of stimulation frequency on force-velocity characteristics of in situ rat medial gastrocnemius muscle. *Exp Physiol* 1998; **83**: 77–84.
- 44 Williams PE, Goldspink G. Connective tissue changes in immobilised muscle. *J Anatomy* 1984; **138**: 343–350.
- 45 Gillette PD, Fell RD. Passive tension in rat hindlimb during suspension unloading and recovery: muscle/joint contributions. *J Appl Physiol* 1996; **81**: 724–730.
- 46 Williams PE, Catanese T, Lucey EG, Goldspink G. The importance of stretch and contractile activity in the prevention of connective tissue accumulation in muscle. *J Anatomy* 1988; **158**: 109–114.

- 47 Pousson M, Perot C, Goubel F. Stiffness changes and fibre type transitions in rat soleus muscle produced by jumping training. *Pflugers Archiv Eur J Physiol* 1991; **419**: 127–130.
- 48 Douglas AJ *et al*. The effects of neuromuscular stimulation on muscle tone at the knee in paraplegia. *Exp Physiol* 1991; **76**: 357–367.
- 49 Maynard Jr FM *et al*. International standards for neurological and functional classification of spinal cord injury. American Spinal Injury Association. *Spinal Cord* 1997; **35**: 266–274.