inconsistent with proposed kinetic mechanisms. This raises some interesting questions on the chemical mechanism of compound I formation.

We thank Hui-Chen J. Lai and Nan-gian Li for their help throughout the study and R. C. Hardison and T H. Kao for reviewing the manuscript. Amino-acid sequence and composition were determined at University of California at Los Angeles Medical School and University of Wisconsin Biophysics Laboratory, respectively. This work was supported in part by a Biomedical Research Support Grant from The Pennsylvania State University and a USDA grant to M.T. M.T. is recipient of a Presidential Young Investigator Award from NSF. C .-P.D.T. is recipient of a Research Career Develoment Award from USPHS. The sequence data in this publication have been submitted to the EMBL/GenBank Libraries under the accession number Y00262.

Received 25 November 1986; accepted 29 January 1987.

- 1. Tien. M. & Kirk. T. K. Science 221, 661-663 (1983).
- 2. Glenn, J. K., Morgan, M. A., Mayfield, M. B., Kuwahara, M. & Gold, M. H. Biochem biophys. Res. Commun. 114, 1077-1083 (1983).
- Tien, M. & Kirk, T. K. Proc. natn. Acad. Sci. U.S.A. 81, 2280-2284 (1984).
- 4. Gold, M. H., Kuwahara, M., Chiu, A. A. & Glenn, J. K. Archs Biochem. Biophys. 234, 353-362 (1984)
- Bumpus, J. A., Tien, M., Wright, D. & Aust, S. D. Science 228, 1434-1436 (1985).
- 6. Kirk, T. K., Croan, S. C., Tien, M., Murtagh, K. & Farrel, R. Enzyme Microb. Technol. 8, 27-32 (1985).
- 7. Gubler, U. & Hoffman, B. J. Gene 25, 263-269 (1983).
- Tien, M. & Kirk, T. K. Meth. Enzym. (in the press).
 Maniatis, T., Fritsch, E. F. & Sambrook, J. Molecular Cloning, a Laboratory Manual (Cold Spring Harbor Laboratory, New York, 1982).
- Sanger, F., Nicklen, S. & Coulson, A. R. Proc. natn. Acad. Sci. U.S.A. 74, 5463-5467 (1977).
 Messing, J. Meth. Enzym. 101, 20-77 (1983).
 Kozak, M. Nucleic Acids Res. 9, 5233-5252 (1981).

- Kaput, J., Goltz, S. & Blobel, G. J. biol. Chem. 257, 15054-15058 (1982).
 Julius, D., Schekman, R. & Thorner, J. Cell 36, 309-318 (1984).
- 15. Bell, G. I., Santerre, R. F. & Mullenbach, G. T. Nature 302, 716-718 (1983).
- 16. Neuberger, A., Gottshalk, A., Marshal, R. D. & Spiro, R. D. in The Glycoproteins: Their

- Composition, Structure and Function Part A (ed. Gottschalk, A.) 450-490 (Elsevier, Amsterdam, 1972).
- 17 Proudfoot N 1 & Brownbee G G Nature 263 211-214 (1976)
- 18. Pelham, H. R. B. & Jackson, R. J. Eur. J. Biochem. 67, 247-256 (1976).
- Tien, M., Kirk, T. K., Bull, C. & Fee, J. A. J. biol. Chem. 261, 1687-1693 (1986).
 Kuila, D., Tien, M., Fee, J. A. & Ondrias, M. R. Biochemistry 24, 3394-3397 (1986).
- 21. Poulos, T. L. et al. J. biol. Chem. 255, 575-580 (1986).
- Dunford, H. B. & Stillman, J. S. Coord. Chem. Rev. 19, 187-251 (1976).
 Poulos, T. L. & Kraut, J. J. biol. Chem. 255, 8199-8205 (1980).
- 24. Chance, B. Arch. Biochem. Biophys. 41, 416-424 (1952).
- Manthey, J. A. & Hager, L. P. J. biol. Chem. 260, 9654-9659 (1985).
 Zhang, Y. Z., Zylstra, G. J., Olsen, R. H. & Reddy, C. A. Biochem. biophys. Res. Commun. 137, 649-656 (1986)
- Chirgwin, J. M., Przybyla, A. E., MacDonald, R. J. & Rutter, W. J. Biochemistry 18, 5294-5299 (1979).
- Bantle, J. A., Maxwell, I. H. & Hahn, W. E. Analyt. Biochem. 72, 413-427 (1976)
- 29. Tu, C.-P. D., Weiss, M. J., Karakawa, W. W. & Reddy, C. C. Nucleic Acids Res. 10, 5407-5419 (1982).
- Laemmli, U. K. Nature 227, 680-685 (1970).
- Towbin, H., Staehelin, T. & Gordon, J. Proc. nam. Acad. Sci. U.S.A. 76, 4350-4354 (1979).
 Welinder, K. G. & Mazza, G. Eur. J. Biochem. 73, 353-358 (1977).
- 33. Welinder, K. G. FEBS Lett. 72, 19-23 (1976).

Corrigendum

A cDNA clone encoding an S-locus-specific glycoprotein from Brassica oleracea

J. B. Nasrallah, T.-H. Kao, M. L. Goldberg & M. E. Nasrallah Nature 318, 263-267 (1985).

FURTHER work on the sequence analysis of λ gt 10 cDNA clones related to pBOS5 has revealed that the nucleotide sequence of the cDNA clone should have:

- an insertion of a C after position 333;
- a change from CG to GC at positions 347-348;
- an insertion of G after position 766;
- an insertion of an A after position 808.

The reading frame is correct as published but the above insertions extend it to the TAG at position 1,252-1,254 raising the number of amino acid residues coded by this cDNA to 418.

MATTERS ARISING

Evidence for climatic attractors

GRASSBERGER1 draws attention to some problems related to the estimate of attractor dimensions in the case of sparse data sets. As parts of this paper may appear to have a controversial character for the unaware reader, in reference to our previous work2 we believe that some comments are in order.

It is surprising that with a number of points as small as 230 for the V28-238 core, Grassberger goes to phase spaces of dimension up to 11. In our own work with ~500 data points we reluctantly went up to dimension six, and considered that beyond this value the results were not reliable. In particular, the estimate of the slopes becomes increasingly difficult—in fact after seeing Grassberger's paper we repeated our computations by taking only one out of two or out of three data points and found no significant trend toward saturation for embedding phase space dimension up to six. We believe therefore that the conclusions drawn by Grassberger with so few data points are subject to caution. In addition it seems unlikely that the V28-239 core can be used for dimension estimates. There are very few raw data points, considering that they span a time interval twice as long as the V28-238

core. Moreover, the measurements beyond 106 yr are less reliable (P. Pestiaux, personal communication).

Dr Grassberger minimizes the role of time lag τ . This may be theoretically correct, but in practice many researchers have pointed out the importance of the choice of τ in the calculation of attractor dimensions^{3,4}. The argument appealing to correlation time is of course legitimate, but in many cases this time may be hard to estimate. It is unfortunate that no specific references regarding these points are provided by the author.

Dr Grassberger also warns against spuriously small dimension estimates arising from smoothed data. It should be realized that in real world complex systems direct access to physical variables may be impossible. In geology in particular, the connection between data (which refer to depth along the core) and physical variables (like temperature or ice volume) is established by appealing to a model. Depending on the type of model (which describes, among others, the sedimentation mechanism) a resolution of for instance, 2,000 yr (our data) or 5,000 yr (Grassberger's data) is achieved. In either case, however, an interpolation is involved, since the raw data need not be equidistant. We regard this as a fact of

life. As all current work on palaeoclimatology is based on such data, we do not see why we should refrain from using

In summary, these issues show how careful one has to be in discussing the behaviour of real world complex systems. They also highlight the need for abundant, high quality geological data. We iterate our strong conviction that climatic attractors do exist, a conviction that is strengthened by independent dimension estimates reported recently by Fraedrich⁵ and Saltzman and co-workers6.

C. NICOLIS

Institut d'Aéronomie Spatiale de Belgique, Avenue Circulaire, 3 1180 Bruxelles, Belgium

G. NICOLIS

Faculté des Sciences de Université Libre de Bruxelles, Campus Plaine, CP 226, Boulevard du Triomphe. 1050 Bruxelles, Belgium

- 1. Grassberger, P. Nature 323, 609-612 (1986).
- Nicolis, C. & Nicolis, G. Nature 311, 529-532 (1984).
- Wolf, A., Swift, J. B., Swinney, H. L. & Vastano, J. A.,
 Physica D16, 285-317 (1985).
 Malmaison, B., Atten, P., Bergé, P. & Dubois, M. J. phys.
- Lett. Paris 44, L897-L902 (1983). Fraedrich, K. J. atmos. Sci. 43, 419-432 (1986).
- Saltzman, B. in Irreversible Phenomena and Dynamical Systems Analysis in Geosciences (Reidel, Dordrecht, 1987).