Fig. 3 Mechanism of action of dipeptide transport forms of a KDO analogue, CMP-KDO synthetase inhibitor in *E. coli*. Ala, Alanine; I, 8-amino-2,6-anhydro-3,8-dideoxy-D-glycero-D-talo-octonic acid; opp, tpp, oligo- and tripeptide permeases; CKS, CMP-KDO synthetase.

After overnight incubation at 37 °C on plates bearing a wild-type *E. coli* W (M26-26) possessing all three permeases, a zone of inhibition (diameter 38 mm) was seen around the disk. Zones of similar diameter were seen around disks placed on permease-deficient mutants of *E. coli* W such as PA 0113 (*dpp*⁻) and PA 0114 (*tpp*⁻), but on plates containing *E. coli* PA 0112 (*opp*⁻), PA 0123 (*dpp*⁻ *opp*⁻) and PA 0128 (*tpp*⁻ *opp*⁻) no inhibition zones were visible. Similarly, an *E. coli* K12 wild type was also sensitive (zone diameter 41 mm), but an *opp*⁻ derivative of it, PA 0183, was sensitive to some degree (zone diameter 13 mm). These data suggest that although the bulk of Ala-Ala-NHdKDO uptake in *E. coli* is by the oligopeptide permease, the compound may also be able to use, to a lesser extent, the *tpp* system of *E. coli* K12 strains.

In the biosynthesis of LPS the donation of KDO from CMP-KDO to an acylated and phosphorylated $\beta(1\rightarrow6)$ -linked D-glucosamine disaccharide known as lipid A precursor¹⁵⁻¹⁷ is an essential prerequisite for the secondary acylation of lipid A, incorporation of core and O-side chain sugars, and the translocation of LPS to the outer membrane^{2,3,15}. At the non-permissive temperature, although *kdsA* and *kdsB* mutants of *Salmonella typhimurium* are unable to produce CMP-KDO, synthesis of lipid A precursor continues unabated. Because the latter cannot be rapidly translocated to the cell exterior^{3,16}, it accumulates, inhibiting growth and ultimately causing the death of the bacterium^{3,15}. *S. typhimurium* AG 701150 (*kdsA*^{ts}) was grown at its permissive and non-permissive temperatures in the presence of N-acetyl-D-[1-¹⁴C]glucosamine, the LPS extracted and separated by TLC. Contact autoradiography (Fig. 2) revealed that at the non-permissive temperature large amounts of lipid A precursor accumulate at the expense of complete LPS. Treatment of the bacterium with Ala-Ala-NHdKDO or Pro-Met-NHdKDO also leads to the accumulation of lipid A precursor.

Lethal levels of these agents had no effect on protein, cell wall, RNA, or DNA synthesis, as measured by incorporation of radiolabelled leucine, diaminopimelic acid, uracil, or adenine into the relevant cellular macromolecules (data not shown). The diagram of LPS biosynthesis in *E. coli* shown in Fig. 3 indicates the mechanisms of drug uptake and hydrolysis and the consequences of subsequent inhibition of CMP-KDO synthetase.

We thank M. J. Osborn for mutants defective in KDO biosynthesis, J. W. Payne for help in determining the mechanism of

peptide uptake and C. Sahlberg for synthesizing the radioactive compounds.

Received 1 April; accepted 27 April 1987.

1. Claesson, A., Luthman, K., Gustafsson, K. & Bondesson, G. *Biochem. biophys. Res. Commun.* **143**, 1063-1068 (1987).
2. Hammond, S. M. in *The Bacterial Cell Surface* (eds Hammond, S. M., Lambert, P. A. & Rycroft, A. N.) 57-82 (Croom Helm, London, 1984).
3. Osborn, M. J. in *The Bacterial Outer Membrane* (ed. Inouye, M.) 15-34 (Wiley, New York, 1979).
4. Högenauer, G. & Woisetschlag, M. *Nature* **293**, 662-664 (1981).
5. Rick, D. & Osborn, M. J. *Proc. natn. Acad. Sci. U.S.A.* **69**, 3756-3760 (1972).
6. Ghalambor, M. A. & Heath, E. C. *J. biol. Chem.* **241**, 3216-3221 (1966).
7. Fickel, T. E. & Gilvarg, C. *Nature* **241**, 161-163 (1973).
8. Ames, B. N. et al. *Proc. natn. Acad. Sci. U.S.A.* **70**, 456-458 (1973).
9. Ringrose, P. S. in *The Scientific Basis of Antimicrobial Chemotherapy* (eds Greenwood, D. & O'Grady, F.) 219-281 (Cambridge University Press, 1985).
10. Berges, D. A. et al. *J. med. Chem.* **29**, 89-95 (1986).
11. Hemsdorf, C. L. & Simmons, C. in *Microorganisms and Nitrogen Sources* (ed. Payne, J. W.) 301-333 (Wiley, New York, 1980).
12. Gibson, M. M., Price, M. & Higgins, C. F. *J. Bact.* **160**, 122-130 (1984).
13. Higgins, C. F. & Gibson, M. M. *Meth. Enzym.* **125**, 365-377 (1986).
14. Payne, J. W. *Biochem. Soc. Trans.* **11**, 794-797 (1983).
15. Rick, P. D., Fung, W.-M., Ho, C. & Osborn, M. J. *J. biol. Chem.* **252**, 4904-4912 (1977).
16. Rick, P. D. & Young, D. A. *J. Bact.* **150**, 456-464 (1982).
17. Raetz, C. R. H., Purcell, S., Meyer, M. V., Quershi, N. & Takayama, K. *J. biol. Chem.* **260**, 16080-16088 (1985).

Errata

Electroreceptors in the platypus

J. E. Gregory, A. Iggo, A. K. McIntyre & U. Proske
Nature **326**, 386-387 (1987).

IN quoting the work of Scheich *et al.*, who measured the threshold voltage gradient for behavioural responses in the platypus, a value of 60 V cm⁻¹ was given. The value should have been 60 μ V cm⁻¹.

Abolition of the expression of inhibitory guanine nucleotide regulatory protein G_i in diabetes

D. Gawler, G. Milligan, A. M. Spiegel, C. G. Unson & M. D. Houslay
Nature **327**, 229-232 (1987).

AN error in the legend to Fig. 2 should be corrected as follows: Membranes are from mature animals in control (lane c), streptozotocin-diabetic (lane a) and streptozotocin-diabetic animals undergoing insulin therapy (lane b) states: DF, dye front.