## H-form DNA and the hairpin-triplex model

SIR-In reporting the occurrence of a novel DNA form, the 'H-form', in supercoiled homopurine-homopyrimidine tracts at low pH, Mirkin et al.1 apparently ignored our previous description of such a structure (the 'hairpin-triplex')<sup>2</sup>. We came up with this model when analysing the precise positions where the enzyme S<sub>1</sub>-nuclease was able to digest a cloned homopurine-homopyrimidine element, normally lying upstream from the human thyroglobulin gene.

The proposed structure, virtually identical to the one depicted in Fig. 3 of ref. 1, was suggested to us by the work of Lee et al.<sup>3</sup> who showed that  $(TC)_{u} \cdot (GA)_{u}$ duplexes could form triple-helical complexes at low pH. Careful examination of data available at that time concerning the digestion of other homopurinehomopyrimidine sequences with S<sub>1</sub>nuclease, as for example the one located in the human  $\alpha_2 - \alpha_1$  globin intergenic region<sup>4</sup>, showed that they also fitted into the proposed model.

We therefore argued<sup>2</sup> that our hairpintriplex structure represented "an alternative to the [then favoured] slippagemechanism hypothesis5 as long as the distance between the repeats still allows the formation of a double-stranded hairpin<sup>2</sup>. As well as setting the facts in their true light, we consider that mention of our data in the context of the work of Mirkin et al. is worthwhile, as both point to the same model while stemming from different methodological approaches.

DANIEL CHRISTOPHE

## IRIBHN,

Campus Erasme, bat. C, 808, route de Lennik, 1070 Bruxelles, Belgium

- Mirkin, S.M. et al. Nature 330, 495–497 (1987).
  Christophe D. et al. Nucleic Acids Res. 13, 5127–5144 (1985).
  Lee J.S., Johnson D.A. & Morgan A.R. Nucleic. Acids Res. 6. 3073-3091 (1979).
- Sor5-3091 (1979).
  Shen C.-K.J. Nucleic Acids Res. 11, 7899–7910 (1983).
  Mace H.A.F., Pelham H.R.B. & Travers A.A. Nature 304,
- 555-557 (1983)

FRANK-KAMENETSKII REPLIES --- Although our *Nature* paper<sup>1</sup> did not cite the paper by Christophe et al.2, its two predecesors3.4 both did. In ref. 3, where we formulated our final model of the H-form, we wrote<sup>3</sup>: "To explain the bimodal cleavage pattern Christophe et al.... also invoked the triple helix." We also properly cited the paper by Lee et al. which Christophe mentions in his letter.

We disagree that the structure claimed in ref. 2 is "virtually identical to the one depicted in Fig. 3 of Mirkin et al.". The Christophe et al. structure is virtually identical the structure proposed by Lee et al. (see Fig. 5 of ref. 5), which we knew before starting the whole business. Indeed, we wrote in the introduction to our very first paper on the subject<sup>6</sup>: "Slippage loops

## SCIENTIFIC CORRESPONDENCE-

., a left-handed helical structure ... triple and quarter helices . . . have been discussed in the literature." The principal difference between the structure of Lee et al. and Christophe et al. on the one hand, and our structure on the other, is the nature of the hairpin. According to the former it is double-stranded, whereas according to the latter it is single-stranded.

Christophe et al.2 invoked the 'hairpintriplex' structure to explain the bimodal pattern of cleavage of their homopurinehomopyrimidine tract by S<sub>1</sub>-endonuclease. The two sites of cleavage are more than 100 base pairs apart and they considered that this 'spacer' region forms a doublestranded hairpin. On the basis of our recent findings<sup>2</sup>, another interpretation of their data seems more plausible. We have found that the two sites of cleavage are actually H-palindromes (underscored in the figure).

<sup>46</sup>CGAAGGAAGGAAAGAACGAAGAAAGAAAAG

<sup>155</sup>AAAGGAAGAAAGGAAGGAAGGAAGGAAGGA

## **Species selection and the** role of the individual

SIR-In opposing the punctuational model of evolution and species selection and in his reply to Eldredge and Gould<sup>2</sup>, Maynard Smith has seriously misrepresented my writings on the subject, making my position appear far more radical than it has ever been.

In promoting the concept of species selection, I have never denied that selection at the level of the individual plays an important role in speciation. In 1979, in addressing quantum speciation, I wrote<sup>3</sup>. "It is guite likely that in the punctational model natural selection will continue to be viewed as the dominant guiding process of generation-by-generation change". And in 1982, I wrote (in italics for emphasis)4: "The punctational model is essentially a description of temporal and spatial patterns, not of population genetic mechanisms, and it is compatible with the possibility that virtually all of the evolution that occurs during divergents speciation is achieved by conventional natural selection among individuals". None of my other writings has been intended to express any other view. Thus, Maynard Smith is mistaken in his claim that the raw material of species selection, as I have defined the process<sup>3-5</sup>, relies strictly on "non-adaptive punctational changes that occur when new species arise"1.

The definition of the word random, as applied to speciation, seems to be at the root of the misunderstanding. Maynard Smith<sup>2</sup> has guoted out of context a statement of mine that refers to "a strong random element" in speciation. Randomness here does not refer to non-adaptive genetic drift or haphazard mutation, as

We therefore believe that these two Hpalindromes, under superhelical stress and at acid pH, form two separate H-forms and these are the S<sub>1</sub>-hypersensitive sites. These H-palindromes are, at least, very similar to those collected in Table 1 of our Nature paper<sup>1</sup>. The interpretation of the S, cleavage pattern in terms of the model of Christophe et al. seems doubtful for several reasons, which are beyond the scope of this short reply. The above discussion clearly shows that these different models entail different interpretations of the same experimental data.

MAXIM FRANK-KAMENETSKII Institute of Molecular Genetics, USSR Academy of Sciences. Moscow 123182, USSR

- Mirkin, S.M. et al. Nature 330, 495–497 (1987).
  Christophe, D. et al. Nucleic Acids Res. 13, 5127–5144 (1985).
- Lyamichev, V.I., Mirkin, S.M. & Frank-Kamenetskii, M.D. I biomolec Struct Dyn 3 667-669 (1986)
- 4. Lyamichev, V.I., Mirkin, S.M. & Frank-Kamenetskii, M.D. J. biomolec. Struct. Dyn. 5, 275-282 (1987).
- Lee, J.S., Woodsworth, M.L., Latimer, L.J.P. & Morgan, A.R. Nucleic Acids Res. 12, 6603–6614 (1984).
- Lyamichev, V.I., Mirkin, S.M. & Frank-Kamenetskii, M.D. J.biomolec.Struct.Dyn. 3, 327-338 (1985).

he seems to have concluded. Again, for historical accuracy, I quote one of my previous statements<sup>4</sup> (omitting the references):

"The random element in speciation to a considerable degree decouples macroevolution from microevolution. The meaning of "random element" has frequently been misunderstood. Although there is often a bias in the direction of speciation within the higher taxon morphogenetic or environmental а tendancy for certain changes to occur rather than others - the direction that the next event will take in any segment of phylogeny nonetheless remains highly unpredictable: a pair of dice may be loaded, but each one still has six sides. Where, when, and in what subenvironment the next speciation event will occur within a lineage is always highly uncertain. Mann has discussed this kind of randomness in the operation of geologic processes. Species selection can operate even without the direction of a speciation event being entirely random. The only requirement is that speciation generate a variety of species upon which selection can operate. As characterized by Mayr, speciation amounts to experimentation by which evolution continually probes the environment. The origin of a particular kind of species at a particular time and place is no guarantee that this kind of species will fare well in a broader environmental context."

It is also important to recognize that selection at the level of the species is not predicted upon the punctuational model. Species selection simply looms larger in this model than in a gradualistic framework, which grants gradual evolution a larger role in the determination of macroevolutionary trends. In fact, although he