


FIG. 2 The measured values of  $\Phi_{ks}$  and  $\Phi_v - \bar{V}_M$  of the sodium salts of oligo- and polynucleotides as a function of the percentage of A+T base pairs. The measurements were made at 1 °C in CsCl (0.2 mol l⁻¹) HEPES buffer (2 mmol l⁻¹), pH 7.8, at concentrations of ~1 mg ml⁻¹. Numbers 1 to 6, deoxyoligonucleotide sequences described in Fig. 1; 7, salmon-sperm DNA; 8, the alternating poly[d(A-T)] · poly[d(A-T)]; and 9, the homopolymer poly(dA) · poly(dT). Note that generally, the lower the values of both  $\Phi_{ks}$  and  $\Phi_v - \bar{V}_M$ , the higher the degree of hydration. All values are calculated per mole of nucleotide. The details of the measurements and preparative procedure are described elsewhere<sup>12,14</sup>. The values of  $\bar{V}_M$  were taken from ref. 18 and the intrinsic volume of sodium ions was not included in the calculations.

both  $\Phi_{ks}$  and  $\Phi_v - \bar{V}_M$  obtained with the oligo(dA) · oligo(dT) and the poly(dA) · poly(dT) duplexes. An important observation is that changing the sequence of the molecule from



resulted in a large decrease in its hydration. These structures differ in only one base-pair stacking contact in the middle of the helix.

The evidence presented here indicates that (dA)<sub>n</sub> · (dT)<sub>n</sub> tracts have an anomalously high hydration in dilute solution. An enhanced hydration has been proposed to be one of major determinants of the anomalous properties of (dA)<sub>n</sub> · (dT)<sub>n</sub><sup>21,22</sup>.

14. Buckin, V. A., Kankiya, B. I., Sarvazyan, A. P. & Uedaira, H. *Nucleic Acids Res.*, in the press.
15. Lo Surdo, A., Shin, Ch., Miller, F. J. *J. chem. Engng. Data* **23**, 187–201 (1978).
16. Ivanova, E. M., Kutiavina, I. V., Pletnev, A. G. & Shamanin, V. A. *Bioorg. Chem.* **8**, 1501–1515 (1982).
17. Maxam, A. G. & Gilbert, W. *Meth. Enzym.* **65**, 499–560 (1986).
18. Pavlov, M. Ya. & Fedorov, B. A. *Biofizika* **28**, 931–936 (1983).
19. Tunis, M. J. B., Hearst, J. E. *Biopolymers* **6**, 1325–1344 (1968).
20. Mrevlishvili, G. M. *Dokl. Akad. Nauk. SSSR* **260**, 761–764 (1981).
21. Chuprina, V. P. *FEBS Lett.* **186**, 98–102 (1985).
22. Chuprina, V. P. *FEBS Lett.* **195**, 363 (1986).

## ERRATA

### Creation of the Uranus rings and dust bands

L. W. Esposito & Joshua E. Colwell

*Nature* **339**, 605–607 (1989)

An error was introduced during the processing of this letter. The two sentences beginning seven lines below equation 3 on page 606 should read: 'The satellites 1986U1–9 range from 20 to 40 km in radius. We consider this a confirmation, but it shows no contradiction between the model disruption rate (calculated from the crater<sup>5</sup>) and the observed moons near the rings.'

### Triple-strand formation in the homopurine:homopyrimidine DNA oligonucleotides d(G-A)<sub>4</sub> and d(T-C)<sub>4</sub>

Ponni Rajagopal & Juli Feigon

*Nature* **339**, 637–640 (1989)

In Fig. 2a of this letter, the A·T base pairs are mislabelled as G·T. In Fig. 2b the indication of Hoogsteen base pairs is omitted, but these are correctly labelled in Fig. 3. Also in Fig. 2b, the 'H-bonded aminos' label should read 'C<sup>+</sup> aminos'.

### Inhibition of exocytosis by intracellularly applied antibodies against a chromaffin granule-binding protein

Felix E. Schweizer, Theo Schäfer, Carlo Tapparelli\*, Marianne Grob, Urs O. Karli, Rolf Heumann\*, Hans Thoenen\*, Richard J. Bookman† & Max M. Burger

Friedrich Miescher Institut, Postfach 2543, CH-4002 Basel, Switzerland

† Howard Hughes Medical Institute, Department of Physiology, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA

\* Present addresses: Sandoz Ltd, Preclinical Research, CH-4002 Basel, Switzerland (C.T.); Abteilung Neurochemie, Max-Planck-Institut für Psychiatrie, D-8033 Martinsried, FRG (R.H. and H.T.)

*Nature* **339**, 709–712 (1989)

In this letter in the 29 June issue, the affiliations of the authors were unclear in the printed version. The correct details appear above.

Received 8 March; accepted 19 June 1989.

1. Saenger, W. *Principles of Nucleic Acid Structure* (Springer, New York, 1984).
2. Berowier, J. A., Zhang, L., Sasse-Dwight, S. & Gralla, J. D. *J. molec. Biol.* **196**, 101–111 (1987).
3. Anderson, J. E., Ptashne, M. & Harrison, S. C. *Nature* **326**, 846–852 (1987).
4. Nelson, H. C. M., Finch, J. T., Luisi, B. F. & Klug, A. *Nature* **330**, 221–226 (1987).
5. Leslie, A. G. W., Arnott, S., Chandrasekaran, R. & Ratliff, R. L. *J. molec. Biol.* **143**, 49–72 (1980).
6. Breslauer, K. J. *et al.* *Proc. Natl. Acad. Sci. U.S.A.* **84**, 8922–8926 (1987).
7. Kunkel, G. R. & Martinson, N. G. *Nucleic Acids Res.* **9**, 6869–6888 (1981).
8. Wu, H. M. & Crothers, D. M. *Nature* **308**, 509–513 (1984).
9. Sarvazyan, A. P. *Ultrasonics* **20**, 151–154 (1982).
10. Conway, B. E. *Ionic Hydration in Chemistry* (Elsevier, Amsterdam, 1981).
11. Roy-Chowdhury, P. *Macromol. chem. Phys.* **C27**(2), 219–252 (1987).
12. Buckin, V. A. *Biophys. Chem.* **29**, 283–292 (1988).
13. Shioi, H., Ogawa, T. & Yoshihashi, H. *J. Am. chem. Soc.* **77**, 4980–4982 (1955).