based on theoretical modelling incorporating various physical, meteorological and physiological effects^{6,7} backed up by extensive empirical tests⁸. Schaefer models atmospheric extinction as a combination of Rayleigh scattering from common gas molecules, absorption by stratospheric ozone and Mie scattering from aerosols. The model produces an estimate of the minimum brightness of the Moon required for 50 per cent of adults with average eyesight to be able to distinguish it against the twilight sky at the optimal time on the evening in question.

IMAGE UNAVAILABLE FOR COPYRIGHT REASONS

Christ on the cross — can we date the event?

Schaefer defines a 'visibility parameter' R as the logarithm of the ratio of the apparent brightness of the Moon to this minimum. Values of R between 0 and 1 are taken to imply that the Moon might have been observable with difficulty; values above 2 are taken to indicate that it would have been observable with ease. The standard error in R expresses uncertainties in the observing conditions. Schaefer's data can be used to generate a list of candidate dates for the crucifixion (see table). These agree completely with some earlier analyses⁹, and when taken together with historical arguments they do not challenge the role of 7 April AD 30 and 3 April AD 33 as clear favourites.

Closer analysis, however, casts severe doubt upon the significance of the lunar eclipse on 3 April AD 33. An eclipsed moon is blood red, and there is Biblical evidence of the Moon appearing like blood on the day of the crucifixion, particularly in Joel's prophecy (Joel 2:31), quoted by Peter (Acts 2:20), that the moon "would be turned to blood before that great and notable day of the Lord comes", a passage that can be interpreted as referring to an event at the time of Christ's death, before the resurrection⁹.

But the eclipse on 3 April AD 33 was only partial. When the umbra covers only a small fraction of the Moon's disk, its blood-red colour is swamped by the far brighter remainder of the Moon and is

unobservable. Furthermore, the Moon was actually already in the final stages of the eclipse when it rose in Jerusalem. The time at which the Moon left the umbra can be determined only to an accuracy of a few minutes, because of cumulative uncertainties in the relationship between the Moon's orbit about the Earth and the Earth's rotation about the Sun, but Schaefer's analysis indicates that the rising moon would first have become visible when at most one per cent of its disk was still in the umbra. Even an experienced observer would be highly unlikely to spot the umbral eclipse. Any reddish colour obvious to all observers as the Moon rose could have been due only to atmospheric filtering at low altitude, an effect that can be seen at any moonrise.

This example draws attention to the importance of theoretical models concerning the observational capabilities of the naked eye. Such studies have considerable significance for archaeoastronomy and the early history of astronomy. For example, many of the high-precision lunar horizon observations postulated by Thom to have been undertaken by Neolithic and Bronze Age Britons¹⁰ are found not to be feasible when the variable effects of refraction are taken into account; similarly, many postulated alignments of megalithic structures upon rising and setting stars¹¹ may be discounted on the basis of uncertainties in the extinction angle¹².

For the dating of the crucifixion, however, the implications are not quite so significant. Given the assumption that the Jewish sabbath has fallen unfailingly on every seventh day since Biblical times, there seems no reason, on present evidence, to cast the net wider than the two dates that have been favoured for years. But the evidence that the eclipse of 3 April AD 33 was not observable implies that the choice between these two candidates must once again be weighed equally, at least on astronomical grounds. □

Clive Ruggles is in the Departments of Archaeology and Computing Studies, University of Leicester, University Road, Leicester LE1 7RH, UK.

1. Bruce, F.F. *New Testament History* (Transl. Green, D.E.) (Black, London, 1968).
2. Ogg, G. *The Chronology of the Public Ministry of Jesus* (Cambridge University Press, 1940).
3. Link, F. *Eclipse Phenomena in Astronomy* (Springer, New York, 1969).
4. Schaefer, B.E. *Q. J. R. astr. Soc.* **31**, 53–67 (1990).
5. Schurer, E., Vermes, G. & Millar, F. *The History of the Jewish People in the Age of Jesus Christ Vol. 1* (Edinburgh University Press, 1973).
6. Bruin, F. *Vistas Astr.* **21**, 331–358 (1977).
7. Schaefer, B.E. in *Proc. I.I.I.T. Conf. on the Lunar Calendar* (ed. Ahmad, I.) **11** (1989).
8. Doggett, L.E. & Schaefer, B.E. *Sky and Telescope* **77**, 373–375 (1989).
9. Humphreys, C.J. & Waddington, W.G. *Nature* **306**, 743–746 (1983).
10. Thom, A. *Megalithic Lunar Observatories* (Oxford University Press, 1971).
11. Thom, A. *Megalithic Sites in Britain* (Oxford University Press, 1967).
12. Schaefer, B.E. *J. Hist. Astr.* **17** (Archaeoastr. Suppl. 10), S32–42 (1986).

The logic of love

TRADITIONAL courtship, says Daedalus, is a collective attempt to pair off society's men and women into the thermodynamically most stable set of couples. Partnerships form, dissolve and exchange, each set of couples a slight improvement on the previous one, except for the occasional disaster which has to be written off as experience. Sooner or later the pairing seems optimal enough to be formalized into a set of marriages. Sadly, this formalization is increasingly premature, and the global optimization continues painfully with further reshuffles in the form of divorces and remarriages.

The whole process reminds Daedalus of the mathematical trick of optimizing a big, multi-solution problem by 'iterative annealing'. Having guessed an initial solution, you carry out a sequence of small random changes governed by a fake 'temperature'. If a small change improves the solution, you accept it. If it doesn't, you accept it with a probability related to the temperature — a cunning trick for 'climbing out' of purely local optima. Keep doing this, while slowly lowering the temperature, until no further anti-optimal climbing can occur. You should then have reached, not necessarily the absolute optimum, but certainly a very good one.

So Daedalus presents Pairwise, his new matrimonial computer program. Atom-like, every client in the program is attracted to each of his or her potential partners by some assumed bond-energy, large or small. In real life (as opposed to atomic physics), there is no requirement for this attraction to be reciprocated. So, since the stability of a couple is determined by the less satisfied partner, the effective bond-energy of any 'pair-molecule' is the lesser of their two attraction-energies. The initial high temperature of the assembly rapidly dissociates the more unstable pairs for recoupling. From this torrid and unremitting promiscuity, the assembly is slowly annealed down to a glacial monogamy. The pairing that finally 'crystallizes out' is thermodynamically stable. Everyone has their best feasible choice.

DREADCO sociologists are keen to try Pairwise out on real people. They plan to ask each marriageable inhabitant of a given community to judge the attractiveness of all possible mates. From these 'bond-energies' Pairwise will find the ideal pairing, and the sociologists will wait to see how accurately it comes true. An even better test bed would be a complex knot of desperate marriages in the course of mutual re-permutation, but such convenient circumstances rarely occur outside the pages of modern novels. As a first step, DREADCO is offering a pairing service for discos, parties and singles bars, to cut out the usual covert manoeuvrings. David Jones