watch someone grabbing a ball as it was to watch a person merely mimicking this action. In the case of object-related movements, some parts of the parietal cortex — Brodmann's areas 39 and 40 — were also activated. As parts of area 40 are active during object manipulation and the parietal lobe has been implicated in action-orientated object description, it is likely that a similar 'pragmatic' analysis takes place when passively watching an object-related action.

How does the existence of this mirror system affect the execution of our own movements? How does it function when observing complicated movement sequences? Does a system like this really help us to improve our backhand or our serve? We don't have rigorous answers for these questions yet, but this should not stop the existence of a mirror system from becoming a new incentive for people like me, who are hopeless during tennis lessons.

Juan Carlos López

References and links

ORIGINAL RESEARCH PAPER Buccino, G. *et al.* Action observation activates premotor and parietal areas in a somatotopic manner: an fMRI study. *Eur. J. Neurosci.* **13**, 400–404 (2001)
FURTHER READING Iacoboni, M. *et al.* Cortical mechanisms of human imitation. *Science* **286**, 2526–2528 (1999) | Rizzolatti, G. & Arbib, M. A. Language within our grasp. *Trends Neurosci.* **21**, 188–194 (1998)
ENCYCLOPEDIA OF LIFE SCIENCES Motor system organization

markedly impaired in the trace conditioning experiments. This is precisely what we would predict if the new cells in the dentate gyrus were indeed involved in memory formation. It remains to be seen whether the behavioural deficit generalizes to other hippocampal tasks. But despite this unresolved issue, it seems that there is now some support for the old idea that some new memories rely on the birth of new cells for their formation.

Rachel Jones

References and links

ORIGINAL RESEARCH PAPER Shors, T. J. *et al.* Neurogenesis in the adult is involved in the formation of trace memories. *Nature* **410**, 372–376 (2001)
FURTHER READING Gross, C. G. Neurogenesis in the adult brain: death of a dogma. *Nature Rev. Neurosci.* **1**, 67–73 (2000) | Macklis, J. P. New memories for new neurons. *Nature* **410**, 314–317 (2001)

CHEMICAL SENSES

A taste of things to come

We have understood olfaction far better than taste ever since the olfactory receptors were cloned ten years ago. But the gustatory system has recently begun to catch up and we are witnessing the first chapters of what promises to be a fascinating story. Scott *et al.* have just published the latest episode of the saga: the characterization of a gene family that encodes candidate gustatory and olfactory receptors in *Drosophila*.

Following the lead of Clyne *et al.*, who discovered this gene family and proposed that it encoded taste receptors, Scott *et al.* identified new members of the family and showed that they are expressed on gustatory neurons that project to the suboesophageal ganglion — a gustatory centre in the fly brain. Similar to the olfactory system, each gustatory receptor is expressed in a small number of gustatory neurons and every gustatory cell seems to express a small number of receptors, perhaps only one. But a more peculiar relationship between the new genes and the previously identified *Drosophila* olfactory receptors is that some members of the new family seem to be olfactory receptors themselves. Indeed, some of the new genes are expressed on olfactory organs and project to a single olfactory glomerulus.

Do the projections from neurons that express the same gustatory receptor mimic the exquisite topographic arrangement that has been described in the olfactory system? Are specific taste modalities in *Drosophila* subserved by particular receptors? The study of Scott *et al.* has been an excellent appetizer but the main course is yet to come.

Juan Carlos López

References and links

ORIGINAL RESEARCH PAPER Scott, K. *et al.* A chemosensory gene family encoding candidate gustatory and olfactory receptors in *Drosophila*. *Cell* **104**, 661–673 (2000)
FURTHER READING Clyne, P. J. *et al.* Candidate taste receptors in *Drosophila*. *Science* **287**, 1830–1834 (2000)
ENCYCLOPEDIA OF LIFE SCIENCES Chemosensory systems | Taste

CELL FATE

Making up your mind

One of the key decisions that any neural stem cell faces is whether to commit to the neuronal or the glial lineage. The factors that regulate this commitment step are beginning to be identified, and the basic helix–loop–helix (bHLH) transcription factors have proved particularly important in this regard. These include the neurogenins (Ngn1 and Ngn2) and the mammalian achaete-scute homologue (Mash1), all of which promote neurogenesis. Indeed, Sun *et al.* recently showed that Ngn1 not only directly activates neurogenic genes, but also blocks the action of transcriptional complexes required for gliogenesis.

In the developing cortex, knocking out *Mash1* or *Ngn2* causes misspecification of individual neuronal subsets, but does not lead to an overall reduction in neurogenesis. However, *Ngn2; Mash1* double mutants show severe cortical defects, with increased and premature production of astrocytes at the expense of neurons. This implies that these two factors act in a redundant fashion to promote the neuronal fate. Now, as reported in *Neuron*, Nieto *et al.* have set out to establish the basis for this redundancy.

Although the telencephalic expression domains of *Ngn2* (dorsal) and *Mash1* (ventral) are considered to be mutually exclusive, some low-level expression of *Mash1* is seen in the cortex, and the gene is also upregulated in the *Ngn2* knockout. Nieto *et al.* found that cortical progenitors are heterogeneous with respect to *Ngn2* expression, prompting them to ask whether *Ngn2* and *Mash1* might normally act in different cell populations. To test this idea, they isolated cortical progenitors from *Mash1* or *Ngn2* knockout mice, sorted them on the basis of *Ngn2* promoter activity and assessed their ability to generate neurons. Knocking out *Ngn2* had little effect on the proportion of neurons that were generated, perhaps because of compensation by upregulation of *Mash1*. However, knocking out *Mash1* led to a reduction in neurogenesis and a conversion of progenitors to astrocytic differentiation, both of which were particularly pronounced in the non-*Ngn2*-expressing fraction.

So, the developing cortex contains a heterogeneous population of progenitor cells that differ at least in their expression of *Ngn2*. It remains to be seen whether there is also a separate *Mash1*-expressing progenitor population. This is a distinct possibility, as *Mash1* is upregulated in the *Ngn2* knockout cortex and, as Nieto *et al.* now show, its effects on neurogenesis seem to be most profound in cells that do not normally express *Ngn2*.

Heather Wood

References and links

ORIGINAL RESEARCH PAPER Nieto, M. *et al.* Neuronal bHLH genes control the neuronal versus glial fate decision in cortical progenitors. *Neuron* **29**, 401–413 (2001)
FURTHER READING Sun, Y. *et al.* Neurogenin promotes neurogenesis and inhibits glial differentiation by independent mechanisms. *Cell* **104**, 365–376 (2001)
ENCYCLOPEDIA OF LIFE SCIENCES Neural development: bHLH genes