was mutated to serine. These were recovered from conditioned medium by protein-A chromatography, cleaved with enterokinase, further purified and confirmed to be monomeric by gel-filtration chromatography as described²⁶. Appropriate binding characteristics for each protein were confirmed by surface plasmon resonance (Biacore) analysis and found to be consistent with published values¹². For CTLA-4, the final cleaved protein comprised residues 3–125 (with the C122S mutation) followed by the octa-peptide DYKDDDDK at the carboxy terminus. For B7-1, the cleaved protein consisted of residues 1–214 with the DYKDDDDK octa-peptide again forming the C terminus.

Crystallization and data collection

The B7-1/CTLA4 complex was formed by incubating a 1:1.5 molar ratio of B7-1 and CTLA4 and purified using a TSK-3000SW size exclusion column (TosoHaas). The purified complex was buffer exchanged and concentrated to 10 mg ml⁻¹ in 20 mM Tris-HCl, pH 8.0. Crystals were grown at 18 °C by the hanging drop method by combining 2 µl of protein solution with 2 µl of 14% PEG 8000, 200 mM magnesium acetate and 100 mM cacodylate, pH 6.3, and equilibrated against 1 ml of the same solution. Crystals appeared in 5 days and were soaked in 16% PEG 8000, 200 mM magnesium acetate, 20 mM Tris-HCl pH 8.0, 100 mM cacodylate, pH 6.3, 20% ethylene glycol for about 1 min before plunging into liquid nitrogen. Crystals belong to space group C222₁ with unit-cell dimensions *a* = 88.5, *b* = 183.4, *c* = 230.8 Å and a calculated solvent content of 58% for two B7-1 and two CTLA-4 molecules per asymmetric unit (adding 82,000 (82K) of glycosylation estimated from SDS-PAGE gel mobility). The 3.0 Å resolution data were collected from a single crystal at 100K at Beamline 5.0.2 at the Advanced Light Source using a Quantum 4 CCD. 725 0.2° rotation images were collected and reduced with HKL2000 (ref. 27) giving statistics outlined in Table 1.

Structure determination and refinement

The structure was solved by molecular replacement in CNS²⁸ using the crystallographic dimer observed in the human B7-1 structure as the search model¹⁰. After rigid body refinement of each B7-1 domain the *R* factor was 55.3%. Initial electron-density maps phased with B7-1 alone showed clear electron density for the two CTLA-4 molecules. The model of murine CTLA-4 monomer (PDB accession number 1dqt) was placed into electron density and was manually rebuilt in QUANTA (Molecular Simulations) to reflect the sequence of human CTLA-4. Non-conserved amino-acid residues and loop regions of CTLA-4 and B7-1, as well as sugar moieties, were entirely built into electron density. A composite annealed omit map was calculated to check the conformation of CTLA-4 and B7-1 loop regions. All refinement procedures were done in CNS and used data from 20.0–3.0 Å. A 2σ cut-off was applied during the refinement because of the anisotropy of the data. Six per cent of the reflections were randomly selected for generation of the *R*_{free} data set (see Table 1). The anisotropy of the data was corrected for also using CNS.

Tight non-crystallographic symmetry restraints were gradually relaxed during the refinement process resulting in a final model with *R*_{work} of 23.2% and *R*_{free} of 25.9%. Continuous electron density was seen for sugar moieties at four N-linked carbohydrate sites on CTLA-4 dimer (Asn 78 and Asn 110) and at ten glycosylation sites on B7-1 dimer (Asn 19, Asn 55, Asn 152, Asn 173 and Asn 192). No saccharides could be modelled at six other expected glycosylation sites on B7-1 (Asn 64, Asn 177 and Asn 198). The final model contains two B7-1 molecules (residues A1–A199 and B1–B199), two CTLA-4 molecules (residues C3–C120 and D3–D120), 20 monosaccharides and shows reasonable geometry: 98.9% of non-glycine φ and ψ angles lie in the allowed regions of the Ramachandran plot (77.6% in the most favoured regions and 21.4% in additionally allowed regions); six amino-acid residues (1.1%) have disallowed angles.

Received 27 December 2000; accepted 9 February 2001.

- Lenschow, D. J., Walunas, T. L. & Bluestone, J. A. CD28/B7 system of T cell costimulation. *Annu. Rev. Immunol.* **14**, 233–258 (1996).
- Greenfield, E. A., Nguyen, K. A. & Kuchroo, V. K. CD28/B7 costimulation: a review. *Crit. Rev. Immunol.* **18**, 389–418 (1999).
- Guinan, E. C. *et al.* Transplantation of anergic histoincompatible bone marrow allografts. *N. Engl. J. Med.* **340**, 1704–1714 (1999).
- Abrams, J. R. *et al.* Blockade of T lymphocyte costimulation with cytotoxic T lymphocyte-associated antigen 4-immunoglobulin (CTLA4Ig) reverses the cellular pathology of psoriatic plaques, including the activation of keratinocytes, dendritic cells, and endothelial cells. *J. Exp. Med.* **192**, 681–694 (2000).
- van der Merwe, P. A., Davis, S. J., Shaw, A. S. & Dustin, M. L. Cytoskeletal polarization and redistribution of cell-surface molecules during T cell antigen recognition. *Semin. Immunol.* **12**, 5–21 (2000).
- Grakoui, A. *et al.* The immunological synapse: a molecular machine controlling T cell activation. *Science* **285**, 221–227 (1999).
- Lee, K.-M. *et al.* Molecular basis of T cell inactivation by CTLA-4. *Science* **282**, 2263–2266 (1998).
- Ostrov, D. A., Shi, W., Schwartz, J.-C. D., Almo, S. C. & Nathenson, S. G. Structure of murine CTLA4 and its role in modulating T cell responsiveness. *Science* **290**, 816–819 (2000).
- Metzler, W. J. *et al.* Solution structure of human CTLA4 and delineation of a CD80/CD86 binding site conserved in CD28. *Nature Struct. Biol.* **4**, 527–531 (1997).
- Ikemizu, S. *et al.* Structure and dimerization of a soluble form of B7-1. *Immunity* **12**, 51–60 (2000).
- Collaborative computational project No. 4. The CCP4 suite: Programs for protein crystallography. *Acta Crystallogr. D* **50**, 760–776 (1994).
- van der Merwe, P. A., Bodian, D. L., Daenke, S., Linsley, P. & Davis, S. J. CD80 (B7-1) binds both CD28 and CTLA-4 with a low affinity and very fast kinetics. *J. Exp. Med.* **185**, 393–403 (1997).
- Peach, R. J. *et al.* Both extracellular immunoglobulin-like domains of CD80 contain residues critical for binding T cell surface receptors CTLA-4 and CD28. *J. Biol. Chem.* **268**, 21181–21187 (1993).
- Janin, J. & Chothia, C. The structure of protein-protein recognition sites. *J. Biol. Chem.* **265**, 16027–16030 (1990).
- Jones, E. Y., Davis, S. J., Williams, A. F., Harlos, K. & Stuart, D. I. Crystal structure at 2.8 Å resolution of a soluble form of the cell adhesion molecule CD2. *Nature* **360**, 232–239 (1992).
- Bodian, D. L., Jones, E. Y., Harlos, K., Stuart, D. I. & Davis, S. J. Crystal structure of the extracellular

- region of the human cell adhesion molecule CD2 at 2.5 Å resolution. *Structure* **15**, 755–766 (1994).
- Wang, J.-h. *et al.* Structure of a heterophilic adhesion complex between the human CD2 and CD58 (LFA-3) counterreceptors. *Cell* **7**, 791–803 (1999).
- van Raaij, M. J., Chouin, E., van der Zandt, H., Bergelson, J. M. & Cusack, S. Dimeric structure of the coxsackievirus and adenovirus receptor D1 domain at 1.7 Å resolution. *Struct. Fold Des.* **8**, 1147–1155 (2000).
- Lawrence, M. C. & Colman, P. M. Shape complementarity at protein/protein interfaces. *J. Mol. Biol.* **234**, 946–950 (1993).
- Garboczi, D. N. *et al.* Structure of the complex between human T-cell receptor, viral peptide and HLA-A2. *Nature* **384**, 134–141 (1996).
- Boyington, J. C., Motyka, S. A., Schuck, P., Brooks, A. G. & Sun, P. D. Crystal structure of an NK cell immunoglobulin-like receptor in complex with its class I MHC ligand. *Nature* **405**, 537–543 (2000).
- Shapiro, L. *et al.* Structural basis of cell–cell adhesion by cadherins. *Nature* **374**, 327–337 (1995).
- Davis, S. J., Ikemizu, S., Wild, M. K. & van der Merwe, P. A. CD2 and the nature of protein interactions mediating cell-cell recognition. *Immunol. Rev.* **163**, 217–236 (1998).
- Griffin, M. D. *et al.* Blockade of T cell activation using a surface-linked single-chain antibody to CTLA-4 (CD152). *J. Immunol.* **164**, 4433–4442 (2000).
- Bachmann, M. F., Kohler, G., Ecabert, B., Mak, T. W. & Kopf, M. Lymphoproliferative disease in the absence of CTLA-4 is not T cell autonomous. *J. Immunol.* **163**, 1128–1131 (1999).
- Somers, W. S., Tang, J., Shaw, G. D. & Camphausen, R. T. Insights into the molecular basis of leukocyte tethering and rolling revealed by structures of P- and E-selectin bound to sLex and PSGL-1. *Cell* **103**, 467–479 (2000).
- Otwinowski, Z. & Minor, W. Processing of X-ray diffraction data collected in oscillation mode. *Methods Enzymol.* **276**, 307–326 (1997).
- Brunger, A. T. *et al.* Crystallography and NMR system (CNS): a new software suite for macromolecular structure determination. *Acta Crystallogr. D* **54**, 905–921 (1998).
- Carson, M. Ribbons 2.0. *J. Appl. Crystallogr.* **24**, 958–961 (1991).
- Esnouf, R. M. An extensively modified version of Molscript that includes enhanced colouring capabilities. *J. Mol. Graph.* **15**, 132–134 (1997).

Acknowledgements

We thank R. Zollner and his colleagues for large scale cell culture, and D. I. Stuart and E. Y. Jones for helpful comments. We also thank the staff at Advanced Light Source for assistance with data collection. S.I. and S.J.D. are supported by the Wellcome Trust and the Human Frontier Science Programme.

Correspondence and requests for materials should be addressed to W.S.S. (e-mail: wsomers@genetics.com) or L.M. (e-mail: lmosyak@genetics.com). The coordinates have been deposited in the Protein Data Bank. The accession code is 1I84.

corrections

Genetic control and evolution of sexually dimorphic characters

Artyom Kopp, Ian Duncan, Dorothea Godt & Sean B. Carroll

Nature **408**, 553–559 (2000).

The name of one author, Dorothea Godt, was omitted from the author list but is included above. She is affiliated with the Department of Zoology, University of Toronto, Canada. In addition, a gene name was misspelled: the correct spelling is *bric à brac*. □

Mitochondrial genome variation and the origin of modern humans

Max Ingman, Henrik Kaessmann, Svante Pääbo & Ulf Gyllenstein

Nature **408**, 708–712 (2000).

The complete mtDNA sequences upon which the analyses are based can be found at our website (<http://www.genpat.uu.se/mtDB/>), dedicated to the analysis of complete mtDNA genomes of humans. The 53 sequences are also available in Genbank with the accession numbers AF346963–AF347015. □

Genetic control and evolution of sexually dimorphic characters in *Drosophila*

Artyom Kopp^{*†}, Ian Duncan[†] & Sean B. Carroll^{*}

^{*} Howard Hughes Medical Institute and Laboratory of Molecular Biology, University of Wisconsin-Madison, 1525 Linden Drive, Madison, Wisconsin 53706-1596, USA

[†] Department of Biology, Washington University, St Louis, Missouri 63130, USA

Sexually dimorphic abdominal pigmentation and segment morphology evolved recently in the *melanogaster* species group of the fruitfly *Drosophila*. Here we show that these traits are controlled by the *bric-a-brac* (*bab*) gene, which integrates regulatory inputs from the homeotic and sex-determination pathways. *bab* expression is modulated segment- and sex-specifically in sexually dimorphic species, but is uniform in sexually monomorphic species. We suggest that *bab* has an ancestral homeotic function, and that regulatory changes at the *bab* locus played a key role in the evolution of sexual dimorphism. Pigmentation patterns specified by *bab* affect mating preferences, suggesting that sexual selection has contributed to the evolution of *bab* regulation.

A key challenge in evolutionary biology is to identify genetic events responsible for morphological change, and to understand how changes at the molecular level affect development and translate into phenotypic diversity. To achieve this, two distinct approaches have been pursued in recent years. First, comparative studies have revealed strong correlations between the expression patterns of individual regulatory genes during development and differences in morphology^{1–7}. Second, direct genetic analysis has been used to estimate the number and identity of genetic loci that contribute to morphological variation within and between species^{8–10}. Despite their respective successes, the two approaches remain far apart because of their different scales of analysis. Comparative studies have concentrated mainly on slowly evolving traits among high-level taxa, but genetic analyses are only possible among closely related species that produce viable and fertile hybrids.

Our approach to bridging this gap between evolutionary genetics and comparative embryology is to analyse and compare the development of rapidly evolving morphological traits. In many animals, secondary sexual characteristics evolve rapidly^{11,12}, making them good candidates for analysis. One such character in *Drosophila* is the pigmentation of adult abdominal segments. In *D. melanogaster*, abdominal pigmentation is sexually dimorphic. Segments 1 to 6 in females and 1 to 4 in males carry only a posterior stripe of dark pigment. However, segments 5 and 6 (A5 and A6) in males are completely pigmented (Fig. 1a and b), giving the species its name. This pattern is of recent evolutionary origin; in most *Drosophila* species, male-specific pigmentation is absent, so that females and males are pigmented identically (Fig. 1c and d). To understand how this new pattern originated and evolved, we have characterized the regulatory circuit that controls its development, and compared its operation in sexually dimorphic and monomorphic species.

Genetic control of male-specific pigmentation

The non-sex-specific striped pigmentation, which is present in most *Drosophila* species, is controlled by the transcription factor *optomotor-blind* (*omb*)¹³. In *omb* mutants, all pigment is lost from segments A1–A4 in males (Fig. 2a), and from all abdominal segments in females (not shown). However, we find that *omb*[–] males have fully pigmented A6, and mostly pigmented A5 (Fig. 2a). Thus, the ancestral striped pigmentation and the newly evolved male-specific pigmentation are controlled by separate genetic pathways.

The dark pigmentation of A5 and A6 in males is controlled by the homeotic gene *Abdominal-B* (*Abd-B*)^{14,15}. In the pupal abdomen, *Abd-B* is expressed at progressively higher levels in A5, A6 and A7,

but is absent from the more anterior segments (Fig. 3a). Ectopic expression of *Abd-B* in A3 and A4 expands male-specific pigmentation to these segments^{14–17} (Fig. 2b). Conversely, loss of *Abd-B* expression from A5–A7 (ref. 18) eliminates male-specific pigmentation, but has no effect on the non-sex-specific pigment stripes (Fig. 2c). *Abd-B* is the primary activator of male pigmentation, but another homeotic gene, *abdominal-A* (*abd-A*), is expressed in A2–A7 (Fig. 3b) and contributes to the specification of A5 and A6 identities^{14–16}.

The development of sexually dimorphic external characteristics is controlled by the *doublesex* (*dsx*) gene^{19,20}. Alternative splicing of the *dsx* transcript produces a male-specific product in males (*dsxM*), and a female-specific product in females (*dsxF*)^{21,22}. Loss of *dsx* function in females results in the development of male-like pigmentation (Fig. 2d), which can be suppressed by heat-shock *dsxF* transgenes^{23,24}. Male-specific pigmentation is therefore expressed by default, and must be actively repressed by *dsxF*.

Thus, the development of sexually dimorphic pigmentation requires integration of homeotic and sex determination gene inputs. In investigating how this integration is achieved, we discovered a newly evolved genetic circuit that appears to be responsible for the origin of male-specific pigmentation.

bric-a-brac represses male-specific pigmentation

A gene near the left tip of the third chromosome contributes to the

Figure 1 Sexually dimorphic pigmentation is of recent evolutionary origin. *D. melanogaster* female (a) and male (b); and *D. willistoni* female (c) and male (d). Most abdominal segments contain pigmented dorsal and unpigmented ventral cuticular plates (tergite (Ter) and sternite (St), respectively) (see Fig. 6a and b for more detail). In the ancestral condition, pigmentation is similar in males and females (c, d). In the derived condition, abdominal segments 5 and 6 (A5 and A6) are fully pigmented in the male (b), but not in the female (a).

variation in female abdominal pigmentation²⁵. In investigating this genetic region, we found that loss of one copy of the *bric-a-brac* (*bab*) locus results in the development of male-specific pigmentation in females (Fig. 2e), but has no effect on the male abdomen. Ectopic pigmentation in heterozygous *bab* females is suppressed by reducing the dosage of *Abd-B* (Fig. 2f), but is not eliminated by loss of *omb* (not shown). This suggests that *bab*⁺ represses the development of male-specific pigmentation in females by opposing the function of *Abd-B*. The *bab* locus contains two closely related genes, *bab1* and *bab2*, which encode putative transcription factors with multiple roles in development^{26,27}. Ectopic pigmentation in females increases in the order *bab1/+* < *bab1/bab1* < *bab1bab2/+* < *bab1bab2/bab1* (not shown), indicating that both genes are involved

Figure 2 Control of male-specific pigmentation by *Abd-B*, *dsx* and *bab*. **a**, Male hemizygous for the *omb* null mutant *l(1)omb²⁸²* lacks segmental pigment stripes but retains pigmentation in A5 and A6. **b**, In *Abd-B^{Mcp} Abd-B^{Sab}* homozygous males, male-specific pigmentation is expanded into A3 and A4. **c**, In *Df(3R)RS4-8/Df(3R)RS1-98* males, loss of *Abd-B* function from A5–A7 eliminates male-specific pigmentation, but has no effect on the non-sex-specific striped pigmentation (see also Fig. 6c). **d**, *dsx¹/Df(3R)dsx2D* chromosomal females show male-like pigmentation of A5 and A6, with the exception of a small anterior-lateral margin of A5 (arrow). **e**, *bab^{Ar07/+}* female shows male-specific pigmentation of A5 and A6. **f**, *bab* phenotype is suppressed by reduction of *Abd-B* dosage in *bab^{Ar07}/Df(3R)RS4-8* females. **g**, Low-level *bab* expression in *UAS-bab2⁴⁻⁶⁶* in the absence of a GAL4 driver results in the loss of male-specific pigmentation, but has no effect on the non-sex-specific pigmentation. **h**, High-level *bab* expression in *pnr-GAL4/UAS-bab2⁴⁻⁵¹* males results in the loss of both male-specific and striped pigmentation (arrows); the latter effect is seen in females as well. An identical phenotype is caused by misexpression of *bab1*. The loss of male-specific pigmentation in this and other *GAL4/UAS-bab* combinations is enhanced by *abd-A Abd-B* deficiencies, and partly suppressed by *abd-A Abd-B* duplications (not shown). **i**, *bab^{Ar07}/bab^{Ar07}* female. Ectopic pigmentation is seen in A2–A7 tergites, A4–A7 sternites, and occasionally in the A1 tergite. An identical pigmentation phenotype is seen in males (not shown).

Figure 3 Regulation of *bab* expression by *Abd-B* and *dsx*. All panels show gene expression in the dorsal abdominal epidermis at 40–45 hours after pupariation, shortly before the onset of cuticle differentiation. **a**, *Abd-B-lacZ* female. Segment boundaries are indicated by arrows; *Abd-B* expression is modulated on a parasegmental rather than segmental basis. In A5, expression is only visible in the polyloid bristle cells. **b**, *Abd-A* is expressed in A2–A7. **c**, **d**, *bab1-lacZ* expression in wild-type female and male, respectively. *bab* expression in A5 and A6 is absent in males (**d**) and downregulated in females (**c**). *bab* expression is first seen at 35 hours after pupariation, shortly before the formation of adult epidermis is completed. *bab* expression is modulated parasegmentally, and is not affected by *hh* or *omb* mutations (not shown). *bab* is also expressed in the ventral abdomen, but at a lower level than in the tergites. Little or no expression is seen in A1. **e**, In *Abd-B^{Mcp} Abd-B^{Sab}/+* males, *bab1-lacZ* is repressed in A3 and A4. **f**, *Bab2* expression is derepressed in A5–A7 in a *Df(3R)RS4-8/Df(3R)RS1-98* male. **g**, *bab1-lacZ* expression in *dsx¹/dsx¹⁸* chromosomal females is similar to that in males (**d**). **h**, Control of sexually dimorphic pigmentation of A5 and A6 by *Abd-B*, *abd-A*, *dsx* and *bab* (see text). It is unclear whether the same circuit operates in A7, as *bab* expression in females is stronger in A7 than in A6. *bab* expression is also present throughout A8–A10, where it is apparently independent of *Abd-B* and *dsx*.

in repressing male pigmentation. For simplicity, we treat the entire locus as one gene, *bab*, unless noted otherwise.

The expression pattern of *bab* at the pupal stage when the adult epidermis develops reflects its sex- and segment-specific function. In females, *bab* expression is strongest in segments A2 and A3, and progressively weaker in A4, A5 and A6 (Fig. 3c). In males, *bab* expression is considerably weaker than in females in all segments. Most strikingly, it is completely absent from A5 and A6 (Fig. 3d). This pattern of *bab* repression correlates with the presence of sex-specific pigmentation in males, and its absence in females.

To test whether *bab*⁺ is sufficient to repress pigmentation, we ectopically expressed the *bab* genes in the pupal abdomen. Low-level expression of *bab*⁺ results in the loss of male-specific pigmentation, but has no other effects on external morphology (Fig. 2g), indicating that differential regulation of *bab* plays a central role in establishing sexual dimorphism. *bab*⁺ can also repress non-sex-specific pigment stripes when expressed at a higher level (Fig. 2h). This suggests that *bab*⁺ acts as a general repressor of pigmentation, but that its effects are overridden by *omb* in the posterior part of each segment. Consistent with this, complete loss of both *bab* genes results in ectopic pigmentation of A2 to A7 in both sexes (Fig. 2i). This phenotype is not caused by expansion of *Abd-B* expression, which appears normal in these mutants (not shown). In *bab* homozygotes, the intensity of pigmentation is higher in the more posterior segments than in those more anterior (Fig. 2i). This suggests that pigmentation does not develop by default in the absence of *bab*, but is actively promoted by *Abd-B* and *abd-A*.

bab integrates regulatory inputs from *Abd-B* and *dsx*

The sexually dimorphic repression of *bab* in the posterior abdomen suggests that *bab* integrates the homeotic and sex determination regulatory inputs. To test this, we examined *bab* expression in *Abd-B* and *dsx* mutant backgrounds. We find that ectopic expression of *Abd-B* in A3 and A4 eliminates *bab* expression from these segments in males (Fig. 3e), and downregulates it in females (not shown). Conversely, *bab* is derepressed in A5–A7 in the mutants that lack *Abd-B* function in these segments (Fig. 3f). Together, these results indicate that *bab* expression in A5 and A6 is normally repressed by *Abd-B*. The slight downregulation of *bab* in A4 (Fig. 3c and d) suggests that it is also weakly repressed by *abd-A*.

In *dsx*[−] intersexes, *bab* is expressed in a male-like pattern (Fig. 3g), suggesting that *dsxF* upregulates *bab* transcription in females. *Abd-B* and *abd-A* expression is identical in males, females and *dsx*[−] intersexes (not shown), indicating that *bab* is regulated independently by homeotic and sex-determination inputs. *dsx*^{Dominant} intersexes, which express both male- and female-specific *dsx* products²⁸, also show male-like expression of *bab* (not shown), indicating that *dsxM* can interfere with *dsxF* function. The two *dsx* isoforms encode transcription factors that bind the same DNA sequence, but have opposite effects on gene expression^{29,30}. *dsx*[−] intersexes differ from males in having a small unpigmented region at the anterior-lateral margin of A5 (refs 19, 23; Fig. 2d), suggesting that *dsxM* may have a slight negative influence on *bab* expression.

Our results suggest that *bab*⁺ regulates sexually dimorphic pigmentation by integrating regulatory inputs from the homeotic genes and the sex determination pathway (Fig. 3h). In this regulatory circuit, *bab*⁺ acts as a general repressor of pigmentation, and *Abd-B* and *abd-A* promote pigmentation in both sexes. In addition, *Abd-B*, and to a lesser extent *abd-A*, repress *bab* transcription. In males, this results in the absence of *bab* from A5 and A6, allowing *Abd-B* and *abd-A* to promote pigmentation in these segments. However, in females, *dsxF* prevents *bab* transcription from being completely repressed by the homeotic genes. As a result, *bab* is present in A5 and A6 in females, where it blocks the ability of *Abd-B* and *abd-A* to promote pigmentation. In A2–A4, *abd-A* alone is not sufficient either to repress *bab* or to overcome its inhibitory effect on pigmentation; thus, only the *omb*-dependent striped pigmentation

is generated. Because *Abd-B*, *abd-A* and *dsx* encode transcription factors, they may regulate *bab* expression directly.

Evolution of sexually dimorphic pigmentation

The central role of *bab* as an integrator of homeotic and sex-determination gene inputs suggests that changes in *bab* regulation may have been responsible for the evolution of sexually dimorphic pigmentation. In the subgenus *Sophophora*, male-specific pigmentation is present only in the *melanogaster* species group. Within this group, sexual dimorphism is seen in all species of the *melanogaster* subgroup and the closely related oriental subgroups, whereas the *ananassae* and *montium* subgroups contain both sexually dimorphic and sexually monomorphic species.

We find that in species with male-specific pigmentation of A5 and A6, *bab* expression is absent or strongly downregulated in these segments in males, but not in females (Fig. 4a and b; Fig. 5). Moreover, in the sexually monomorphic species outside the *melanogaster* species group, *bab* expression is identical in both sexes and in all segments from A2 to A7 (Fig. 4c and d; Fig. 5). This correlation suggests that changes in the regulation of *bab* by *Abd-B* and *dsx* played an important role in the origin of sexually dimorphic pigmentation.

Surprisingly, in some species of the *montium* subgroup, such as *D. kikkawai*, *bab* expression in A5 and A6 is downregulated in males despite the absence of sex-specific pigmentation (Fig. 5). This suggests that the function of *bab*⁺ may not be limited to the control of pigmentation. To test whether *bab*⁺ regulates other morphological characters in abdominal segments, we analysed the phenotypes produced by gain and loss of *bab* function in *D. melanogaster*.

Homeotic function of *bab*

In *D. melanogaster*, A5, A6 and A7 differ from the more anterior segments not only in pigmentation, but also in the size and shape of tergites and sternites and in the distribution of bristles and trichomes (Fig. 6a and b). These characteristics, which are especially pronounced in the male, are under the control of *Abd-B*, and the differences among segments are thought to be specified by the different levels of *Abd-B* expression^{14–18}. Ectopic expression of *Abd-B* induces posterior characters in anterior segments, while the loss of *Abd-B* function from A5–A7 transforms these segments to a more anterior identity (Fig. 6c).

We find that *bab*⁺ regulates segment shape and bristle and trichome patterns in a manner reciprocal to *Abd-B*. Loss of *bab*⁺ function in females enhances posterior characteristics in A6, A7 and A8 (Fig. 6d). No phenotype is seen in males, consistent with the absence of *bab* expression in posterior segments. Conversely, ectopic expression of *bab* transforms A6 and A7 to a more anterior identity in both males (Fig. 6e) and females (see Supplementary Information). These observations suggest that *bab*⁺ acts as an antagonist of

Figure 4 Repression of *bab* correlates with the presence of male-specific pigmentation. *Bab2* expression is modulated in the sexually dimorphic species *D. biarmipes* (a, female; b, male), but not in the sexually monomorphic *D. willistoni* (c, female; d, male).

Abd-B homeotic function, and that posterior abdominal characters are determined by the balance between *Abd-B* and *bab* activities.

This model predicts that evolutionary changes in *bab* regulation should result in morphological transformation of *Abd-B*-expressing segments. Indeed, we find that the entire suite of characteristics that distinguishes A5 and A6 from the more anterior segments in *D. melanogaster* is of recent evolutionary origin. In *D. willistoni*, *bab* is expressed strongly in A5 and A6 in males (Fig. 4d), whereas *Abd-B* is expressed in the same pattern as in *D. melanogaster* (not shown). As predicted, A5 and A6 are almost identical to the more anterior, non-*Abd-B*-expressing segments in the males of this species (Fig. 6f). In contrast, the *melanogaster* species group shows great diversity of bristle and trichome patterns in posterior abdominal segments (Table 1). The two main lineages within this group show different patterns of evolution. In the clade composed of the *melanogaster* and oriental subgroups, male-specific pigmentation and bristle and trichome patterns have evolved in a concerted fashion (Table 1, group A). However, in the *ananassae* + *montium*

lineage, these characteristics vary independently of each other, and sexually dimorphic bristle and trichome patterns are sometimes observed in species that do not show visible modulation of *bab* expression (Table 1, group B, compare with Fig. 5). This suggests that evolutionary changes have occurred not only in *bab* regulation, but also in the target genes of *bab* and in other genes regulated by *Abd-B* and *dsx*. We also note that suppression of A7 development in males has occurred earlier in evolution than visible modulation of *bab* expression, despite the ability of *bab* to override this suppression (Fig. 6e).

Sexually dimorphic pigmentation affects mating preferences

The rapid evolution of sexually dimorphic pigmentation and segment morphology may have been driven by sexual selection. We therefore tested whether male-specific pigmentation confers a competitive advantage in *D. melanogaster* males. Surprisingly, we find that UAS-*bab2*⁴⁻⁶⁶ males, which lack male-specific pigmentation but are otherwise normal (Fig. 2g), enjoy the same mating

Figure 5 Modulation of *bab* expression correlates with sexual dimorphism. Phylogenetic hypothesis is based on information from various sources⁴³⁻⁴⁶. The divergence time between *melanogaster* and *obscura* species groups has been estimated at 25 million years⁴⁷. A6 pigmentation is variable in the females of *D. kikkawai* and *D. rufa* (asterisk), and in both females and males of *D. auraria* and *D. jambulina* (double asterisk). Bab2 is repressed in A5 and A6 in the males of sexually dimorphic species (red), but not in the sexually monomorphic species (blue). Exceptions to this rule are found among species of

the *montium* subgroup, in which Bab2 is downregulated in the absence of male-specific pigmentation (yellow). Expression was only analysed in the dorsal abdomen. We cannot rule out that species listed as having unmodulated *bab* expression do in fact have subtle modulation that is below the resolution of our methods. For instance, a twofold reduction in the dosage of *bab*, which causes a dramatic pigmentation phenotype in the females of *D. melanogaster* (Fig. 2e), is not detectable by antibody staining. For the same technical reason, quantitative differences between sexes and species were not studied.

success as wild-type males (27 versus 32 matings; $\chi^2 = 0.42$; $P > 0.05$). Thus, although male pigmentation may have been important in the past, it appears to have little or no effect on female mating preferences in extant *D. melanogaster*.

However, we find that *D. melanogaster* males discriminate strongly against heterozygous *bab* females, which have ectopic male-specific pigmentation but are otherwise normal (Fig. 2e), compared with females with lightly pigmented A5 and A6 (23 versus 105 matings; $\chi^2 = 52.53$; $P < 0.001$). Importantly, white mutant males, which are visually impaired, mate equally with *bab*/+ and lightly pigmented females (34 versus 39 matings; $\chi^2 = 0.34$; $P > 0.05$), suggesting that discrimination against *bab* heterozygous females is due to their pigmentation. These results suggest that female pigmentation is important in determining their attractiveness to males, and that the absence of male-specific pigmentation in females may be maintained by sexual selection.

A model of evolution of a genetic circuit under sexual selection

Our findings indicate that changes in *bab* regulation have played an important part in the evolution of abdominal segment morphology (Fig. 6g). The presence of *bab* expression in all *Drosophila* species examined suggests that its roles in antagonizing the homeotic function of *Abd-B* and repressing pigmentation are ancestral.

However, in the ancestral condition, *bab* expression was independent of *Abd-B* and *dsx*, resulting in sexually monomorphic pigmentation and segment morphology. In the *melanogaster* species group, *bab* evolved to be under the control of *Abd-B* and *dsx*. This eliminated *bab* from *Abd-B*-expressing segments in the male and resulted in a major transformation of male segment morphology. Subsequent diversification of pigmentation, bristle and trichome patterns was probably driven both by the fine-tuning of *bab* regulation and by changes in the downstream targets of *bab* and *Abd-B*.

Two features of this genetic circuit make it highly plastic and evolvable. First, the adult phenotype is sensitive to quantitative changes in *bab* expression. Second, the level of *bab* expression is determined by the balance between *Abd-B* and *dsxF* inputs. If *bab* is regulated directly by *Abd-B* and *dsx*, then the evolution of sexually dimorphic pigmentation and segment morphology may ultimately be traced to the acquisition and modification of binding sites for the *Abd-B* and *Dsx* proteins in the *cis*-regulatory region of *bab*. Thus, even a subtle molecular change could be expressed phenotypically and become subject to selection.

This evolutionary model is further supported by the presence of intraspecific genetic variation in sexually dimorphic pigmentation in many extant species^{31,32}. In at least one case, there is strong

Figure 6 The homeotic function of *bab*. Photomicrographs of the cuticles are available in the Supplementary Information. **a**, Wild-type male of *D. melanogaster*. A7 is greatly reduced and lacks both tergite and sternite. In addition, the A6 sternite lacks bristles and has a unique horseshoe-like shape, and most of the A6 tergite is devoid of the small hairs (trichomes) that cover the more anterior segments. Male genitalia develop from A9, whereas A8 development is repressed. **b**, In a wild-type female, A7 sternite and A7 and A6 tergites are only partly covered by trichomes; the A7 sternite also has a distinctive shape. Female genitalia develop from A8, whereas A9 development is repressed. **c**, In *Df(3R)RS4–8/Df(3R)RS1–98* males, A5–A7 are transformed towards A4 identity¹⁸ (indicated in red). The presence of sternite and tergite in A7 is restored, A6 and A7 tergites are completely covered by trichomes, and A6 and A7 sternites are rectangular and carry bristles. **d**, In *bab^{Ar07}/bab^{Ar07}* females, expression of posterior morphological characters is increased (blue). Trichomes are absent from the A7 tergite and sternite, and the number of bristles on A6 and A7 sternites is reduced. The A6 sternite also acquires a shape reminiscent of the A6 in the male, or A7 in the female. A weaker phenotype is seen in *bab^{Ar07}/+* (not shown). The thorn bristles normally present on the vaginal plates of A8 (**b**) are replaced by long, wavy bristles typical of the male genital arch (A9). *bab⁻* females are nevertheless distinguishable from males in the morphology of A6 and particularly A7,

indicating that *dsxM* and/or *dsxF* regulate additional target genes in the abdomen. **e**, Ubiquitous *bab* expression in *C765-GAL4/UAS-bab^{24–51}* males transforms A6 and A7 to a more anterior identity (red). The presence of tergite and sternite in A7 is restored, A6 sternite acquires a rectangular shape and a number of bristles, and the entire A6 tergite is covered by trichomes. A similar phenotype is caused by misexpression of *bab1* (not shown). The phenotype is enhanced by *abd-A Abd-B* deficiencies, and partly suppressed by *abd-A Abd-B* duplications (not shown). Ectopic *bab* has no effect in the genitalia and analia of either sex (segments A8–A10). The homeotic function of *bab* is apparently limited to the adult stage, because ectopic *bab* expression has no effect in the larval cuticle (not shown). **f**, In *D. willistoni* males, the A6 sternite is rectangular and carries numerous bristles, and the A6 tergite is covered by dense trichomes. We note the similarity to **c** and **e**. **g**, The role of *bab* in the evolution of sexually dimorphic segment morphology. We suggest that sexual dimorphism evolved through the acquisition of two new regulatory interactions, shown in red. The number of genetic steps involved in this transition is unclear. In the *ananassae* and *montium* subgroups, *bab* regulation may differ strongly between closely related species (Fig. 5), indicating that the new genetic circuit retained considerable plasticity.

Table 1 Male-specific characters in two lineages of *melanogaster* species group*

Species	A5/A6 pigmentation (A6 only in <i>montium</i> subgroup)	Loss of A6 tergite trichomes	Loss of A6 sternite bristles
A			
<i>D. melanogaster</i>	+	+	+
<i>D. biarmipes</i>	+	+	+
<i>D. takahashii</i>	+	+	+
<i>D. eugracilis</i>	+	+	+
<i>D. ficusphila</i>	+	+	+
<i>D. elegans</i>	+	—	+
<i>D. fuyamai</i>	+	+	+
B			
<i>D. ananassae</i>	—	—	+
<i>D. atripex</i>	+	—	+
<i>D. leontia</i>	—	—	+
<i>D. bocki</i>	—	+	+
<i>D. auraria</i>	+/-†	+	—
<i>D. rufa</i>	+	+	—
<i>D. burlai</i>	+	+	+
<i>D. nikananu</i>	—	+	+

* Photomicrographs of some species are available in the Supplementary Information.

† A6 pigmentation is variable in *D. auraria* males.

evidence that allelic differences at the *bab* locus contribute to this variation. Females found in natural populations of *D. melanogaster* vary widely in the extent of A6 pigmentation, ranging from near-zero to 100% (ref. 25). The locus with the largest effect on this variation has been mapped²⁵—to the exact position of *bab*. These observations suggest that sexually dimorphic pigmentation evolved through fixation of intraspecific genetic variants at the *bab* locus.

Fixation of new *bab* alleles was probably driven initially by “runaway” sexual selection. In this case, a slight female preference for a weakly pronounced male character would initiate a positive feedback loop that would rapidly increase both the expression of the male character and the female preference for it³³. This self-reinforcing mechanism can drive rapid character divergence and create new species through sexual isolation^{34–37}. Male-specific pigmentation could evolve by this mechanism, with increasingly discriminating females selecting for increasingly dark males. However, once fixed, sexual characteristics can lose their significance as they are overtaken by newly evolving signals and as females become habituated and “resistant” to old characters³⁸. This may explain our finding that male pigmentation has no effect on mating success in extant *D. melanogaster*.

Whereas the runaway model explains the evolution of male sexual characters, it does not account for the absence of these characters in females, that is, sexual dimorphism. However, sexual dimorphism can be produced effectively by counter-selection against male-specific traits in females³³. Consistent with this, we find that *D. melanogaster* males discriminate against females that have male-like pigmentation. In most *Drosophila* species, including *D. melanogaster*, males seek out females at feeding sites and attempt to court as many as possible. Courting other males is not only disadvantageous in competition for females, but may also carry a direct cost³⁹. Thus, males are probably selected for an ability to avoid courting other males, and pigmentation may be used to identify females at a distance.

The evolution of *bab* regulation offers a tractable model of how selection creates new morphological characters through changes in DNA sequence. Analysis of the *cis*-regulatory elements of *bab* in sexually dimorphic and monomorphic species will help to clarify the molecular basis of morphological divergence between these taxa. □

Methods

Anterior is upward in all figures. Adult abdomens were cut along the dorsal midline and mounted flat so that ventral cuticle is on the left and dorsal cuticle is on the right. Pupal dissections, X-Gal staining⁴⁰ and antibody staining⁴ were performed as described.

Drosophila strains and reagents

Abd-B^{Mcp} Abd-B^{Sab} is a double gain-of-function mutation that results in ectopic *Abd-B* expression in A3 and A4 (ref. 15). *Df(3R)RS4–8* removes the *cis*-regulatory elements responsible for *Abd-B* expression in A5–A7; in *Df(3R)RS4–8/Df(3R)RS1–98* heterozygotes, these segments lack *Abd-B* function¹⁸. *Abd-B-lacZ* enhancer trap HCJ199 (ref. 41) reproduces the pattern of *Abd-B* protein expression (not shown). *Abd-A* protein expression was detected by monoclonal antibody Dmabd-A.1 (ref. 42).

bab^{Ar07} (F. Laski, unpublished work) is a small deletion that inactivates both *bab* genes. *UAS-bab^{24–66}* (D. Godt, unpublished work) is inserted into the 5' regulatory region of the ecdyson-inducible gene *ftz-F1* (data not shown). The phenotype produced by this line in the absence of a GAL4 driver is presumably caused by low-level expression of *bab2* driven by *ftz-F1* regulatory elements during metamorphosis. In the presence of GAL4, this line generates phenotypes similar to other *UAS-bab* insertions, including *UAS-bab^{4–51}*. *pnr-GAL4* drives gene expression in a stripe along the dorsal midline in all abdominal segments. *bab1-lacZ* enhancer trap²⁶ reproduces the expression of both *bab* genes. *Bab1* and *Bab2* proteins, detected by specific antibodies (F. Laski, unpublished work), are expressed in similar patterns in all species examined (*melanogaster*, *virilis*, *willistoni*, *ananassae* and *eugracilis*) (data not shown).

Mating choice experiments

Virgin males and females, 3–5 days old, were placed together in a 95 × 24mm glass vial, and matings were observed visually. For male-choice experiments, a single male was combined with two females of different genotypes. For female-choice experiments, a single female was combined with two males of different genotypes. Only matings that occurred in the first 30 min were counted; the number of unmated groups was also recorded. The *bab* genotypes used were *bab^{Ar07}/+*, *bab^{Ar07}/TM3*, *bab^{FC}/bab-lacZ* and *Df(3L)Ar12–1/+*. Strains in which females have lightly pigmented A6 were isolated from various laboratory stocks, and were not standard wild-type strains. For some experiments, *bab^{Ar07}* was introgressed into a ‘light’ strain for 3–4 generations. Oregon-R strain was used as a source of wild-type males. For the most part, identical results were obtained with different *bab* alleles and wild-type strains, as well as following the introgression of *bab^{Ar07}* into a lightly pigmented strain, suggesting that mating discrimination is not due to differences in genetic backgrounds.

In experiments with one ‘light’ strain, wild-type males mated equally with *bab^{Ar07}/+* and lightly pigmented females (18 versus 22 matings; $\chi^2 = 0.4$; $P > 0.05$). In these experiments, males courted both types of females repeatedly; however, courtship was quickly aborted in most cases, and nearly 50% of males did not mate within 1 hour. The reasons for this behaviour are unclear.

Received 30 June; accepted 10 October 2000.

- Warren, R., Nagy, L., Selegue, J., Gates, J. & Carroll, S. Evolution of homeotic gene regulation and function in flies and butterflies. *Nature* **372**, 458–461 (1994).
- Burke, A. C., Nelson, C. E., Morgan, B. A. & Tabin, C. *Hox* genes and the evolution of vertebrate axial morphology. *Development* **121**, 333–346 (1995).
- Averof, M. & Akam, M. *Hox* genes and the diversification of insect-crustacean body plans. *Nature* **376**, 420–423 (1995).
- Brakefield, P. M. *et al.* Development, plasticity and evolution of butterfly eyespot patterns. *Nature* **384**, 236–242 (1996).
- Averof, M. & Patel, N. H. Crustacean appendage evolution associated with changes in *Hox* gene expression. *Nature* **388**, 682–686 (1997).
- Stern, D. L. A role of *Ultrabithorax* in morphological differences between *Drosophila* species. *Nature* **396**, 463–466 (1998).
- Sucena, E. & Stern, D. L. Divergence of larval morphology between *Drosophila sechellia* and its sibling species caused by *cis*-regulatory evolution of *ovo/shaven-baby*. *Proc. Natl Acad. Sci. USA* **97**, 4530–4534 (2000).

8. True, J. R., Stam, L. F., Zeng, Z. -B. & Laurie, C. C. Quantitative genetic analysis of divergence in male secondary sexual traits between *Drosophila simulans* and *Drosophila mauritiana*. *Evolution* **51**, 816–832 (1997).
9. Gurganus, M. C., Nuzhdin, S. V., Leips, J. W. & MacKay, T. F. High-resolution mapping of quantitative trait loci for sternopleural bristle number in *Drosophila melanogaster*. *Genetics* **152**, 1535–1604 (1999).
10. Lyman, R. F., Lai, C. & MacKay, T. F. Linkage disequilibrium mapping of molecular polymorphisms at the *scabrous* locus associated with naturally occurring variation in bristle number in *Drosophila melanogaster*. *Genet. Res.* **74**, 303–311 (1999).
11. Anderson, M. *Sexual Selection* (Princeton Univ. Press, Princeton, 1994).
12. Eberhard, W. G. Copulatory courtship and cryptic female choice in insects. *Biol. Rev.* **66**, 1–31 (1991).
13. Kopp, A. & Duncan, I. Control of cell fate and polarity in the adult abdominal segments of *Drosophila* by *optomotor-blind*. *Development* **124**, 3703–3714 (1997).
14. Sanchez-Herrero, E., Vernos, I., Marco, R. & Morata, G. Genetic organization of the *Drosophila* bithorax complex. *Nature* **313**, 108–113 (1985).
15. Duncan, I. M. The bithorax complex. *Annu. Rev. Genet.* **21**, 285–319 (1987).
16. Celniker, S. E., Sharma, S., Keelan, D. J. & Lewis, E. B. The molecular genetics of the bithorax complex of *Drosophila*: cis-regulation in the *Abdominal-B* domain. *EMBO J.* **9**, 4277–4286 (1990).
17. Crosby, M. A., Lundquist, E. A., Tautvydas, R. M. & Johnson, J. The 3' regulatory region of the *Abdominal-B* gene: genetic analysis supports a model of reiterated and interchangeable regulatory elements. *Genetics* **134**, 809–824 (1993).
18. Hopmann, R., Duncan, D. & Duncan, I. Transvection in the *iab-5,6,7* region of the bithorax complex of *Drosophila*: homology independent interactions in *trans*. *Genetics* **139**, 815–833 (1994).
19. Baker, B. S. & Ridge, K. A. Sex and the single cell. I. On the action of major loci affecting cell determination in *Drosophila melanogaster*. *Genetics* **94**, 383–423 (1980).
20. Cline, T. W. & Meyer, B. J. Vive al difference: males vs. females in flies vs. worms. *Annu. Rev. Genet.* **30**, 637–702 (1996).
21. McKeown, M. Sex differentiation: the role of alternative splicing. *Curr. Opin. Genet. Dev.* **2**, 299–303 (1992).
22. Burtis, K. C. The regulation of sex determination and sexually dimorphic differentiation in *Drosophila*. *Curr. Opin. Cell Biol.* **5**, 1006–1014 (1993).
23. Jursnich, V. A. & Burtis, K. C. A positive role in differentiation for the male doublesex protein of *Drosophila*. *Dev. Biol.* **155**, 235–249 (1993).
24. Waterbury, J. A., Jackson, L. L. & Schedl, P. Analysis of the doublesex female protein in *Drosophila melanogaster*: Role in sexual differentiation and behavior and dependence on intersex. *Genetics* **152**, 1653–1667 (1999).
25. Robertson, A., Briscoe, D. A. & Louw, J. H. Variation in abdomen pigmentation in *Drosophila melanogaster* females. *Genetica* **47**, 73–76 (1977).
26. Godt, D., Couderc, J. -L., Cramton, S. E. & Laski, F. Pattern formation in the limbs of *Drosophila*: *bric a bric* is expressed in both a gradient and a wave-like pattern and is required for specification and proper segmentation of the tarsus. *Development* **119**, 799–812 (1993).
27. Pointud, J. C. Fly base personal communications report (<http://flybase.bio.indiana.edu/bin/fbpcq.html?FBrf0104845>) (1998).
28. Nagoshi, R. N. & Baker, B. S. Regulation of sex-specific RNA splicing at the *Drosophila doublesex* gene: cis-acting mutations in exon sequences alter sex-specific RNA splicing patterns. *Genes Dev.* **4**, 89–97 (1990).
29. Burtis, K. C., Coschigano, K. T., Baker, B. S. & Wensink, P. C. The Doublesex proteins of *Drosophila melanogaster* bind directly to a sex-specific yolk protein gene enhancer. *EMBO J.* **10**, 2577–2582 (1991).
30. Coschigano, K. T. & Wensink, P. C. Sex-specific transcriptional regulation by the male and female doublesex proteins of *Drosophila*. *Genes Dev.* **7**, 42–54 (1993).
31. Bock, I. R. Taxonomy of the *Drosophila bipectinata* species complex. *Univ. Texas Publ.* **7103**, 273–280 (1971).
32. Ohnishi, S. & Watanabe, T. K. Genetic analysis of color dimorphism in the *Drosophila montium* subgroup. *Jpn J. Genet.* **60**, 355–358 (1985).
33. Fisher, R. A. *The Genetical Theory Of Natural Selection* (Dover, New York, 1958).
34. Carson, H. L. in *Ecological Genetics: The Interface* (ed. Brussard, P. F.) 93–107 (Springer, New York, 1978).
35. Lande, R. Models of speciation by sexual selection on polygenic traits. *Proc. Natl Acad. Sci. USA* **78**, 3721–3725 (1981).
36. Kirkpatrick, M. Sexual selection and the evolution of female choice. *Evolution* **36**, 1–12 (1982).
37. Carson, H. L. in *Evolutionary Genetics* (eds Singh, R. S. & Krimbas, C. B.) 495–512 (Cambridge Univ. Press, Cambridge, 1999).
38. Holland, B. & Rice, W. R. Chase-away sexual selection: antagonistic seduction versus resistance. *Evolution* **52**, 1–7 (1998).
39. Cordts, R. & Partridge, L. Courtship reduces longevity of male *Drosophila melanogaster*. *Anim. Behav.* **52**, 269–278 (1996).
40. Kopp, A. & Duncan, I. Technical tips for analyzing gene expression in the pupal abdomen. *Dros. Info. Serv.* (in the press).
41. Bender, W. & Hudson, A. P. Element homing to the *Drosophila* bithorax complex. *Development* **127**, 3981–3992 (2000).
42. Kellerman, K. A. Mutations affecting the stability of the *fushi tarazu* protein of *Drosophila*. *Genes Dev.* **4**, 1936–1950 (1990).
43. Bock, I. R. & Wheeler, M. R. The *Drosophila melanogaster* species group. *Univ. Texas Publ.* **7213**, 1–102 (1972).
44. Bock, I. R. Current status of the *Drosophila melanogaster* species-group (Diptera). *Syst. Entomol.* **5**, 341–356 (1980).
45. Pelandakis, M. & Solignac, M. Molecular phylogeny of *Drosophila* based on ribosomal RNA sequences. *J. Mol. Evol.* **37**, 525–543 (1993).
46. Toda, M. J. *Drosophilidae* (Diptera) in Myanmar (Burma). VII. The *Drosophila melanogaster* species-group, excepting the *D. montium* species-subgroup. *Orient. Insects* **25**, 69–94 (1991).
47. Russo, C. A. M., Takezaki, N. & Nei, M. Molecular phylogeny and divergence times of drosophilid species. *Mol. Biol. Evol.* **12**, 391–404 (1995).

Supplementary information is available on Nature's World-Wide Web site (<http://www.nature.com>) or as paper copy from the London editorial office of Nature.

Acknowledgements

We thank F. Laski for *bab* stocks and antibodies; D. Godt for UAS-*bab* lines; W. Bender, S. Celniker and E. Sanchez-Herrero for the *Abd-B* enhancer trap and antibodies; J. David, Y. Fuyama, M. Kimura, J. Roote and the Bowling Green stock centre for various *Drosophila* species; L. Olds for the artwork; B. Holland, S. Nuzhdin, M. Servadio and J. True and T. Wittkopp for discussions. A.K. is a Howard Hughes Medical Institute fellow of the Life Sciences Research Foundation; I.D. is supported by an NIH grant; S.B.C. is an investigator at the Howard Hughes Medical Institute.

Correspondence and requests for materials should be addressed to S.B.C. (e-mail: sbcarrol@facstaff.wisc.edu).

10. Russell, D. A. & Dong, Z.-M. A nearly complete skeleton of a new troodontid dinosaur from the Early Cretaceous of the Ordos Basin, Inner Mongolia, People's Republic of China. *Can. J. Earth Sci.* **30**, 2163–2173 (1994).
11. Currie, P. J. Bird-like characteristics of the jaws and teeth of troodontid theropods (Dinosauria, Saurischia). *J. Vert. Paleont.* **7**, 72–81 (1987).
12. Clark, J. M., Perle, M. & Norell, M. A. The skull of *Erlicosaurus andrewsi*, a Late Cretaceous "Segnosaur" (Theropoda: Therizinosauridae from Mongolia). *Am. Mus. Novit.* **3115**, 1–39 (1994).
13. Ostrom, J. H. Osteology of *Deinonychus antirrhopus*, an unusual theropod dinosaur from the Lower Cretaceous of Montana. *Peabody Mus. Nat. Hist. Bull.* **30**, 1–165 (1969).
14. Wellnhofer, P. Das siebte Exemplar von *Archaeopteryx* aus den Solnhofener Schichten. *Archaeopteryx* **11**, 1–47 (1993).
15. Currie, P. J. New information on the anatomy and relationships of *Dromaeosaurus albertensis* (Dinosauria: Theropoda). *J. Vert. Paleont.* **15**, 576–591 (1995).
16. Martin, L. D. in *Origins of the Higher Groups of Tetrapods* (eds Schultze, H.-P. & Trueb, L.) 485–540 (Cormstock, Ithaca and London, 1991).
17. Zhou, Z. & Martin, L. D. Feathered dinosaur or bird?—a new look at the hand of *Archaeopteryx*. *Smithson. Contrib. Paleobiol.* **89**, 289–293 (1999).
18. Forster, C. A., Sampson, S. D., Chiappe, L. M. & Krause, D. W. The theropod ancestry of birds: new evidence from the Late Cretaceous of Madagascar. *Science* **279**, 1915–1919 (1998).
19. Makovicky, P. & Sues, H.-D. Anatomy and phylogenetic relationships of the theropod dinosaur *Microvenator celer* from the Lower Cretaceous of Montana. *Am. Mus. Novit.* **3240**, 1–27 (1998).
20. Holtz, T. R. Jr. The phylogenetic position of the Tyrannosauridae: implications for theropod systematics. *J. Paleont.* **68**, 1100–1117 (1994).
21. Norell, M. A. & Makovicky, P. J. Important features of the dromaeosaur skeleton: information from a new specimen. *Am. Mus. Novit.* **3215**, 1–28 (1997).
22. Currie, P. J. & Peng, J.-H. A juvenile specimen of *Saurornithoides mongoliensis* from the Upper Cretaceous of northern China. *Can. J. Earth Sci.* **30**, 2224–2230 (1994).
23. Currie, P. J., Rigby, J. K. & Sloan, R. E. in *Dinosaur Systematics: Perspectives and Approaches* (eds Carpenter, K. & Currie, P. J.) 108–125 (Cambridge Univ. Press, Cambridge, 1990).
24. Wellnhofer, P. A new specimen of *Archaeopteryx* from the Solnhofen Limestone. *Nat. Hist. Mus. Los Angeles County Sci. Ser.* **36**, 3–23 (1992).
25. Zhou, Z.-H. & Hou, L.-H. *Confuciusornis* and the early evolution of birds. *Vertebrata Palasiatica* **36**(2), 136–146 (1998).
26. Buffetaut, E., Suteethorn, V. & Tong, H.-Y. The earliest known tyrannosaur from the Lower Cretaceous of Thailand. *Nature* **381**, 689–691 (1996).
27. Xu, X., Tang, Z.-L. & Wang, X.-L. A therizinosaurid dinosaur with integumentary structures from China. *Nature* **399**, 350–354 (1999).
28. Clark, J. M., Hopson, J. A., Hernandez, R., Fastovsky, D. E. & Montellano, M. Foot posture in a primitive pterosaur. *Nature* **391**, 886–889 (1998).
29. Chiappe, L. M. Climbing *Archaeopteryx*? A response to Yalden. *Archaeopteryx* **15**, 109–112 (1997).
30. Chatterjee, S. *The Rise of Birds* (John Hopkins Univ. Press, Baltimore, 1997).

Supplementary information is available on Nature's World-Wide Web site (<http://www.nature.com>) or as paper copy from the London editorial office of Nature.

Acknowledgements

We thank D. Unwin, L. Chiappe, X.-C. Wu, J. Clark and L. Witmer for comments; M.-M. Chang and Y.-Q. Wang for assistance during the course of the work; H.-J. Wang for preparing the specimen; R.-S. Li for making the drawings; and J. Zhang and H.-L. You for taking the photographs. This work was supported by the Special Funds for Major State Basic Research Projects of China, and research grants from the National Geographic Society of the United States, Chinese Natural Science Foundation, and the Chinese Academy of Sciences.

Correspondence and requests for materials should be addressed to X.X. (e-mail: xxu@midwest.com.cn).

Mitochondrial genome variation and the origin of modern humans

Max Ingman*, Henrik Kaessmann†, Svante Pääbo† & Ulf Gyllenstein*

* Department of Genetics and Pathology, Section of Medical Genetics, Rudbeck Laboratory, University of Uppsala, S-751 85 Uppsala, Sweden

† Max Planck Institute for Evolutionary Anthropology, Inselstrasse 22, D-04103 Leipzig, Germany

The analysis of mitochondrial DNA (mtDNA) has been a potent tool in our understanding of human evolution, owing to characteristics such as high copy number, apparent lack of recombination¹, high substitution rate² and maternal mode of inheritance³. However, almost all studies of human evolution based on mtDNA sequencing have been confined to the control region, which constitutes less than 7% of the mitochondrial

genome. These studies are complicated by the extreme variation in substitution rate between sites, and the consequence of parallel mutations⁴ causing difficulties in the estimation of genetic distance and making phylogenetic inferences questionable⁵. Most comprehensive studies of the human mitochondrial molecule have been carried out through restriction-fragment length polymorphism analysis⁶, providing data that are ill suited to estimations of mutation rate and therefore the timing of evolutionary events. Here, to improve the information obtained from the mitochondrial molecule for studies of human evolution, we describe the global mtDNA diversity in humans based on analyses of the complete mtDNA sequence of 53 humans of diverse origins. Our mtDNA data, in comparison with those of a parallel study of the Xq13.3 region⁷ in the same individuals, provide a concurrent view on human evolution with respect to the age of modern humans.

The molecular clock hypothesis postulates that DNA sequence evolution is roughly constant over time in all evolutionary lineages. We used a test⁸ that compares the log likelihoods of trees reconstructed with and without the molecular clock assumption to examine the supposition that the mtDNA lineages evolve at 'clock-like' rates. The human mtDNA sequences, excluding the D-loop, have evolved at roughly constant rates ($P = 0.094$), and a relative rates test⁹, using a gorilla sequence as an outgroup, demonstrates that there is also no significant difference between the evolutionary rate of human and chimpanzee mtDNAs ($P = 0.123$), excluding the D-loop. In contrast, the D-loop has not evolved at a constant rate across all human lineages ($P < 0.001$), and is consequently less suitable for dating evolutionary events. Therefore, unless specifically mentioned, we have excluded the D-loop from the analyses that follow.

Figure 1 The relationship between linkage disequilibrium, measured by $|D'|$ versus distance between nucleotide sites for all 53 complete human mtDNA genomes. Values of ± 1.0 have been removed. **a**, Individuals of African descent ($n = 1,719$ comparisons), $R^2 = 0.001$; **b**, only non-African individuals ($n = 741$ comparisons), $R^2 = 0.005$.

Figure 2 Neighbour-joining phylogram based on complete mtDNA genome sequences (excluding the D-loop). Data was constructed using PAUP*4.0 Beta (Sinauer Associates) and bootstrapped with 1,000 replicates (bootstrap values shown on nodes). The population origin of the individual is given at the twigs. Branches have been colour coded

as in Fig. 4. Individuals of African descent are found below the dashed line and non-Africans above. The node marked with an asterisk refers to the MRCA of the youngest clade containing both African and non-African individuals.

Table 1 Summary of statistical parameters for the mtDNA

	Data set	Length	<i>n</i>	<i>S</i>	MPSD	π
Total	All humans	16,553	53	657	61.1	3.7×10^{-3}
	Non-Africans	16,555	32	358	38.5	2.3×10^{-3}
	Africans	16,556	21	367	76.7	4.6×10^{-3}
D-Loop	All humans	1,118	53	141	17.2	1.5×10^{-2}
	Non-Africans	1,118	32	103	12.8	1.1×10^{-2}
	Africans	1,121	21	77	19.7	1.8×10^{-2}
Rest	All humans	15,435	53	516	43.9	2.8×10^{-3}
	Non-Africans	15,437	32	255	25.7	1.7×10^{-3}
	Africans	15,448	21	290	57.0	3.7×10^{-3}

mtDNA data are given as entire sequences, or separated into D-loop and rest (all but the D-loop), and further separated into groups of African and non-African individuals. Length, aligned sequence length excluding gaps; *n*, number of sequences; *S*, number of segregating sites; MPSD, mean pairwise sequence difference; and π , genetic diversity.

From the mean genetic distance between all the humans and the one chimpanzee sequence (0.17 substitutions per site) and the assumption, based on palaeontological¹⁰ and genetic¹¹ evidence, of a divergence time between humans and chimpanzees of 5 Myr, the mutation rate (μ) for the mitochondrial molecule, excluding the D-loop, is estimated to be 1.70×10^{-8} substitutions per site per year.

On the basis of the correlation between linkage disequilibrium and distance between sites, it has been claimed that mtDNA sequences show signs of recombination^{12,13}. These analyses can be criticized for the methodology used¹⁴, particularly for the use of a linkage disequilibrium measure (Hill and Robertson measure, r^2) that doesn't take allele frequency into account. We examined linkage

disequilibrium among all informative sites in our set of complete mtDNA genomes (including the D-loop) using a standard estimate (D') that allows all variable positions to be examined¹⁵. For this analysis, we studied the sequences of Africans and non-Africans separately. There is no correlation between D' and nucleotide distance between sites in the 53 sequences (African correlation coefficient, $R^2 = 1 \times 10^{-3}$; non-African $R^2 = 5 \times 10^{-3}$) (Fig. 1). We also analysed the association between r^2 and distance, and again there is no evidence of a correlation ($R^2 = 2.23 \times 10^{-6}$, and 1×10^{-3} , respectively; data not shown). Thus, it does not appear to be necessary to consider recombination as contributing to the evolution of mtDNA.

The two main hypotheses for the evolution of modern humans agree that *Homo erectus* spread from Africa around 2 Myr ago. The 'recent African origin' hypothesis^{16,17} states that anatomically modern humans originated in Africa 100,000–200,000 years ago and subsequently spread to the rest of the world, replacing archaic human forms with little or no genetic mixing. The alternative, 'multi-regional' hypothesis proposes that the transformation to anatomically modern humans occurred in different parts of the world, and supports this with fossil evidence of cultural and morphological continuity between archaic and modern humans outside Africa¹⁸. There are, of course, variants of these two basic hypotheses that introduce additional assumptions, such as gene flow, that make the hypotheses increasingly difficult to test.

Support for a recent African origin of modern humans has been provided by a number of mtDNA studies^{16,17,19,20}; however, these results have been troubled by the lack of statistical support for tree topology, especially the deep African branches^{21,22}. Lacking sufficiently strong empirical data, it is impossible to confidently place the root of modern human mtDNA lineages in sub-Saharan Africa. The neighbour-joining (NJ) tree constructed from our mtDNA sequences has a strongly supported basal branching pattern (Fig. 2). The three deepest branches lead exclusively to sub-Saharan mtDNAs, with the fourth branch containing both Africans and non-Africans. The deepest, statistically supported branch (NJ bootstrap = 100) provides compelling evidence of a human mtDNA origin in Africa.

Figure 3 Mismatch distributions of pairwise nucleotide differences between mtDNA genomes (excluding the D-loop). **a**, African; **b**, non-African.

The amount of mtDNA sequence diversity (π) among Africans (3.7×10^{-3} nucleotide differences per site) is more than twice that among non-Africans (1.7×10^{-3}) (Table 1), corroborating earlier studies of the D-loop¹⁶ and nuclear loci²⁴. Also notable is the contrast between the deep branches of African mtDNAs and the 'star-like' phylogeny of non-African mtDNAs (Fig. 2). This high African diversity might result from either a considerably larger effective population size or a significantly longer genetic history. The 'star-like' phylogeny of the non-African sequences suggests a

Figure 4 Data matrices showing all informative nucleotide positions, in decreasing order of frequency. Left, the whole mtDNA genome, excluding the D-loop. Right, the D-loop. The

population bottleneck, potentially associated with the colonization of Eurasia from Africa. The date of this exodus from Africa can be estimated if the departing group subsequently experienced a population expansion. The mtDNA mismatch distributions for Africans and non-Africans indicate a marked difference in population history for the two groups²⁵ (Fig. 3). Mitochondrial DNAs from individuals of African origin show a ragged distribution consistent with constant population size, whereas the bell-shaped distribution of the non-African comparisons clearly indicates a recent popula-

tion expansion. The assumption of constant population size can be verified by tests of selective neutrality that examine the correlation between the mean pairwise sequence difference (MPSD) and the number of segregating sites (S)^{26,27}. In the African group, we cannot reject this assumption (Fu and Li's $D = -1.17$ (ref. 26); Tajima's $D = -1.22$ (ref. 27)), consistent with the premise that the population has been of roughly constant size. However, it can be rejected in the non-African group ($D = -4.02$; $D = -2.28$), indicating that this group has experienced a period of population growth²⁵. The

trees on the left are cladograms with the same topology and numbering of individuals as the tree in Fig. 2. Individuals of African descent are found below the dashed line and

non-Africans above. The four major groups of sequences have been colour coded as in Fig. 2. Blocks denote groups of nucleotides that are identical in several sequences.

time when the expansion began was estimated ($\tau = 20.23$) to be about 1,925 generations ago²⁸. Assuming a generation time of 20 yr this equates to 38,500 yr BP, a date that coincides with the onset of a period of cultural change about 35,000–40,000 years ago²⁹. This involves, for example, the first appearance of regional cultural variation and the acceleration of artefactual change.

The age of the most recent common ancestor (MRCA) for mtDNA, on the basis of the maximum distance between two humans (5.82×10^{-3} substitutions per site between the Africans Mkamba and San), is estimated to be $171,500 \pm 50,000$ yr BP. We can also estimate the age of the MRCA for the youngest clade that contains both African and non-African sequences (Fig. 2, asterisk) from the mean distance of all members of that clade to their common node (8.85×10^{-4} substitutions per site) as $52,000 \pm 27,500$ yr BP. Because genetic divergence is expected to precede the divergence of populations, this date can be considered as the lower bound for an exodus from Africa.

Notably, a group of six African sequences (Fig. 4a, sequences 33–38) are genetically distant to those of other Africans, but share a common ancestor with non-Africans. These lineages represent descendants of a population that evidently gave rise to all the non-African lineages. Whether the ancestors of these six extant lineages originally came from a specific geographic region is not possible to determine, but we note that these sequences are from five populations that are now geographically unrelated.

Our study of the entire mitochondrial genome has some significant distinctions from previous studies of the D-loop. Most notably, the sequences outside of the D-loop evolve in a roughly 'clock-like' manner, enabling a more accurate measure of mutation rate, and therefore improved estimates of times to evolutionary events. Also of importance is the strong statistical support for the tree topology that has been lacking in earlier investigations. The difference between the D-loop and the rest of the molecule is visually evident in the contrast between the jumbled arrangement of polymorphic sites in the D-loop and the clear haplotypes defined by the sites in the rest of the molecule (Fig. 4).

The use of largely the same individuals in this study, as in that of the nuclear Xq13.3 (ref. 7) region, provides a unique opportunity to compare the information gained from the two genetic systems. Because the X chromosome has an effective population size three times that of mitochondria, the MRCA of an X-chromosomal locus is expected to be three times higher. Thus, the age of the MRCA of Xq13.3 is in agreement with the mtDNA data (mtDNA: 171,500; Xq13.3: 479,000 yr BP). The results from Xq13.3 also concur with the mtDNA data with respect to the greater genetic diversity found among African individuals relative to non-Africans, but Xq13.3 shows a considerably lower difference in diversity between the two groups (mtDNA: African, 3.9×10^{-3} , non-African, 1.7×10^{-3} ; Xq13.3: 3.5×10^{-4} , 3.05×10^{-4}). Owing to its lower substitution rate, only 33 segregating sites were present in 69 sequences of 10.2 kilobases (kb) at Xq13.3, as compared with the 657 segregating sites in the mitochondrial dataset. Comparisons of the mtDNA and Xq13.3 sequences carried by specific individuals show little correlation between the two loci, as expected from the different modes of inheritance. For example, the two Warao Indians showing the highest similarity in mtDNA have, in fact, two of the most divergent sequences studied at the Xq13.3 locus. Others, such as the Saami and Mandenka sequences, are closely related at Xq13.3 but have relatively high mitochondrial divergence.

Our results indicate that the field of mitochondrial population genomics will provide a rich source of genetic information for evolutionary studies. Nevertheless, mtDNA is only one locus and only reflects the genetic history of females. For a balanced view, a combination of genetic systems is required. With the human genome project reaching fruition, the ease by which such data may be generated will increase, providing us with an evermore detailed understanding of our genetic history. □

Methods

Sampling strategy and mtDNA sequences

To assess the global genetic diversity in humans, while analysing a restricted number of samples, we studied 53 individuals representing 14 of the major linguistic phyla. This sampling strategy attempts to avoid the bias inherent in selecting individuals on the basis of current world demographics, such as current population size or geographic location⁵. To provide an opportunity for comparison, we selected, where possible, the same individuals as those used by a previous study⁷ for the analysis of the Xq13.3 region. All the complete mtDNA sequences are unique and vary in length from 16,558 to 16,576 base pairs (bp). From nearly 900 kb sequenced, 5 heteroplasmic sites were confidently identified. We identified a total of 657 segregating sites (141 in the D-loop; 516 outside) among these 53 individuals, of which 283 (80 in the D-loop; 203 outside) showed the same polymorphism in at least 2 individuals (Fig. 4). The pairwise sequence distances between mtDNAs, corrected for multiple substitutions⁴, vary from 6.0×10^{-5} substitutions per site between two South American Indians (Warao) to 6.8×10^{-3} substitutions per site between two Africans (Mbenzele pygmy and San). The average distance between mtDNA genomes is 3.8×10^{-3} substitutions per site.

PCR primers and sequencing

The primers we used for polymerase chain reaction (PCR) amplification have been described³⁰. Sequencing was performed on the PCR products directly using BigDye (Applied Biosystems) chemistry. Separation of sequencing ladders was performed on the ABI 377 instrument for automated fragment analysis. We sequenced both forward and reverse strands. Sequence analysis was performed using Sequencing Analysis 3.3 (Applied Biosystems), and sequence alignment was made with Sequencher 3.1.1 (Gene Codes).

Accession numbers

The chimpanzee sequence (GenBank accession no. D38113 and the gorilla sequence (accession no. D38114) that we used as outgroups were obtained from a public mitochondrial sequence database (MITOMAP: <http://www.gen.emory.edu/mitomap.html>).

Received 9 June; accepted 30 August 2000.

1. Olivio, P. D., Van de Walle, M. J., Laipis, P. J. & Hauswirth, W. W. Nucleotide sequence evidence for rapid genotypic shifts in the bovine mitochondrial DNA D-loop. *Nature* **306**, 400–402 (1983).
2. Brown, W. M., George, M. Jr & Wilson, A. C. Rapid evolution of animal mitochondrial DNA. *Proc. Natl Acad. Sci. USA* **76**, 1967–1971 (1979).
3. Giles, R. E., Blanc, H., Cann, H. M. & Wallace, D. C. Maternal inheritance of human mitochondrial DNA. *Proc. Natl Acad. Sci. USA* **77**, 6715–6719 (1980).
4. Tamura, K. & Nei, M. Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Mol. Biol. Evol.* **10**, 512–526 (1993).
5. Maddison, D. R., Ruvo, M. & Swofford, D. L. Geographic origins of human mitochondrial DNA: phylogenetic evidence from control region sequences. *Syst. Biol.* **41**, 111–124 (1992).
6. Torroni, A. et al. mtDNA analysis reveals a major late Paleolithic population expansion from southwestern to northeastern Europe. *Am. J. Hum. Genet.* **62**, 1137–1152 (1998).
7. Kaessmann, H., Heissig, F., von Haeseler, A. & Paabo, S. DNA sequence variation in a non-coding region of low recombination on the human X chromosome. *Nature Genet.* **22**, 78–81 (1999).
8. Strimmer, K. & von Haeseler, A. Quartet puzzling: a quartet maximum-likelihood method for reconstructing tree topologies. *Mol. Biol. Evol.* **13**, 964–969 (1996).
9. Sarich, V. M. & Wilson, A. C. Generation time and genomic evolution in primates. *Science* **179**, 1144–1147 (1973).
10. Andrews, P. Evolution and environment in the Hominoidea. *Nature* **360**, 641–646 (1992).
11. Kumar, S. & Hedges, S. B. A molecular timescale for vertebrate evolution. *Nature* **392**, 917–920 (1998).
12. Awadalla, P., Eyre-Walker, A. & Smith, J. M. Linkage disequilibrium and recombination in hominid mitochondrial DNA. *Science* **286**, 2524–2525 (1999).
13. Eyre-Walker, A., Smith, N. H. & Smith, J. M. How clonal are human mitochondria? *Proc. R. Soc. Lond. B* **266**, 477–483 (1999).
14. Kumar, S., Hedrick, P. & Stoneking, M. Questioning evidence for recombination in human mitochondrial DNA. *Science* **288**, 1931 (2000).
15. Lewontin, R. C. The interaction of selection and linkage. I. General considerations; heterotic models. *Genetics* **49**, 49–67 (1964).
16. Vigilant, L., Stoneking, M., Harpending, H., Hawkes, K. & Wilson, A. C. African populations and the evolution of human mitochondrial DNA. *Science* **253**, 1503–1507 (1991).
17. Cann, R. L., Stoneking, M. & Wilson, A. C. Mitochondrial DNA and human evolution. *Nature* **325**, 31–36 (1987).
18. Wolpoff, M. H. In *The Human Revolution: Behavioural and Biological Perspectives on the Origins of Modern Humans* (eds Mellars, P. & Stringer, C.) 62–108 (Princeton Univ. Press, Princeton, New Jersey, 1989).
19. Horai, S., Hayasaka, K., Kondo, R., Tsugane, K. & Takahata, N. Recent African origin of modern humans revealed by complete sequences of hominid mitochondrial DNAs. *Proc. Natl Acad. Sci. USA* **92**, 532–536 (1995).
20. Ruvo, M. et al. Mitochondrial COII sequences and modern human origins. *Mol. Biol. Evol.* **10**, 1115–1135 (1993).
21. Templeton, A. R. Human origins and analysis of mitochondrial DNA sequences. *Science* **255**, 737 (1992).
22. Nei, M. Age of the common ancestor of human mitochondrial DNA. *Mol. Biol. Evol.* **9**, 1176–1178 (1992).
23. Saitou, N. & Nei, M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**, 406–425 (1987).

24. Zietkiewicz, E. *et al.* Genetic structure of the ancestral population of modern humans. *J. Mol. Evol.* **47**, 146–155 (1998).
25. Rogers, A. R. & Harpending, H. Population growth makes waves in the distribution of pairwise genetic differences. *Mol. Biol. Evol.* **9**, 552–569 (1992).
26. Fu, Y. X. & Li, W. H. Statistical tests of neutrality of mutations. *Genetics* **133**, 693–709 (1993).
27. Tajima, F. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* **123**, 585–595 (1989).
28. Rozas, J. & Rozas, R. DnaSP, DNA sequence polymorphism: an interactive program for estimating population genetics parameters from DNA sequence data. *Comput. Appl. Biosci.* **11**, 621–625 (1995).
29. Klein, R. G. *The Human Career: Human Biological and Cultural Origins* (Univ. Chicago Press, Chicago, 1989).
30. Reider, M. J., Taylor, S. L., Tobe, V. O. & Nickerson, D. A. Automating the identification of DNA variations using quality-based fluorescence re-sequencing: analysis of the human mitochondrial genome. *Nucleic Acids Res.* **26**, 967–973 (1998).

Acknowledgements

We thank M. Stoneking for his advice regarding the analysis of recombination, and L. Cavalli-Sforza, G. Destro-Bisol, L. Excoffier, T. Jenkins, K. Kidd, J. Kidd, G. Klein, R. Mahabber, V. Nasidze, E. Poloni, H. Soodyall, M. Stoneking, M. Voevoda and S. Wells for samples. This work was supported by grants from Swedish Natural Sciences Research Council and Beijer Foundation.

Correspondence and requests for materials should be addressed to U.G. (e-mail: ulf.gyllensten@genpat.uu.se).

The *HIC* signalling pathway links CO₂ perception to stomatal development

Julie E. Gray*, Geoff H. Holroyd†, Frederique M. van der Lee‡, Ahmad R. Bahrami*, Peter C. Sijmons§, F. Ian Woodward||, Wolfgang Schuch¶ & Alistair M. Hetherington†

* Department of Molecular Biology and Biotechnology, University of Sheffield, Sheffield S10 2TN, UK

† Department of Biological Sciences, University of Lancaster, Lancaster LA1 4YQ, UK

‡ Zeneca Mogen, PO Box 628, 2300 AP Leiden, The Netherlands

§ Cellscreen, PO Box 134, 6700 AC Wageningen, Netherlands

|| Department of Animal and Plant Sciences, University of Sheffield, Sheffield, S10 2TN, UK

¶ Zeneca Wheat Improvement Centre, Norwich Research Park, Colney, Norwich NR4 7UH, UK

Stomatal pores on the leaf surface control both the uptake of CO₂ for photosynthesis and the loss of water during transpiration. Since the industrial revolution, decreases in stomatal numbers in parallel with increases in atmospheric CO₂ concentration have provided evidence of plant responses to changes in CO₂ levels caused by human activity^{1,2}. This inverse correlation between stomatal density and CO₂ concentration also holds for fossil material from the past 400 million years³ and has provided clues to the causes of global extinction events⁴. Here we report the

identification of the *Arabidopsis* gene *HIC* (for high carbon dioxide), which encodes a negative regulator of stomatal development that responds to CO₂ concentration. This gene encodes a putative 3-keto acyl coenzyme A synthase—an enzyme involved in the synthesis of very-long-chain fatty acids⁵. Mutant *hic* plants exhibit up to a 42% increase in stomatal density in response to a doubling of CO₂. Our results identify a gene involved in the signal transduction pathway responsible for controlling stomatal numbers at elevated CO₂.

As part of a promoter trap screen aimed at isolating tissue-specific genes in *Arabidopsis thaliana*^{6,7}, we identified an individual plant designated *hic* in which the β -glucuronidase (*GUS*) reporter gene was expressed only in the guard cells (Fig. 1) and nowhere else in the plant. Southern analysis of *hic* genomic DNA using a *GUS* gene probe detected a single band, suggesting that there was only one *GUS* gene insertion in this line (data not shown).

We next investigated the phenotype of the *hic* mutant and found it to be different from previously described *Arabidopsis* stomatal development mutants^{8,9}. *hic* stomata had no obvious phenotype and the plants were not wilting, suggesting that guard-cell function was not impaired. *hic* guard cells increased turgor in response to light and reduced turgor in response to the addition of the plant hormone abscisic acid in stomatal bioassays (data not shown). *hic* plants were indistinguishable from the parental ecotype when analysed by infrared thermography (data not shown); however, analysis of plants that had been grown under differing CO₂ concentrations showed that the *HIC* gene affects stomatal development.

We grew *hic* and C24, the parental ecotype, at elevated and ambient concentrations of CO₂, and measured the stomatal index and density on the leaf surface. A reduction in stomatal index and density in response to elevated CO₂ is well characterized in some species, including three ecotypes of *Arabidopsis*^{1,2}, and indicates that the plant has the potential for increased water-use efficiency with increasing CO₂. Table 1 shows that growth of *hic* plants under elevated CO₂ induced marked increases in both stomatal index and

Figure 1 *GUS* expression in *hic* *Arabidopsis* plants is specific to guard cells. Part of a *hic* leaf histochemically stained for *GUS* activity shows guard-cell-specific expression at low (a; scale bar, 100 μ m) and higher magnification (b; scale bar, 10 μ m).

Table 1 Effect of ambient and elevated CO₂ on stomatal and epidermal cell density and stomatal index

Experiment	Line	Stomatal density (n mm ⁻²)				Epidermal cell density (n mm ⁻²)				Stomatal index (%)			
		Ambient	Elevated	Difference (<i>P</i>)	% change	Ambient	Elevated	Difference (<i>P</i>)	% change	Ambient	Elevated	Difference (<i>P</i>)	% change
1	C24	280 (9)	233 (8)	<0.001	-17.0	1,047 (9)	902 (8)	<0.001	-13.8	21.1 (9)	20.5 (8)	NS	-3.1
	<i>hic</i>	231 (8)	320 (8)	<0.001	+38.0	891 (8)	902 (8)	NS	+1.2	20.8 (8)	26.0 (8)	<0.005	+25
2	C24	265 (7)	245 (9)	NS	-7.8	1,056 (7)	1,028 (9)	NS	-2.7	20.0 (7)	19.1 (9)	NS	-4.4
	<i>hic</i>	257 (9)	324 (10)	<0.001	+25.2	1,047 (9)	1,061 (10)	NS	+1.3	19.8 (9)	23.2 (10)	<0.001	+17.8
3	C24	324 (9)	245 (7)	<0.001	-24.5	1,322 (9)	1,058 (7)	<0.005	-20	19.6 (9)	18.7 (7)	NS	-4.7
	<i>hic</i>	273 (7)	387 (9)	<0.001	+41.8	1,053 (7)	1,092 (9)	NS	+3.7	20.5 (7)	26.2 (9)	<0.001	+27.6
4	C24	273 (12)	283 (12)	NS	+3.1	1,052 (12)	1,049 (12)	NS	-0.3	20.6 (12)	21.2 (12)	NS	+3.2
	<i>hic</i>	266 (12)	351 (12)	<0.001	+31.8	1,035 (12)	1,062 (12)	NS	+2.6	20.6 (12)	24.9 (12)	<0.001	+21.2

Data are presented for four separate experiments in the *hic* mutant and the parental ecotype (C24). The average density or stomatal index is presented for each treatment. The numbers in brackets indicate the number of individual plants in each treatment. Treatments were compared using the Mann-Whitney Rank Sum Test (SigmaStat 2.03). NS, not significant.