this diversity is located in discrete gene clusters that are spread throughout the different genomes. In contrast to this diversity, these enteric microorganisms exhibit marked synteny in their large-scale genomic organization, bearing in mind that *E. coli* and *S. enterica* diverged about 100 Myr ago²⁸. The conserved genes may be a reflection of the basic lifestyle of the bacteria, requiring intestine colonization, environmental survival and transmission. The unique gene clusters probably contribute to adaptation to environmental niches and to pathogenicity. The pseudogene complement of *S. typhi* has implications for our understanding of the tight host restriction of this organism, and raises the question of whether it may be possible to eradicate *S. typhi* and typhoid fever altogether. □

Methods

Salmonella typhi CT18 was isolated in December 1993, at the Mekong Delta region of Vietnam, from a 9-year-old girl who was suffering from typhoid. The strain was isolated from blood using routine culture methods²³, and after serological and metabolic confirmation of the strain as *S. typhi* it was immediately frozen in glycerol at -70 °C. The genome sequence was obtained from 97,000 end sequences (giving 7.9× coverage) derived from several pUC18 genomic shotgun libraries (with insert sizes ranging from 1.4 to 4.0 kb) using dye terminator chemistry on ABI377 automated sequencers. This was supplemented with 0.7× sequence coverage from M13mp18 libraries with similar insert sizes. End sequences from a larger insert plasmid (pSP64; 1.9× clone coverage, 10–14-kb insert size) and lambda (lambda-FIX-II; 0.4× clone coverage, 20–22-kb insert size) libraries were used as a scaffold, and the final assembly was verified by comparison with restriction-enzyme digest patterns using pulsed-field gel electrophoresis (data not shown). Total sequence coverage was 9.1×. The sequence was assembled, finished and annotated as described²⁹, using Artemis³⁰ to collate data and facilitate annotation. In addition we used a genefinder that was trained specifically for *S. typhi*, which uses a hidden Markov model with modules for the coding region, start and stop codons, and the ribosome-binding site (T.S.L. and A.K., unpublished data). The genome and proteome sequences of *S. typhi* and *S. typhimurium* or *E. coli* were compared in parallel to identify deletions and insertions using the Artemis Comparison Tool (ACT) (K. Rutherford, unpublished data; see also <http://www.sanger.ac.uk/Software/ACT/>). Pseudogenes had one or more mutations that would ablate expression, and were identified by direct comparison with *S. typhimurium*; each of the inactivating mutations was subsequently checked against the original sequencing data.

Received 4 April; accepted 20 July 2001.

- Ivanhoff, B. Typhoid fever, global situation and WHO recommendations. *Southeast Asian J. Trop. Med. Public Health* **26**, (Suppl. 2) 1–6 (1995).
- Reeves, M. W., Evans, G. M., Heiba, A. A., Plikaytis, B. D. & Farmer, J. J. Clonal nature of *Salmonella typhi* and its genetic relatedness to other salmonellae as shown by multilocus enzyme electrophoresis, and proposal of *Salmonella bongori* comb. nov. *J. Clin. Microbiol.* **27**, 313–230 (1989).
- Parry, C., Wain, J., Chinh, N. T., Vinh, H. & Farrar, J. J. Quinolone-resistant *Salmonella typhi* in Vietnam. *Lancet* **351**, 1289 (1998).
- Neidhardt, F. C. & Curtiss, R. *Escherichia coli* and *Salmonella: Cellular and Molecular Biology* (ASM Press, Washington DC, 1996).
- Rosenberg, S. M., Longrich, S., Gee, P. & Harris, R. S. Adaptive mutation by deletions in small mononucleotide repeats. *Science* **265**, 405–407 (1994).
- Townsend, S. M. *et al.* *Salmonella enterica* serovar Typhi possesses a unique repertoire of fimbrial gene sequences. *Infect. Immun.* **69**, 2894–2901 (2001).
- Hensel, M., Hinsley, A. P., Nikolaus, T., Sawers, G. & Berks, B. C. The genetic basis of tetrathionate respiration in *Salmonella typhimurium*. *Mol. Microbiol.* **32**, 275–287 (1999).
- Blanc-Potard, A. B., Solomon, F., Kayser, J. & Groisman, E. A. The SPI-3 pathogenicity island of *Salmonella enterica*. *J. Bacteriol.* **181**, 998–1004 (1999).
- Tsolis, R. M. *et al.* Identification of a putative *Salmonella enterica* serotype typhimurium host range factor with homology to IpaH and YopM by signature-tagged mutagenesis. *Infect. Immun.* **67**, 6385–6393 (1999).
- Miao, E. A. & Miller, S. I. A conserved amino acid sequence directing intracellular type III secretion by *Salmonella typhimurium*. *Proc. Natl Acad. Sci. USA* **97**, 7539–7544 (2000).
- Bakshi, C. S. *et al.* Identification of SopE2, a *Salmonella* secreted protein which is highly homologous to SopE and involved in bacterial invasion of epithelial cells. *J. Bacteriol.* **182**, 2341–2344 (2000).
- Kingsley, R. A. & Baumber, A. J. Host adaptation and the emergence of infectious disease: the *Salmonella* paradigm. *Mol. Microbiol.* **36**, 1006–1014 (2000).
- Sanderson, K. E. & Liu, S. L. Chromosomal rearrangements in enteric bacteria. *Electrophoresis* **19**, 569–572 (1998).
- Blattner, F. R. *et al.* The complete genome sequence of *Escherichia coli* K-12. *Science* **277**, 1453–1474 (1997).
- Marcus, S. L., Brumell, J. H., Pfeifer, C. G. & Finlay, B. B. *Salmonella* pathogenicity islands: big virulence in small packages. *Microbes Infect.* **2**, 145–156 (2000).
- Perna, N. T. *et al.* Genome sequence of enterohaemorrhagic *Escherichia coli* O157:H7. *Nature* **409**, 529–533 (2001).
- Wong, K. K. *et al.* Identification and sequence analysis of a 27-kilobase chromosomal fragment containing a *Salmonella* pathogenicity island located at 92 minutes on the chromosome map of *Salmonella enterica* serovar typhimurium LT2. *Infect. Immun.* **66**, 3365–3371 (1998).
- Hashimoto, Y., Li, N., Yokoyama, H. & Ezaki, T. Complete nucleotide sequence and molecular characterization of ViaB region encoding Vi antigen in *Salmonella typhi*. *J. Bacteriol.* **175**, 4456–4465 (1993).
- Mirol, S. *et al.* Isolation of a temperate bacteriophage encoding the type III effector protein SopE from an epidemic *Salmonella typhimurium* strain. *Proc. Natl Acad. Sci. USA* **96**, 9845–9850 (1999).

- Zhang, X. L. *et al.* *Salmonella enterica* serovar typhi uses type IVB pili to enter human intestinal epithelial cells. *Infect. Immun.* **68**, 3067–3073 (2000).
- McClelland, M. *et al.* Complete genome sequence of *Salmonella enterica* serovar Typhimurium LT2. *Nature*, **413**, 852–856 (2001).
- Sherburne, C. K. *et al.* The complete DNA sequence and analysis of R27, a large IncHI plasmid from *Salmonella typhi* that is temperature sensitive for transfer. *Nucleic Acids Res.* **28**, 2177–2186 (2000).
- Wain, J. *et al.* Quantitation of bacteria in blood of typhoid fever patients and relationship between counts and clinical features, transmissibility, and antibiotic resistance. *J. Clin. Microbiol.* **36**, 1683–1687 (1998).
- Hu, P. *et al.* Structural organization of virulence-associated plasmids of *Yersinia pestis*. *J. Bacteriol.* **180**, 5192–5202 (1998).
- Perry, R. D. & Fetherston, J. D. *Yersinia pestis*—etiologic agent of plague. *Clin. Microbiol. Rev.* **10**, 35–66 (1997).
- Prentice, M. B. *et al.* *Yersinia pestis* pFra shows biovar-specific differences and recent common ancestry with a *Salmonella enterica* serovar Typhi plasmid. *J. Bacteriol.* **183**, 2586–2594 (2001).
- Jing, D. H., Dong, F., Latham, G. J. & von Hippel, P. H. Interactions of bacteriophage T4-coded primase (gp61) with the T4 replication helicase (gp41) and DNA in primosome formation. *J. Biol. Chem.* **274**, 27287–27298 (1999).
- Doolittle, R. F., Feng, D. F., Tsang, S., Cho, G. & Little, E. Determining divergence times of the major kingdoms of living organisms with a protein clock. *Science* **271**, 470–477 (1996).
- Parkhill, J. *et al.* Complete DNA sequence of a serogroup A strain of *Neisseria meningitidis* Z2491. *Nature* **404**, 502–506 (2000).
- Rutherford, K. *et al.* Artemis: sequence visualization and annotation. *Bioinformatics* **16**, 944–945 (2000).

Supplementary information is available on Nature's World-Wide Web site (<http://www.nature.com>) or as paper copy from the London editorial office of Nature.

Acknowledgements

We would like to acknowledge the support of the Sanger Centre core sequencing and informatics groups. We are grateful to the *S. typhimurium* sequencing group at Washington University for sharing sequence and analysis data before publication. A.K. and T.S.L. were supported by a grant from the Danish National Research Foundation. This work was supported by the Wellcome Trust through its Beowulf Genomics initiative and by a Wellcome Trust Programme Grant to G.D.

Correspondence and requests for materials should be addressed to J.P. (e-mail: parkhill@sanger.ac.uk) or G.D. (e-mail: g.dougan@ic.ac.uk). The sequences have been submitted to EMBL with the accession numbers AL513382 (chromosome), AL513383 (pHCM1) and AL513384 (pHCM2).

Complete genome sequence of *Salmonella enterica* serovar Typhimurium LT2

Michael McClelland*, Kenneth E. Sanderson†, John Spieth‡, Sandra W. Clifton‡, Phil Latreille‡, Laura Courtney‡, Steffen Porwollik*, Johar Ali‡, Mike Dante‡, Feiyu Du‡, Shunfang Hou‡, Dan Layman‡, Shawn Leonard‡, Christine Nguyen‡, Kelsi Scott‡, Andrea Holmes‡, Neenu Grewal‡, Elizabeth Mulvaney‡, Ellen Ryan‡, Hui Sun‡, Liliana Florea§, Webb Miller§, Tamberlyn Stoneking‡, Michael Nhan‡, Robert Waterston‡ & Richard K. Wilson‡

* Sidney Kimmel Cancer Center, 10835 Altman Row, San Diego, California 92121, USA

† Department of Biological Sciences, University of Calgary, Calgary, Alberta T2N 1N4, Canada

‡ Genome Sequencing Center, Washington University School of Medicine, 4444 Forest Park Boulevard, St Louis, Missouri 63108, USA

§ Department of Computer Science and Engineering, The Pennsylvania State University, University Park, Pennsylvania 16802, USA

Salmonella enterica subspecies I, serovar Typhimurium (*S. typhimurium*), is a leading cause of human gastroenteritis, and is used as a mouse model of human typhoid fever¹. The incidence of non-typhoid salmonellosis is increasing worldwide^{2–4}, causing millions of infections and many deaths in the human population each year. Here we sequenced the 4,857-kilobase (kb) chromosome

Present address: Celera Genomics, Rockville, Maryland 20850, USA.

and 94-kb virulence plasmid of *S. typhimurium* strain LT2. The distribution of close homologues of *S. typhimurium* LT2 genes in eight related enterobacteria was determined using previously completed genomes of three related bacteria, sample sequencing of both *S. enterica* serovar Paratyphi A (*S. paratyphi* A) and *Klebsiella pneumoniae*, and hybridization of three unsequenced genomes to a microarray of *S. typhimurium* LT2 genes. Lateral transfer of genes is frequent, with 11% of the *S. typhimurium* LT2 genes missing from *S. enterica* serovar Typhi (*S. typhi*), and 29% missing from *Escherichia coli* K12. The 352 gene homologues of *S. typhimurium* LT2 confined to subspecies I of *S. enterica*—containing most mammalian and bird pathogens³—are useful for studies of epidemiology, host specificity and pathogenesis. Most of these homologues were previously unknown, and 50 may be exported to the periplasm or outer membrane, rendering them accessible as therapeutic or vaccine targets.

The genus *Salmonella* comprises two species: *S. enterica*, which is subdivided into over 2,000 serovars, and *Salmonella bongori*. Some serovars of *S. enterica*, such as *S. typhi*, cause systemic infections and typhoid fever, whereas others, such as *S. typhimurium*, cause gastroenteritis. Some serovars, such as *S. typhi*, are host specialists that infect only humans, whereas others such as *S. typhimurium*, are host generalists that occur in humans and many other mammalian species. Domestic animals act as a reservoir for the food-borne spread of host-generalist serovars, which accounts for the high incidence of non-typhoid *Salmonella* infections worldwide. Estimated costs of food-borne diseases in the United States (with salmonellosis a major component) range from 4.8 to 23 billion dollars³.

Salmonella typhimurium strain LT2, the principal strain for cellular and molecular biology in *Salmonella*, was isolated in the 1940s and used in the first studies on phage-mediated transduction¹. Attenuated mutants of *S. enterica* may be used as live oral vaccines against *Salmonella* infection, to express antigens from other pathogens, and to deliver proteins to solid tumours⁶.

The general characteristics of the *S. typhimurium* LT2 genome are summarized in Table 1. The full dataset is presented as Supplementary Information and is available also at <http://genome.wustl.edu/gsc/Projects/S.typhimurium>.

The overall similarity of *S. typhimurium* LT2 to eight other enterobacterial genomes is summarized in Table 2. We compared *S. typhimurium* LT2 to three other fully sequenced genomes: *S. typhi*⁷, the principal cause of typhoid in humans; *E. coli* K12 (ref. 8), a non-pathogen; and *E. coli* O157:H7, an enterohaemorrhagic strain⁹. (*Escherichia coli* is a member of the closest known genus to *Salmonella*.) We sequenced genome samples over 97% coverage from *S. enterica* serovar *S. paratyphi* A, a frequent cause of typhoid, and *K. pneumoniae*, an opportunistic human pathogen closely related to *Salmonella*. Visual representations of the comparisons of *S. typhimurium* LT2 to the sequenced and sampled genomes are available at both gene and sequence resolution using STM-Enteric and STM-Menteric web tools (<http://galapagos.cse.psu.edu/enterix>)^{10,11}. A total of 4,330 complete open reading frames (ORFs) from the *S. typhimurium* LT2 genome were amplified using specific PCR primers with appended, common 5' ends. These ORFs were microarrayed on glass and probed with fluorescently labelled

Table 2 Coding sequence homologues shared by *S. typhimurium* LT2 and eight other strains of enterobacteria

Organism	Data source	Homologues of STM CDS (%) ^a	Median homology of reciprocal best hit CDS (%)	
			DNA	Amino acid
<i>S. enterica</i> serovar <i>S. typhimurium</i> LT2	Complete sequence	100	100	100
<i>S. typhi</i> CT18	Complete sequence ⁷	89	98	99%
<i>S. paratyphi</i> A	~97% sequence sample/microarray	87/89	98	99
<i>S. paratyphi</i> B	Microarray	92	—	—
<i>S. arizonae</i>	Microarray	83	—	—
<i>S. bongori</i>	Microarray	85	—	—
<i>E. coli</i> K12	Complete sequence ⁸	71	80	90
<i>E. coli</i> O157:H7	Complete sequence ⁹	73	80	89
<i>K. pneumoniae</i>	~97% sequence sample	73	76	88

^aFor sequenced genomes, reciprocal best hits, excluding unsampled regions; for microarrays, signal ratio of *S. typhimurium* LT2 with genome is 3:1 or greater and based on roughly 4,330 *S. typhimurium* LT2 CDS. Raw data is available at <http://genome.wustl.edu/gsc/Projects/S.typhimurium>.

genomic DNA¹² from *S. bongori* and from *S. enterica* serovars *S. paratyphi* A, *S. paratyphi* B and *S. arizonae*, to determine the presence of homologues of *S. typhimurium* LT2 genes. The microarray data for *S. paratyphi* A was over 98% concordant with data from the *S. paratyphi* A genomic sequence (see Supplementary Information). *Salmonella typhimurium* LT2, *S. typhi*, *S. paratyphi* A and *S. paratyphi* B are all in subspecies I of *S. enterica*, which colonizes mammals and birds and causes 99% of *Salmonella* infections in humans^{5,13}. *Salmonella arizonae* is in subspecies IIIa, which colonizes reptiles and also, rarely, humans; *S. bongori*, the only other species of *Salmonella*, does not cause disease in mammals.

Genomic comparisons among the four completed genomes (*S. typhimurium* LT2, *S. typhi*, *E. coli* K12 and *E. coli* O157:H7) reveal that they are collinear for most genes except for inversions over the terminus of replication (TER)^{9,14}. *Salmonella typhimurium* LT2 has a 588-kb inversion compared with *E. coli* K12. Rearrangements between the *rrn* operons are common in *S. typhi*^{7,15}, but are not found in *E. coli* K12, *E. coli* O157:H7 or *S. typhimurium* LT2, or in most other *Salmonella* strains. Large, stable duplications are rare in enterobacterial genomes other than structural RNA genes and transposable elements. In *S. typhimurium* LT2 the largest duplicated region of coding sequences (CDS) is the cytochrome *c* biogenesis locus (*ccm*, 7.5 kb). The two copies are 99% identical at the DNA level and 100% identical for amino acids.

The chromosomes of enteric bacteria are mosaics, composed of collinear regions interspersed with 'loops' or 'islands' unique to certain species; the islands sometimes encode pathogenicity functions (called *Salmonella* pathogenicity islands, SPIs)^{9,16}. Of the 4,489 CDS and pseudogenes annotated in the *S. typhimurium* LT2 chromosome, at least 2,466 (55%) have a close homologue in all eight other enterobacterial genomes that we examined. These homologous genes (shown in red in Fig. 1) along with structural RNAs constitute 2.5 Mb of the *S. typhimurium* LT2 genome.

Table 3 summarizes the distribution of the larger islands among the eight genomes, on the basis of sequence and microarray analysis. SPIs 1–5, previously described in *S. typhimurium* (see refs 1, 17), are listed along with many newly detected large islands and the subset of smaller islands that encode fimbriae or potential virulence factors. Fifteen of the islands are adjacent to a transfer RNA. In fact, almost half of the individually encoded tRNAs are adjacent to an island. At least seven islands encode integrase-like proteins, which are often found in islands¹⁶. *Salmonella typhimurium* LT2 contains four

Table 1 Genome essentials

Parameter	Chromosome	Plasmid pSLT
Size (bp)	4,857,432	93,939
G+C content	53%	53%
rRNA clusters	7	0
tRNAs	85	0
tRNA pseudogene	1	0
Structural RNAs	11	1
CDS (including pseudogenes)	4,489	108
CDS pseudogenes	39	6

functional prophages: Gifsy-1 and -2 (known to have a role in infection^{18,19}) and Fels-1 and -2 (ref. 12). The insertion site of these prophages was predicted from the sequence and was confirmed by microarray-based comparison of labelled DNA from strains that were cured of prophage. These phage are not present in the other eight genomes, although homologues of a few genes exist, presumably in related phage. A previously unknown phage, or phage remnant, that included the *S. typhimurium* LT2 genes *STM4201* to *STM4219* was detected in *S. typhimurium* LT2 by homology to other phage.

Most strains of *S. typhimurium* contain a plasmid of about 90 kb. The plasmid of strain LT2 is called pSLT²⁰. Out of 108 annotated CDS and pseudogenes in pSLT, only three have a close homologue in *S. typhi*, *S. paratyphi* A or *S. paratyphi* B, as expected owing to these strains lacking the plasmid. A search through GenBank revealed 50 pSLT genes that have a close homologue in plasmids from other *Salmonella* serovars. Many homologues of genes in the *tra* operon of the F-factor of *E. coli* K12 were identified, which presumably are responsible for the self-transmissibility of pSLT at rates of up to 3×10^{-4} (ref. 21). The copy number of the pSLT plasmid was estimated as 2.75, using the relative sequence coverage of the plasmid versus the chromosome in the shotgun phase of sequencing, and estimated as 1.4–3.1 under a variety of growth conditions, when measured by average signal intensity on microarrays (data not shown).

Fimbriae on the cell surface mediate adhesion to host cells²². *Salmonella typhimurium* LT2 contains 12 putative operons of the chaperone–usher assembly class: *stc* (called *yehABCD* in *E. coli* K12), *bcf*, *fim*, *lpf*, *saf*, *stb*, *std*, *stf*, *sth*, *sti*, *stj*, all of which are located on the chromosome, and *pef*, which is located on the plasmid. Operons *bcf*, *fim*, *lpf* and *pef* were reported earlier for *S. typhimurium* LT2 and shown to be functional; *saf*, *stb*, *std* and *sth* were detected only by hybridization²². The *sti* and *stj* operons were previously undetected; thus, six of the operons were not previously sequenced and two were not previously detected. *Salmonella typhimurium* LT2 also has the *csg* operon (originally called *agf*) for curli fimbriae, from the nucleator-dependent assembly pathway, but genes for type IV fimbriae⁷ are not detected. Table 3 describes the taxonomic distribution of close homologues of the fimbriae gene clusters in the other eight enterobacterial genomes, but does not include information on the presence of more divergent homologues or orthologues in the other genomes.

Complete sequencing of many closely related genomes, such as *E. coli* K12, *E. coli* O157:H7, *S. typhi* and *S. typhimurium* LT2 (see Table 2), aids the detection of pseudogenes, because a frameshift or stop codon is recognizable only if the gene is collinear with a functional, homologous gene in another genome. This allowed the detection of at least 204 pseudogenes in *S. typhi*⁷, whereas the *S. typhimurium* LT2 chromosome has only about 39. The large number of pseudogenes in *S. typhi* may contribute to or be a consequence of the restriction of *S. typhi* to growth in humans alone⁷, whereas *S. typhimurium* LT2, with a broad host range, has far fewer pseudogenes. Pseudogenes may be unrecognized when a close, intact homologue is unavailable, as is true for 11% or more of the *S. typhimurium* LT2 and *S. typhi* genomes (Table 2). Other potential pseudogenes are encoded across insertion/deletion events that distinguish *S. typhimurium* LT2 and *S. typhi*, such as *S. typhimurium* LT2 gene *STM0098*, which is split by an insertion of about 2,000 bp in *S. typhi* and *S. paratyphi* (or a deletion of this size in *S. typhimurium* LT2). Such genes have been annotated as insertion/deletion events rather than pseudogenes (see Supplementary Information). Finally, pseudogenes might actually have at least one functional fragment. Nevertheless, these caveats do not alter the conclusion that *S. typhimurium* LT2 has far fewer pseudogenes than its close relative *S. typhi*.

The consequences of loss of function in pseudogenes in *S. typhimurium* LT2 is usually unclear, as the function of the known, intact homologue in another organism is often unknown. However, there are pseudogenes in *S. typhimurium* LT2 for maltose

regulation (*malXY*²³) and for trehalose metabolism (*treB*), where the normal allele can substitute in the maltose pathway²⁴; thus, *S. typhimurium* LT2 has disrupted maltose pathways with multiple, independent mutations. Histidine is not used as a carbon source in LT2, although all of the genes are present; this may be explained by a pseudogene mutation in *hutU*. Some of the pseudogenes are in potentially redundant systems: the *dcoA* pseudogene shows over 98% amino-acid identity to another putative sodium ion pump, oxaloacetate decarboxylase alpha chain (*STM3352*); *cutF*

Table 3 Islands and prophages

Start (by STM gene number)	Number of genes	Homologues	Comments
2584*	53	†	Gifsy-1 phage
1005*	52	†	Gifsy-2 phage, <i>sodC</i> , <i>msgA</i> homologue
2865	50	SAL‡	SPI1. Type III secretion system
2694*	47	†	Fels-2 phage
1379§	44	SAL‡	SPI2. Type III secretion system, part missing in SBO
0266*§	42	SS1†	<i>saf</i> fimbriae, <i>virG</i> , <i>sapA</i> , <i>tia</i> and <i>clpB</i> homologues
0893*	40	†	Fels-1 phage, <i>sodC</i> homologue
2019	40	SEN§, KPN	<i>cbi</i> and <i>pdu</i> clusters
0328	37	SS1†	<i>stb</i> fimbriae, <i>ail</i> homologue
1528	35	SAL‡	<i>pqaA</i>
1239§	31	SAL‡	<i>envEF</i> , <i>msgA</i> , <i>pagCD</i>
2741*	27	STM†, KPN‡	Adjacent Fels-2
0014	25	SAL‡	<i>bcf</i> fimbriae
4195	25	†	Many phage homologues
2770	19	SAL‡	<i>fljAB</i> , <i>hin</i> , <i>iroDEN</i> , <i>virK</i> , <i>pipB</i> homologue
4418	19	SPB†	<i>srj</i>
1854*	18	SAL‡	<i>sopE2</i> , <i>pagOK</i>
3117§	18	SAL‡	
3191§	18	SAL	
2230§	16	SPB‡	<i>oafA</i> , <i>sspH2</i> , <i>msgA</i> homologues
4305§	16	SAL‡	<i>phoN</i>
1911	15	SAL, <i>E. coli</i>	<i>flh</i> , flagella, <i>che</i> , <i>mot</i>
1171	14	SAL‡	<i>flg</i> flagella
0715	13	SS1†	Cell wall biogenesis?
3752§	13	SS1†, SBO‡	SPI3 (part) <i>mgtBC</i>
0052	12	SS1†, KPN‡	<i>citDEF</i>
2082	12	SS1	
3025§	12	SS1†	<i>std</i> fimbriae
4440	11	SAL‡, KPN	
0543*§	10	SEN†	<i>ftm</i> fimbriae
1629	10	SAL‡	<i>sseJ</i>
1664	10	SAL‡	<i>invC</i> homologue
3766	10	SS1†, SBO‡	SPI3 (part)
4488*§	9	†	
1087§	8	SAL	SPI5. <i>pip</i> cluster and <i>sopB</i>
3351	8	SEN, KPN	
0426	7	SS1, KPN	<i>phn</i>
0571	7	SAL‡, KPN	
1127§	7	SEN	
1139	7	SAL, <i>E. coli</i>	<i>csg</i> curli fimbriae
3527	7	SPB†	
0195	6	SPA, SPB	<i>stf</i> fimbriae
0854	6	SEN	
1491	6	SAL	
2175	6	SAL, ECH	
2340	6	SAL	
2513	6	SAL‡	<i>shdA</i> , <i>sinIH</i>
3079	6	SPB	
3779	6	SS1†	SPI3 (part)
4010	6	STY, SPA	
4257	6	SAL‡	SPI4
4591	6	SAL‡	<i>sth</i> fimbriae
1328	5	SAL‡	<i>wbbH</i>
2357	5	SEN†	<i>oadABG</i>
3636	5	SPB, SBO	<i>lpf</i> fimbriae
4571	5	†	<i>stj</i> fimbriae
0174	4	SPB	<i>sti</i> fimbriae
1113	4	SEN	<i>scsABCD</i>
1599	4	SAL‡	<i>pdgL</i> , <i>ugtL</i> , <i>sifB</i>
2063	4	SAL	<i>phsABC</i> , <i>sopA</i>
2149	4	STY, SPB	<i>stc</i> fimbriae
2396§	4	SEN, KPN	<i>pgtABCEP</i>

Distribution of *S. typhimurium* LT2 CDS in eight other genomes. Not all islands consisting of five genes or fewer are listed. STM, *S. typhimurium* LT2; SPA, *S. paratyphi* A; SPB, *S. paratyphi* B; STY, *S. typhi*; KPN, *K. pneumoniae*; ECH, *E. coli* O157:H7; SBO, *S. bongori*; SS1 includes STM, SPA, SPB and STY; SEN includes SS1 and *S. arizonae*; and SAL includes SEN and *S. bongori*.

* Encodes integrase.

† Some genes in this cluster have close homologues outside the group.

‡ Some genes in this cluster do not have close homologues in all members of the group.

§ Adjacent to a tRNA.

(STM0241), which encodes the copper homeostasis protein, is a pseudogene, but *cutC* (which remains intact) is sufficient for the same function²⁵.

Genes found only in *Salmonella* (see Table 3 for examples) may have been recruited since the divergence from *Escherichia coli* and *Klebsiella*, or they may have been lost from these genera. There are 1,106 *S. typhimurium* LT2 CDS in this class (marked in green in Fig. 1) that have a close homologue in at least one of the other five *Salmonella* examined. Many *S. typhimurium* LT2 genes associated with pathogenesis are in this class, including invasion genes such as *inv* and *prg*; proteins exported by the type III secretion system, such as SopB, SopD, SopE2, SipA, SipB, SipC and SptP; and some secretory system genes, such as members of *ssa* and *sse*. Most of these genes have homologues in *S. bongori*, indicating that these characteristics are maintained by even the most divergent of *Salmonella*. A subset of 352 *S. typhimurium* LT2 CDS (8%) have close homologues in one or more of the other three subspecies I genomes (*S. typhi*, *S. paratyphi* A and *S. paratyphi* B) but not in *S. arizonae*, *S. bongori*, *E. coli* K12, *E. coli* O157:H7 or *K. pneumoniae*. These genes, most of which were unknown previously, may include genes for specialization of subspecies I to warm-blooded hosts. Among these 352 *S. typhimurium* LT2 CDS are *bigA*, *envF*, *shdA*, *sifAB*, *sinH*, *srff*, *srgAB* (homologues of *ail* and *clpB*), and some genes in the fimbrial clusters, *saf*, *stb*, *stc*, *std*, *stf* and *sti*. Only 70 genes (20%) are named, whereas 49% of all CDS are named, indicating that the group has remained largely unstudied. There are 121 *S. typhimurium* LT2 CDS that have no close homologues in any of the other eight genomes, only a few of which are named, including individual genes from the *pef* and *stj* fimbriae, *spvR*, and a homologue of *mvpA*.

CDS rich in A+T are almost threefold over-represented among genes that have no close homologues outside subspecies I, confirming previous observations made with a smaller set of such genes²⁶. It

is not obvious why CDS confined to only some *Salmonella* should so often be A+T-rich. There is no such bias among G+C-rich genes. If some of these genes are of recent foreign origin then perhaps there is a preferential A+T-rich source for recruited genes.

Attenuated *S. typhimurium* has been used both as a vaccine and for expression of heterologous proteins for vaccines against other organisms²⁷. Proteins that are predicted to be exported to the periplasm or outer membrane, or beyond, are of interest for vaccines and therapy, as they would be exposed for targeting. PSORT²⁸, which predicts cellular location from protein sequence, predicts 251 outer-membrane proteins and 347 periplasmic proteins in *S. typhimurium* LT2, 182 of which are missing from *E. coli* K12, *E. coli* O157:H7 or *K. pneumoniae*, and about 50 of which are confined to subspecies I.

Antigens from all of the predicted surface and secreted CDS can be cloned, expressed on cells, and tested for immunological response, as was done recently for CDS conserved among *Neisseria meningitidis*²⁹. As a first step, we have amplified by PCR almost all the *S. typhimurium* LT2 CDS in a manner suitable for expression, and the distribution of close homologues among eight other enterobacterial genomes has been determined for each gene. □

Methods

Genome sequencing

The complete 4.8-Mb genome of *S. enterica* serovar Typhimurium strain LT2 (American Type Culture Collection, ATCC, number 700720) was sequenced using a combination of two approaches. The first approach was fivefold, whole-genome shotgun sampling, and the second was three- to tenfold sampling of gel-purified restriction fragments ranging in size from 43.3 to 570 kb¹⁴. Sonicated and size-fractionated DNA (approximately 1.5 kb) was cloned into M13 and plasmid vectors. Subclones were sequenced using dye-primer and dye-terminator chemistry on ABI 377 and 3700 sequencing machines. We assembled 92,648 sequence reads, representing ~7.7-fold final coverage, using the PHRAP assembly program (P. Green, unpublished data). The assembled genome sequence is in agreement

Figure 1 The *Salmonella enterica* serovar Typhimurium LT2 genome. **a**, The chromosome. Base pairs are indicated outside the outer circle. The outer two circles represent the coding orientation, with the forward strand on the outside and the reverse strand on the inside. Red indicates close homologues in all eight genomes. Green indicates genes with a close homologue in at least one other *Salmonella* (*S. typhi*, *S. paratyphi* A, *S. paratyphi* B, *S. arizonae* or *S. bongori*) but not in *E. coli* K12, *E. coli*

O157:H7 and *K. pneumoniae*. Blue indicates genes present only in *S. typhimurium* LT2. Grey indicates other combinations. The black inner circle is the G+C content; the purple/yellow innermost circle is the GC bias. The positions of the origin of replication (ORI) and terminus (TER) are shown. **b**, The plasmid pSLT. Base pairs are indicated outside the outer circle. The plasmid is not to scale. The colouring scheme is the same as for **a**.

with the physical map of the *S. typhimurium* LT2 genome¹⁴, with reads from both ends of about 372 lambda DASH-II clones with inserts of 13–20 kb, and with a *S. typhimurium* LT2 bacterial artificial chromosome library (prepared by R. Wing) restriction finger-printed at the Genome Sequencing Center. Two other genomes were partially sequenced using whole-genome, shotgun sample sequencing: 27,670 sequence reads of *S. enterica* serovar Paratyphi A strain RKS4993 (ATCC 9510) yielded a 3.7-fold coverage (97.5%) for this ~4.6 Mb genome, having 894 contigs with a predicted average gap size of ~200 bp; 36,848 sequence reads of *K. pneumoniae* strain MGH78578 (ATCC 700721), a clinical isolate from sputum, yielded a 3.9-fold depth of coverage (98%) for this ~5.6 Mb genome, resulting in 711 contigs with a predicted average gap size of ~160 bp.

Annotation

The primary annotation database was Acedb (<http://www.acedb.org>). Protein gene predictions were performed using GeneMark (<http://opal.biology.gatech.edu/GeneMark>) and Glimmer (<http://www.tigr.org/softlab/glimmer/glimmer.html>). Predicted proteins were searched against the protein family database, Pfam 5.5 (<http://pfam.wustl.edu>), against the COG database to find putative orthologues in other completed genomes³⁰, and against the protein localization prediction software, PSORT²⁸. The annotation was compared with the *S. typhi* annotation from the Sanger Centre and all differences in CDS predictions and start predictions were reassessed, on the basis of choice of start codon, length, conservation in Enterobacteriaceae and presence of identifiable motifs. The details of microarray construction and hybridization with genomic probes is described elsewhere¹². Strains, lambda and BAC clones are available from the *Salmonella* Genetic Stock Centre, <http://www.ualgary.ca/~kesander>.

Received 10 April; accepted 24 September 2001.

- Neidhardt, F. C. (ed. in chief) *Escherichia coli and Salmonella: Cellular and Molecular Biology* (ASM, Washington DC, 1996).
- Chalker, R. B. & Blaser, M. J. A review of human salmonellosis: III. Magnitude of *Salmonella* infection in the United States. *Rev. Infect. Dis.* **10**, 111–124 (1988).
- Todd, E. Epidemiology of foodborne illness: North America. *Lancet* **336**, 788–790 (1990).
- Cooke, E. M. Epidemiology of foodborne illness: UK. *Lancet* **336**, 790–793 (1990).
- Selander, R. K., Li, J. & Nelson, K. in *Escherichia coli and Salmonella: Cellular and Molecular Biology* (eds Neidhardt, F. C. et al.) 2691–2707 (ASM, Washington DC, 1996).
- Bermudes, D., Low, B. & Pawelek, J. Tumor-targeted *Salmonella*. Highly selective delivery vectors. *Adv. Exp. Med. Biol.* **465**, 57–63 (2000).
- Parkhill, J. et al. Complete genome sequence of a multiple drug resistant *Salmonella enterica* serovar Typhi CT18. *Nature* **413**, 848–852 (2001).
- Blattner, F. R. et al. The complete genome sequence of *Escherichia coli* K-12. *Science* **277**, 1453–1474 (1997).
- Perna, N. T. et al. Genome sequence of enterohaemorrhagic *Escherichia coli* O157:H7. *Nature* **409**, 529–533 (2001).
- Florea, L. et al. Web-based visualization tools for bacterial genome alignments. *Nucleic Acids Res.* **28**, 3486–3496 (2000).
- McClelland, M. et al. Comparison of the *Escherichia coli* K-12 genome with sampled genomes of a *Klebsiella pneumoniae* and three *Salmonella enterica* serovars, Typhimurium, Typhi and Paratyphi. *Nucleic Acids Res.* **28**, 4974–4986 (2000).
- Porwollik, S. et al. The Δ -uvrB mutations in the Ames strains of *Salmonella* span 15 to 119 genes. *Mut. Res.* **483**, 1–11 (2001).
- Popoff, M. Y., Bockemuhl, J. & Brenner, F. W. Supplement 1999 (no. 43) to the Kauffmann-White scheme. *Res. Microbiol.* **151**, 893–896 (2000).
- Liu, S. L., Hessel, A. & Sanderson, K. E. The *Xba*I-*Bln*I-*Ceu*I genomic cleavage map of *Salmonella typhimurium* LT2 determined by double digestion, end labelling, and pulsed-field gel electrophoresis. *J. Bacteriol.* **175**, 4104–4120 (1993).
- Liu, S. L. & Sanderson, K. E. Highly plastic chromosomal organization in *Salmonella typhi*. *Proc. Natl Acad. Sci. USA* **93**, 10303–10308 (1996).
- Ochman, H., Lawrence, J. G. & Groisman, E. A. Lateral gene transfer and the nature of bacterial innovation. *Nature* **405**, 299–304 (2000).
- Blanc-Potard, A. B., Solomon, F., Kayser, J. & Groisman, E. A. The SPI-3 pathogenicity island of *Salmonella enterica*. *J. Bacteriol.* **181**, 998–1004 (1999).
- Stanley, T. L., Ellermeier, C. D. & Schlauch, J. M. Tissue-specific gene expression identifies a gene in the lysogenic phage Gifsy-1 that affects *Salmonella enterica* serovar Typhimurium survival in Peyer's patches. *J. Bacteriol.* **182**, 4406–4413 (2000).
- Figuerola-Bossi, N., Uzzau, S., Maloroli, D. & Bossi, L. Variable assortment of prophages provides a transferable repertoire of pathogenic determinants in *Salmonella*. *Mol. Microbiol.* **39**, 260–272 (2001).
- Matsui, H. et al. Virulence plasmid borne *spvB* and *spvC* genes can replace the 90-kilobase plasmid in conferring virulence to *Salmonella enterica* serovar Typhimurium in subcutaneously inoculated mice. *J. Bacteriol.* **183**, 4652–4658 (2001).
- Ahmer, B. M., Tran, M. & Heffron, F. The virulence plasmid of *Salmonella typhimurium* is self-transmissible. *J. Bacteriol.* **181**, 1364–1368 (1999).
- Townsend, S. M. et al. *Salmonella enterica* serovar Typhi possesses a unique repertoire of fimbrial gene sequences. *Infect. Immun.* **69**, 2894–2901 (2001).
- Reidl, J. & Boos, W. The *malX* *malY* operon of *Escherichia coli* encodes a novel enzyme II of the phosphotransferase system recognizing glucose and maltose and an enzyme abolishing the endogenous induction of the maltose system. *J. Bacteriol.* **173**, 4862–4876 (1991).
- Decker, K., Gerhardt, F. & Boos, W. The role of the trehalose system in regulating the maltose regulon of *Escherichia coli*. *Mol. Microbiol.* **32**, 777–788 (1999).
- Gupta, S. D., Lee, B. T., Camakaris, J. & Wu, H. C. Identification of *cutC* and *cutF* (*nlpE*) genes involved in copper tolerance in *Escherichia coli*. *J. Bacteriol.* **177**, 4207–4215 (1995).
- Lan, R. & Reeves, P. R. Gene transfer is a major factor in bacterial evolution. *Mol. Biol. Evol.* **13**, 47–55 (1996).
- Bumann, D., Hueck, C., Aebischer, T. & Meyer, T. F. Recombinant live *Salmonella* spp. for human vaccination against heterologous pathogens. *FEMS Immunol. Med. Microbiol.* **27**, 357–364 (2000).

- Nakai, K. & Horton, P. PSORT: a program for detecting sorting signals in proteins and predicting their subcellular localization. *Trends Biochem. Sci.* **24**, 34–36 (1999).
- Pizza, M. et al. Identification of vaccine candidates against serogroup B meningococcus by whole-genome sequencing. *Science* **287**, 1816–1820 (2000).
- Tatusov, R. L. et al. The COG database: new developments in phylogenetic classification of proteins from complete genomes. *Nucleic Acids Res.* **29**, 22–28 (2001).

Supplementary information is available on Nature's World-Wide Web site (<http://www.nature.com>) or as paper copy from the London editorial office of Nature.

Acknowledgements

This work was supported by National Institutes of Health grants to R.W., M.M. and W.M. Nine ORFs were added to the annotation courtesy of L. Umayam and O. White of the Comprehensive Microbial Resource. We thank T. Lynch, M. Becker, G. Duckels and P. Minx for library production; J. Welsh, E. Koonin, R. Tatusov, H. Salgado, J. Collado-Vides and K. Nakai for assistance with computer analysis; S. Sims for assistance with primer design; M. Sekhon and the GSC Mapping Group for BAC mapping; S.-L. Liu for large restriction fragments of *S. typhimurium* LT2; A. McKay, B. Pearson, J. Roth and M. Riley for advice; and J. Parkhill and G. Dougan for sharing data from the *S. typhi* project.

Correspondence and requests for materials should be addressed to M.M. (e-mail: mmccllland@skcc.org). The sequence of the *S. typhimurium* LT2 chromosome and the annotated pSTL plasmid are deposited in GenBank under accession numbers AE006468 and AE006471, respectively.

Frizzled-7 signalling controls tissue separation during *Xenopus* gastrulation

Rudolf Winklbauer*†‡, Araceli Medina*†, Rajeeb K. Swain† & Herbert Steinbeisser*

† Max-Planck-Institute for Developmental Biology, Department of Cell Biology, Spemann Strasse 35, D-72076 Tübingen, Germany

* These authors contributed equally to this work

Cell signalling through Frizzled receptors has evolved to considerable complexity within the metazoans. The Frizzled-dependent signalling cascade comprises several branches, whose differential activation depends on specific Wnt ligands, Frizzled receptor isoforms and the cellular context. In *Xenopus laevis* embryos, the canonical β -catenin pathway contributes to the establishment of the dorsal–ventral axis¹. A different branch, referred to as the planar cell polarity pathway, is essential for cell polarization during elongation of the axial mesoderm by convergent extension². Here we demonstrate that a third branch of the cascade is independent of Dishevelled function and involves signalling through trimeric G proteins and protein kinase C (PKC). During gastrulation, Frizzled-7 (Fz7)-dependent PKC signalling controls cell-sorting behaviour in the mesoderm. Loss of zygotic Fz7 function results in the inability of involuted anterior mesoderm to separate from the ectoderm, which leads to severe gastrulation defects. This result provides a developmentally relevant *in vivo* function for the Fz/PKC pathway in vertebrates.

In *Xenopus*, maternal Fz7 acts in the Wnt/ β -catenin pathway and contributes to dorsalization of the embryo³. To study the role of zygotic *Xenopus* Fz7, we attempted a 'loss-of-function' approach that depends on blocking translation of the Fz7 messenger RNA by an antisense morpholino oligonucleotide directed against the 5'

‡ Present address: Department of Zoology, University of Toronto, 25 Harbord Street, Toronto, Ontario M5S 3G5, Canada.