Table 1 Appearance of hammerhead clones during selection

Round	Frequency*	Activity† (min ⁻¹)	No.‡	Progenitors§
5	0.02	0.0026 ± 0.000057	40	1/39
7	0.17	0.026 ± 0.0021	29	20/5
9	0.67	0.086 ± 0.0022	33	23/8
11	1.00	0.166 ± 0.0053	30	17/0
12	0.98	0.322 ± 0.014	55	22/1
13	—	0.655 ± 0.022	—	—
		0.0381 ± 0.0031		
14	0.93	0.835 ± 0.054	30	19/1
		0.050 ± 0.00098		
16	0.77	0.289 ± 0.031	30	7/2
		0.0649 ± 0.0069		

* Observed frequency of hammerhead-containing clones.

† Estimated pool activity (mean ± s.d.).

‡ Number of sequenced clones.

§ Number of distinct, independent sequences with/without hammerhead motif.

|| Activity in 0.5 mM MgCl₂ at pH 7.2.

efficiently by the RT–PCR procedure we used, or must be restricted to specific substrate sequences not included in our selection. Our results are consistent with those of Tang and Breaker⁷, who observed many distinct ribozymes with activities lower than 0.1 min⁻¹. In contrast, another selection experiment¹⁰, in which the random-sequence region contained only 18 nucleotides, yielded only hammerhead ribozymes at both low and high activities. However, the stringent size constraint on the random-sequence region, and the constraints placed on cleavage site recognition, may have ruled out the appearance of other ribozymes. There are several natural classes of self-cleaving ribozymes other than the hammerhead, including the hairpin ribozyme¹¹, the hepatitis delta ribozyme (HDV)^{12,13}, and the *Neurospora* VS motif¹⁴. We wondered why none of these alternative highly active ribozymes was observed in our selection. These ribozymes are all significantly more complex than the hammerhead ribozyme, and are thus expected to occur much less frequently in selection experiments, in the absence of selective factors other than rate of cleavage. Thus, if selective pressure requires only that self-cleavage activity be in the range of 0.1–1.0 min⁻¹, the expected result, both *in vitro* and in nature, is the emergence of the hammerhead ribozyme as the most probable solution. Our results show that, despite the dominance of contingency (historical accident) in some recent discussions of evolutionary mechanisms¹⁵, purely chemical constraints (that is, the ability of only certain sequences to carry out particular functions) can lead to the repeated evolution of the same macromolecular structures. □

Methods

Selection procedure

Pool RNA dissolved in 10 mM Tris–HCl buffer at pH 8.0 and 1 mM EDTA (TE) was heated to 85 °C then cooled to ambient temperature. Cleavage reactions were started by adding pool RNA to reaction buffer such that the final composition was 1 μM RNA, 5 mM MgCl₂, 50 mM Tris–HCl at pH 7.8, 10 mM DTT and 1 U μl⁻¹ RNasin. Reaction times were 17–20 h at 26 °C in rounds 1–5 and in the subsequent rounds were adjusted to produce 1% cleavage. Reactions were stopped by addition of two volumes of 8 M urea, 2 mM Tris–HCl at pH 7.5 and 75 mM EDTA. Samples were resolved by electrophoresis in 8% denaturing polyacrylamide gels alongside a marker RNA to identify the position of the cleaved products; this region was excised and the RNA was isolated. RNAs were reverse transcribed using SuperScript II (Gibco–BRL), amplified by PCR and transcribed with T7 RNA polymerase (0.5 mM nucleoside triphosphates, 40 mM Tris–HCl at pH 7.9, 6 mM MgCl₂, 2 mM spermidine, 10 mM dithiothreitol and 1 U μl⁻¹ RNasin) in the presence of 50 μM of the blocking oligonucleotide (5′-TACTGACTGGGTCCAG-3′). After transcription, RNAs were purified by denaturing polyacrylamide gels, eluted, ethanol precipitated, dissolved in TE, and stored at –80 °C.

Cleavage site determination

RNAs were denatured with heat, cooled to ambient temperature, incubated in cleavage buffer, and then reverse transcribed with a 5′-³²P-labelled primer (5′-CGTTGTGGTCTATC-3′). Samples mixed with formamide stop buffer were analysed by electrophoresis in 12% denaturing polyacrylamide gels.

Received 25 May; accepted 7 August 2001.

- Pabon-Pena, L. M., Zhang, Y. & Epstein, L. M. Newt satellite 2 transcripts self-cleave by using an extended hammerhead structure. *Mol. Cell. Biol.* **11**, 6109–6115 (1991).
- Zhang, Y. & Epstein, L. M. Cloning and characterization of extended hammerheads from a diverse set of caudate amphibians. *Gene* **172**, 183–190 (1996).
- Ferbyre, G., Smith, J. M. & Cedergren, R. Schistosoma satellite DNA encodes active hammerhead ribozymes. *Mol. Cell. Biol.* **18**, 3880–3888 (1998).
- Rojas, A. A. *et al.* Hammerhead-mediated processing of satellite pDo500 family transcripts from *Dolichopoda* cave crickets. *Nucleic Acids Res.* **28**, 4037–4043 (2000).
- Diener, T. O. Circular RNAs: relics of precellular evolution? *Proc. Natl Acad. Sci. USA* **86**, 9370–9374 (1989).
- Williams, K. P., Ciafre, S. & Tocchini-Valentini, G. P. Selection of novel Mg²⁺-dependent self-cleaving ribozymes. *EMBO J.* **14**, 4551–4557 (1995).
- Tang, J. & Breaker, R. R. Structural diversity of self-cleaving ribozymes. *Proc. Natl Acad. Sci. USA* **97**, 5784–5789 (2000).
- Thomson, J. B., Tuschl, T. & Eckstein, F. in *Catalytic RNA* (eds Eckstein, F. & Lilley, D. M. J.) 173–196 (Springer, Heidelberg, 1997).
- Ruffner, D. E., Stormo, G. D. & Uhlenbeck, O. C. Sequence requirements of the hammerhead RNA self-cleavage reaction. *Biochemistry* **29**, 10695–10702 (1990).
- Conaty, J., Hendry, P. & Lockett, T. Selected classes of minimised hammerhead ribozyme have very high cleavage rates at low Mg²⁺ concentration. *Nucleic Acids Res.* **27**, 2400–2407 (1999).
- Hampel, N. in *Progress in Nucleic Acids Research and Molecular Biology* (ed. Moldave, K.) 1–38 (Academic, London, 1998).
- Been, M. D. *Cis-* and *trans-*acting ribozymes from a human pathogen, hepatitis delta virus. *Trends Biochem. Sci.* **19**, 251–256 (1994).
- Thill, G., Vasseur, M. & Tanner, N. K. Structural and sequence elements required for the self-cleavage activity of the hepatitis delta virus ribozyme. *Biochemistry* **32**, 4254–4262 (1993).
- Beattie, T. L., Olive, J. E. & Collins, R. A. A secondary-structure model for the self-cleaving region of *Neurospora* VS RNA. *Proc. Natl Acad. Sci. USA* **92**, 4686–4690 (1995).
- Gould, S. J. *Wonderful Life: The Burgess Shale and the Nature of History* 292–323 (Norton, New York, 1989).
- Long, D. M. & Uhlenbeck, O. C. Self-cleaving catalytic RNA. *FASEB J.* **7**, 25–30 (1993).

Acknowledgements

We thank P. Svec for assistance in cloning and sequencing, and members of the laboratory for their comments on the manuscript. This work was supported in part by a grant from the National Institutes of Health. J.W.S. is an investigator of the Howard Hughes Medical Institute.

Correspondence and requests for materials should be addressed to J.W.S. (e-mail: szostak@molbio.mgh.harvard.edu).

corrections

Retrotransposition of a bacterial group II intron

Benoit Cousineau, Stacey Lawrence, Dorie Smith & Marlene Belfort

Nature **404**, 1018–1021 (2000).

In this Letter, we demonstrated active group II intron transposition to multiple ectopic sites and showed that movement to these new sites occurs via an RNA intermediate in a process termed retrotransposition. Of interest was whether retrotransposition occurs by reverse splicing of the intron RNA into RNA, single-stranded DNA and/or double-stranded DNA targets. Our results suggested that retrotransposition occurs predominantly by the intron reverse-splicing into an RNA target. However, subsequent experiments using a different intron donor have indicated that the selection method for retrotransposition events can strongly bias the interpretation of pathway (K. Ichyanagi, A. Beauregard and M.B., unpublished results). The new data do not point to RNA-targeting as the principal means for intron retrotransposition, and DNA-based pathways must be considered as playing an important role. □

pol IV (20 nM) or pol V (25–50 nM). The running-start nucleotide was absent in standing-start reactions. Reactions were run at 37 °C for 5 min and quenched by adding 20 µl of 20 mM EDTA in 95% formamide. Primer–template extension products were heat denatured and run on a 16% polyacrylamide denaturing gel. We measured integrated gel-band intensities by phosphorimaging using ImageQuant software (Molecular Dynamics). Nucleotide incorporation rates opposite both undamaged and lesion target sites were determined, and apparent V_{max}/K_m values and nucleotide misincorporation frequencies were calculated as described¹⁶. We carried out fidelity measurements using standing-start reactions on undamaged linear M13mp7 DNA templates annealed to 30mer ³²P-labelled primers with ATP (100 µM). The following template sequences were used: 3'...GTTGC CG...; 3'...CGTAGCC...; 3'TCGTAGC...; 3'TATCAAC, where the base in bold is the template target site at which the nucleotide misincorporation ratio, f_{inc} , was calculated.

Processivity measurements

We measured the processivity of pol IV and pol V Mut on a single-stranded circular DNA (M13 mp7) primed with a 5'-³²P-labelled 30-mer primer as described²⁹. A fixed amount of substrate (10 nM) was incubated with decreasing amounts of pol IV or pol V at 37 °C for 3 min. If indicated, other proteins included in the reactions were: RecA (5 µM), SSB (1.5 µM), β (200 nM) and γ-complex (50 nM). ATP (1 mM) is added in reactions containing RecA and/or γ-complex. Reactions were initiated by adding four dNTPs (100 µM), quenched after 5 min and analysed on a 12% denaturing polyacrylamide gel. Reaction products were quantified by phosphorimaging. Processivities were measured at polymerase concentrations, where product lengths remained stable, and the number of primers used remained less than 10%. We computed processivity as a ratio of the total number of nucleotides added in the reaction to the number of extended DNA primers. High enzyme/DNA ratios are required to carry out primer extension (Fig. 3), probably because the polymerases bind to the long regions of single-stranded DNA. We have previously observed cooperative binding of pol V to single-stranded DNA³⁰.

Received 13 December 1999; accepted 4 February 2000.

1. Tang, M. *et al.* Biochemical basis of SOS mutagenesis in *Escherichia coli*: reconstitution of *in vitro* lesion bypass dependent on the UmuD₂C mutagenic complex and RecA protein. *Proc. Natl Acad. Sci. USA* **95**, 9755–9760 (1998).
2. Tang, M. *et al.* UmuD₂C is an error-prone DNA polymerase, *Escherichia coli* pol V. *Proc. Natl Acad. Sci. USA* **95**, 8919–8924 (1999).
3. Wagner, J. *et al.* The *dinB* gene encodes a novel *Escherichia coli* DNA polymerase (DNA pol IV). *Mol. Cell* **40**, 281–286 (1999).
4. Friedberg, E. C., Walker, G. C. & Siede, W. *DNA Repair and Mutagenesis* 407–522 (ASM, Washington, 1995).
5. Kato, T. & Shinoura, Y. Isolation and characterization of mutants of *Escherichia coli* deficient in induction of mutagenesis by ultraviolet light. *Mol. Gen. Genet.* **156**, 121–131 (1977).
6. Bonner, C. A., Hays, S., McEntee, K. & Goodman, M. F. DNA polymerase II is encoded by the DNA damage-inducible *dinA* gene of *Escherichia coli*. *Proc. Natl Acad. Sci. USA* **87**, 7663–7667 (1990).
7. Brotcorne-Lannoye, A. & Maenhaut-Michel, G. Role of RecA protein in untargeted UV mutagenesis of bacteriophage λ: Evidence for the requirement for the *dinB* gene. *Proc. Natl Acad. Sci. USA* **83**, 3904–3908 (1986).
8. Kim, S.-R. *et al.* Multiple pathways for SOS-induced mutagenesis in *Escherichia coli*: An over-expression of *dinB/dinP* results in strongly enhancing mutagenesis in the absence of any exogenous treatment to damage DNA. *Proc. Natl Acad. Sci. USA* **94**, 13792–13797 (1997).
9. Knippers, R. DNA polymerase II. *Nature* **228**, 1050–1053 (1970).
10. Rangarajan, S., Woodgate, R. & Goodman, M. F. A phenotype for enigmatic DNA polymerase II: a pivotal role for pol II in replication restart in UV-irradiated *Escherichia coli*. *Proc. Natl Acad. Sci. USA* **96**, 9224–9229 (1999).
11. Lindahl, T. DNA repair enzymes. *Annu. Rev. Biochem.* **51**, 61–87 (1982).
12. Banerjee, S. K., Christensen, R. B., Lawrence, C. W. & LeClerc, J. E. Frequency and spectrum of mutations produced by a single *cis-syn* thymine–thymine cyclobutane dimer in a single-stranded vector. *Proc. Natl Acad. Sci. USA* **85**, 8141–8145 (1988).
13. Lawrence, C. W., Borden, A., Banerjee, S. K. & LeClerc, J. E. Mutation frequency and spectrum resulting from a single abasic site in a single-stranded vector. *Nucleic Acids Res.* **18**, 2153–2157 (1990).
14. LeClerc, J. E., Borden, A. & Lawrence, C. W. The thymine–thymine pyrimidine–pyrimidone (6–4) ultravioletlight photoproduct is highly mutagenic and specifically induces 3' thymine-to-cytosine transitions in *Escherichia coli*. *Proc. Natl Acad. Sci. USA* **88**, 9685–9689 (1991).
15. Rajagopalan, M. *et al.* Activity of the purified mutagenesis proteins UmuC, UmuD', and RecA in replicative bypass of an abasic DNA lesions by DNA polymerase III. *Proc. Natl Acad. Sci. USA* **89**, 10777–10781 (1992).
16. Creighton, S., Bloom, L. B. & Goodman, M. F. in *Methods Enzymol.* Vol. 262 (ed. Campbell, J. L.) 232–256 (Academic, San Diego, 1995).
17. Smith, C. A. *et al.* Mutation spectra of M13 vectors containing site-specific *cis-syn*, *trans-syn-I*, (6–4), and dewar pyrimidine photoproducts of thymidyl-(3'→5')-thymidine in *Escherichia coli* under SOS conditions. *Biochemistry* **35**, 4146–4154 (1996).
18. Lawrence, C. W., Banerjee, S. K., Borden, A. & LeClerc, J. E. T–T Cyclobutane dimers are misinstructive, rather than non-instructive, mutagenic lesions. *Mol. Gen. Genet.* **166**, 166–168 (1990).
19. Fijalkowska, I. J., Dunn, R. L. & Schaaper, R. M. Genetic requirements and mutational specificity of the *Escherichia coli* SOS mutator activity. *J. Bacteriol.* **179**, 7435–7445 (1997).
20. Bloom, L. B. *et al.* Fidelity of *Escherichia coli* DNA polymerase III holoenzyme: The effects of β, γ complex processivity proteins and ε proofreading exonuclease on nucleotide misincorporation efficiencies. *J. Biol. Chem.* **272**, 27919–27930 (1997).
21. Sommer, S., Boudsocq, F., Devoret, R. & Bailone, A. Specific RecA amino acid changes affect RecA–UmuD' interaction. *Mol. Microbiol.* **28**, 281–291 (1998).
22. Menetski, J. P. & Kowalczykowski, S. C. Interaction of RecA protein with single-stranded DNA: Quantitative aspects of binding affinity modulation by nucleotide cofactors. *J. Mol. Biol.* **181**, 281–295 (1985).
23. Reuven, N. B., Arad, G., Maor–Shoshani, A. & Livneh, Z. The mutagenic protein UmuC is a DNA

polymerase activated by UmuD', RecA, and SSB and is specialized for translesion replication. *J. Biol. Chem.* **274**, 31763–31766 (1999).

24. Radman, M. Enzymes of evolutionary change. *Nature* **401**, 866–869 (1999).
25. McDonald, J. P. *et al.* Novel human and mouse homologs of *Saccharomyces cerevisiae* DNA polymerase η. *Genomics* **60**, 20–30 (1999).
26. Gerlach, V. L. *et al.* Human and mouse homologs of *Escherichia coli* DinB (DNA polymerase IV), members of the UmuC/DinB superfamily. *Proc. Natl Acad. Sci. USA* **96**, 11922–11927 (1999).
27. Naktinis, V., Turner, J. & O'Donnell, M. A molecular switch in a replication machine defined by an internal competition for protein rings. *Cell* **84**, 137–145 (1996).
28. Jing, Y., Kao, J. F.-L. & Taylor, J.-S. Thermodynamic and base-pairing studies of matched and mismatched DNA dodecamer duplexes containing *cis-syn*, (6–4) and Dewar photoproducts of TT. *Nucleic Acids Res.* **26**, 3845–3853 (1998).
29. Bambara, R. A., Fay, P. J. & Mallaber, L. M. Methods of analyzing processivity. *Methods Enzymol.* **262**, 270–280 (1995).
30. Bruck, L., Woodgate, R., McEntee, K. & Goodman, M. F. Purification of a soluble UmuD' complex from *Escherichia coli*. Cooperative binding of UmuD' to single-stranded DNA. *J. Biol. Chem.* **271**, 10767–10774 (1996).

Acknowledgements

This work was supported by National Institutes of Health grants to M.F.G., M.O. and J.-S.T.).

Correspondence and requests for materials should be addressed to M.F.G. (e-mail: mgoodman@mizar.usc.edu).

Retrotransposition of a bacterial group II intron

Benoit Cousineau, Stacey Lawrence, Dorie Smith & Marlene Belfort

Molecular Genetics Program, Wadsworth Center, New York State Department of Health, and School of Public Health, State University of New York at Albany, PO Box 22002, Albany, New York 12201-2002, USA

Self-splicing group II introns may be the evolutionary progenitors of eukaryotic spliceosomal introns^{1–7}, but the route by which they invade new chromosomal sites is unknown. To address the mechanism by which group II introns are disseminated, we have studied the bacterial Ll.LtrB intron from *Lactococcus lactis*⁸. The protein product of this intron, LtrA, possesses maturase, reverse transcriptase and endonuclease enzymatic activities^{9–11}. Together with the intron, LtrA forms a ribonucleoprotein (RNP) complex which mediates a process known as retrohoming¹¹. In retrohoming, the intron reverse splices into a cognate intronless DNA site. Integration of a DNA copy of the intron is recombinase independent but requires all three activities of LtrA¹¹. Here we report the first experimental demonstration of a group II intron invading ectopic chromosomal sites, which occurs by a distinct retrotransposition mechanism. This retrotransposition process is endonuclease-independent and recombinase-dependent, and is likely to involve reverse splicing of the intron RNA into cellular RNA targets. These retrotranspositions suggest a mechanism by which spliceosomal introns may have become widely dispersed.

The group II intron donor in these experiments is a twintron/Kan^R variant of Ll.LtrB, carrying the self-splicing group I *td* intron and a kanamycin resistance (Kan^R) gene (Fig. 1a, b)¹¹. The Kan^R marker facilitates selection of group II intron mobility events. The group I intron allows identification of those mobility events that have proceeded through an RNA intermediate and in which the group I intron has therefore been lost. The donor was first tested for retrohoming using *L. lactis* and a two-plasmid system with a recipient plasmid containing the cognate Ll.LtrB homing site (Fig. 1a)¹¹. After RNP expression under the control of a nisin-inducible promoter¹², inheritance of the Kan^R-marked intron occurred at 40% or 58% per recipient in two different wild-type hosts (Fig. 1c, crosses 1 and 3, H%). Of these homing products, 98%

had lost the group I intron (Fig. 1c, crosses 1 and 3, GI loss %), indicating that retrohoming was at least 39% or 57% in the two *L. lactis* hosts (Fig. 1c, crosses 1 and 3, RH%). Thus, retrohoming of Ll.LtrB in *L. lactis* is extremely efficient, with a tenfold increase over uninduced levels¹¹ of the RNP, and a greater than 100-fold increase in Kan^R recombinants above a donor expressing non-functional LtrA (Fig. 1c, compare crosses 3 and 4). Retrohoming levels in an Endo⁻ mutant that retains reverse transcriptase and maturase activity

(J. San Filippo and A. M. Lambowitz, unpublished) dropped to 2% of the LtrA⁺ parent (Fig. 1c, compare crosses 3 and 5). This agrees with the behaviour of this mutant in *Escherichia coli* (5% of LtrA⁺, J. San Filippo and A. M. Lambowitz, unpublished). Furthermore, retrohoming levels remained high in $\Delta recA$ cells lacking an intact homologous recombination system, at 30% per recipient, or 74% of the efficiency in the *recA*⁺ wild-type host (Fig. 1c, compare crosses 1 and 2). These results are consistent with an endonuclease-mediated recombination-independent mechanism¹¹.

The donor plasmid for detecting group II intron movement to ectopic chromosomal sites has the nisin-inducible twintron/Kan^R expression cassette in an erythromycin-resistant (Erm^R) temperature-sensitive (Ts) shuttle vector. This plasmid is maintained in *L. lactis* at 30 °C but cannot replicate at 37 °C (Fig. 1b¹³). Therefore Kan^R transposition events into the chromosome were selected after increasing the temperature and then screened for plasmid curing (Erm^S) and the presence of both introns. Genomic DNA containing retrotransposed group II introns was then isolated from colonies that were Kan^R-Erm^S, group II intron positive and group I intron negative. After cloning the Kan^R genomic fragments, we determined the insertion sites of the intron by sequencing the flanking DNA.

We characterized eight different ectopic insertion sites of Ll.LtrB in the chromosome of *L. lactis* (Fig. 2, 1–8). Two of these sites were at different positions in 23S ribosomal RNA. Also, one was in the *hdsM* gene (methylase; Genbank AF034786), four were in open reading frames (ORFs) of unknown function and one was in undefined sequence (D. Ehrlich and A. Sorokin, personal communication). Remarkably, insertions 1–7 were all in the sense orientation of the gene.

Analysis of the sequences flanking retrotransposed introns indicates that the base pairings between the EBS1 sequence of the intron

Figure 1 Retrohoming and retrotransposition in *L. lactis*. **a**, Retrohoming assay. Ll.LtrB group II intron, red; group I *td* intron, grey; E1 and E2, Ll.LtrB exons; RNPs, LtrA protein and intron RNA; pNIS, nisin-inducible promoter. **b**, Retrotransposition assay. Recipient is bacterial chromosome. **c**, Retrohoming and retrotransposition levels. H, homing, $H = \text{Kan}^R\text{Cam}^R/\text{Cam}^R \times 100$ (ref. 11); GI loss, group I intron loss; RH, retrohoming, $RH = H \times \text{GI loss}$; T, transposition, RT, retrotransposition, $RT = T \times \text{GI loss}$; Rel WT = RH or RT relative to 100 for wild-type host (A or B) and intron donor (LtrA⁺). T and RT are expressed per total cell number (although relative values in different crosses are accurate, absolute values may be elevated by Kan^R enrichment). All assays were repeated three times concomitantly. Standard errors for H% are: cross 1, 5.92; cross 2, 3.60; cross 3, 5.22; cross 4, not calculated; cross 5, 0.13. Standard errors for $T \times 10^{-2}$ are: cross 1, 0.62; cross 2, 0.02; cross 3, 1.06; cross 4, 0.03; cross 5, 0.32.

Figure 2 Ectopic insertions of Ll.LtrB in the *L. lactis* chromosome. Top, Interaction between group II intron RNP and target DNA. Dotted lines, base pairings (EBS1/IBS1 and EBS2/IBS2) between intron RNA (red lariat) and DNA target; breaks in each DNA strand, cleavage sites. Three activities of LtrA protein: reverse transcriptase, RT; endonuclease, E; maturase, M. Bottom, wild-type retrohoming site (WT) and insertion sites of eight ectopic integrants (1–8). White nucleotides on black background represent identity with wild type, or the possibility of intron RNA interacting via G–U base pairs (asterisk). U, undefined sequence. Bidirectional arrows point at protein recognition sites G–21 and T+5.

and the IBS1 sequence 5' to the intron insertion site determine the specificity of LL.LtrB integration (Fig. 2). Conservation of nucleotides indicates that LL.LtrB may invade new sites by reverse splicing downstream of locations resembling its natural exon I sequence, and that the IBS1/EBS1 interaction is more important than the IBS2/EBS2 interaction, as in fungal systems^{14,15}. In accord with reverse splicing into the ectopic target sites, most of the retrotransposed introns were also able to undergo some degree of forward splicing, as determined by primer extension analysis of *L. lactis* RNA containing retrotransposed LL.LtrB. With an intron-specific primer (Fig. 3)¹⁰, complementary DNA corresponding to excised intron lariat, an indicator of splicing, was observed in most cases. Where the lariat is not detected there may be an absence of splicing, or simply low expression of the interrupted gene. The ectopic insertion of LL.LtrB near the 5' end of a 23S rRNA gene (Fig. 2, site 7) allowed us to estimate the splicing efficiency of LL.LtrB from this site, because the unspliced precursor was detectable. The ratio of spliced intron lariat to total RNA (precursor + lariat) was estimated at 0.1, compared with 0.8 for the intron at its native site.

Significantly, determinants of LtrA protein binding in the exons are not discernable. In retrohoming, protein recognition of the DNA target site is required at position -21(G) for reverse splicing and at position +5(T) for antisense-strand cleavage (Fig. 2, bottom)¹⁶. The lack of apparent protein recognition sites, particularly at T+5 in exon II, which is required for LtrA-mediated cleavage, suggests that retrotransposition might occur by an alternative pathway to retrohoming, a pathway where the LtrA endonuclease activity is dispensable.

To explore different pathways in ectopic intron mobility, we performed retrotransposition assays in the presence and absence of *L. lactis* RecA function and LtrA endonuclease activity. Whereas retrohoming proceeded at 74% of wild-type levels in *recA*⁻ cells, retrotransposition of LL.LtrB in the *recA*⁻ strain was reduced to 20% of that in a *recA*⁺ host (Fig. 1c, compare crosses 1 and 2, Rel WT), indicating that, unlike retrohoming, a significant proportion of retrotransposition events followed a recombination-dependent pathway. Furthermore, the LtrA endonuclease mutant, which strongly inhibited retrohoming (2% of the LtrA⁺ donor), was

greater than 20-fold more active in retrotransposition (41% of the LtrA⁺ donor; Fig. 1c, compare crosses 3 and 5).

Therefore only a fraction (20%) of retrotransposition events appear to occur via an illegitimate RecA-independent retrohoming mechanism (Fig. 4a). The retrohoming-like pathway, involving reverse splicing into double-stranded DNA (step 1a), cDNA synthesis (steps 2a and 3a), second-strand synthesis (step 4a) and ligation (step 5a), would be independent of homologous recombination, but dependent on the endonuclease activity of LtrA (step 1a)¹¹. This minor pathway would explain the endonuclease requirement of some retrotransposition events (Fig. 1c, compare crosses 3 and 5). However, the endonuclease-independence of 41% of retrotransposition events indicates that second-strand cleavage may not be required and that reverse splicing might occur into single-stranded nucleic acid, either single-stranded DNA (for example at a replication fork or an actively transcribed gene) or RNA. Whereas integration into DNA would probably involve the replication machinery but not recombination, integration into RNA is predicted to be RecA-dependent (Fig. 4b), as observed for most retrotransposition events. Reverse-splicing into RNA (step 1b) would require recombinase function, as for example after cDNA synthesis (steps 2b and 3b), where the double-stranded DNA undergoes a double crossover event (step 4b). Resolution (step 5b) would yield the retrotransposition product, with the intronless chromosomal allele being replaced by the intron-containing allele.

Interestingly, residual retrotransposition events in a *recA*⁻ background (20%; Fig. 1c, cross 2) that typify the DNA-based pathway

Figure 3 Splicing of LL.LtrB at ectopic sites. Primer extension was performed on total RNA¹⁰ from strain NZ9800ΔLL.LtrB containing ectopic chromosomal insertions (grown at 25 °C). Lengths of cDNA bands, synthesized with intron-specific primer (P), are indicated in nucleotides on the left. They correspond to precursor, excised intron lariat and an unknown *L. lactis* product (X) used as loading control. The positive control (+) has two independent transcription starts (black arrowheads)^{10,11}. 0, intronless *L. lactis* (NZ9800ΔLL.LtrB); lane numbers correspond to insertion sites in Fig. 2. Dashed lines, cDNAs; black bar, primer; upward-directed arrows, splice sites.

Figure 4 Mobility pathways. RNPs interact and promote reverse splicing of the intron into (a) dsDNA and (b) RNA. The RNP is depicted on a grey background with the intron lariat (red) and the trifunctional protein, LtrA, with reverse transcriptase (RT), endonuclease (E) and maturase (M) activities. Functions required exclusively by one pathway are shown in green. a, 1a, reverse splicing into dsDNA and endonuclease cleavage; 2a and 3a, cDNA synthesis; 4a, second-strand synthesis; 5a, repair. b, 1b, reverse splicing into RNA; 2b, cDNA synthesis; 3b, second-strand synthesis; 4b, recombination; 5b, resolution.

(Fig. 4a), plus those events with an Endo⁻ mutant (41%; Fig. 1c, cross 5) that are characteristic of the RNA-based pathway (Fig. 4b), fall short of 100%. This observation may be due to compromised reverse transcriptase and maturase activity in the endonuclease mutant, or to additional pathways to those shown in Fig. 4.

Retrotransposition into an RNA target is not only supported by the RecA-dependence and endonuclease-independence of many of the events, but is also consistent with the following observations. First, all the ectopic insertions are in the sense orientation of the respective genes, in accord with reverse splicing into expressed RNA (Fig. 2). Second, two of eight events are into a 23S rRNA gene. Ribosomal RNAs in a variety of organisms from all three kingdoms contain many introns at different positions, so these abundant RNAs provide many targets for reverse splicing and intron dissemination¹⁷. Third, autocatalytic introns are able to reverse-splice into RNA *in vitro* and *in vivo*^{17–23}. Regardless of pathway, the dispensability of the endonuclease activity of LtrA in retrotransposition reduces the specificity of site recognition by the RNPs to a small number of base pairs, as few as 8 out of 13 of the IBS1-IBS2 residues (Fig. 2), thereby promoting the promiscuity of group II introns and their ability to invade new sites.

The mechanism of DNA-targeted retrohoming and target-primed cDNA synthesis of group II introns (Fig. 4a) has led to comparisons to non-long-terminal-repeat (LTR) retrotransposons, which also encode proteins with reverse transcriptase and endonuclease activities^{24,25}. However, there are clear differences between retrotransposition of group II introns and non-LTR retrotransposons. First, there are striking mechanistic distinctions, with RNA being a possible target of a retrotransposing group II intron, and DNA being the exclusive target of a non-LTR retrotransposon. Second, reverse splicing of group II introns during retrotransposition guarantees a basic level of splicing at the newly invaded site (Fig. 3). Such events would be less deleterious to the host than gene invasion by non-LTR retrotransposons, thereby facilitating group II intron spread.

The discovery of group II introns in bacteria reinforced previous arguments for the antiquity of these catalytic RNA elements^{3,26}. The group II introns might have been pivotal in the evolution of eukaryotic genomes. On the one hand, the parallels in target-primed cDNA-synthesis and associated protein enzymes indicate that they may share evolutionary history with non-LTR retrotransposons^{24,25,27,28}, which occupy >17% of the human genome²⁹. On the other hand, because of similarities in splicing pathways and RNA components, group II introns may be ancestral to the spliceosome-dependent nuclear introns², which form >16% of the human genome (N. J. Schisler and J. Palmer, personal communication). Because of likeness in splicing chemistry, the mode of retrotransposition of group II introns in bacteria by reverse splicing into RNA may provide clues to the means of dispersal of spliceosomal introns. As an RNA-targeted mechanism is independent of endonuclease function, and because reverse splicing into RNA ensures forward splicing to maintain gene expression, this pathway is the most plausible mode of dissemination of spliceosomal introns in modern genomes of extant eukaryotes. □

Methods

Retrohoming assay

Spc^RKan^R products were selected from crosses between Cam^RKan^R donors (pLEIt^dKR⁺) and Spc^R recipients (pMN1343)¹¹. Cells were grown at 30 °C until the optical density at 600 nm was 0.5, then induced with nisin¹² for 3 h before plating under selective conditions. Assays were performed in *L. lactis* strain NZ9800 (wild-type host A) or NZ9800ΔLLtrB lacking the resident LLtrB intron (wild-type host B). Intron donors were wild-type (LtrA⁺), carried an LtrA deletion that removes all three activities (ΔLtrA, a >500-amino-acid deletion designated ΔORF¹⁰) or an LtrA mutant lacking endonuclease activity (Endo⁻ YRT mutant of J. San Filippo and A. M. Lambowitz, personal communication). Spc, spectinomycin; Cam, chloramphenicol.

Retrotransposition assay

The donor plasmid pGIIt^dKR⁺, a twintron/Kan^R derivative of temperature-sensitive pG⁺host5 (Erm^R)¹⁵, was transformed into *L. lactis*. Transformants were grown at 30 °C, and RNPs induced with nisin. After 3 h, cultures were diluted 100-fold into fresh medium and shifted to 37 °C for about 24 h to inhibit plasmid replication and stimulate plasmid loss. Retrotransposition events were enriched after 100-fold dilution in GM17-Kan medium (Kan^R) by overnight growth at 37 °C, and selected on kanamycin-containing plates. Kan^RErm^R colonies were analysed for group II intron acquisition and group I intron loss with ³²P-labelled region-specific dsDNA probes.

Received 2 December 1999; accepted 10 February 2000.

- Jacquier, A. Self-splicing group II and nuclear pre-mRNA introns: how similar are they? *Trends Biochem. Sci.* **15**, 351–354 (1990).
- Sharp, P. A. Five easy pieces. *Science* **254**, 663 (1991).
- Cavalier-Smith, T. Intron phylogeny: a new hypothesis. *Trends Genet.* **7**, 145–148 (1991).
- Roger, A. J. & Doolittle, W. F. Why introns-in-pieces? *Nature* **364**, 289–290 (1993).
- Weiner, A. M. mRNA splicing and autocatalytic introns: distant cousins or the products of chemical determinism? *Cell* **72**, 161–164 (1993).
- Michel, F. & Ferat, J. Structure and activities of group II introns. *Annu. Rev. Biochem.* **64**, 435–461 (1995).
- Hetzer, M., Wurzer, G., Schwyen, R. J. & Mueller, M. W. Trans-activation of group II intron splicing by nuclear U5 snRNA. *Nature* **386**, 417–420 (1997).
- Mills, D. A., Manias, D. A., McKay, L. L. & Dunny, G. M. Homing of a group II intron from *Lactococcus lactis* subsp. *lactis* ML3. *J. Bacteriol.* **179**, 6107–6111 (1997).
- Lambowitz, A. M., Caprara, M. G., Zimmerly, S. & Perlman, P. S. In *The RNA World* 451–485 (Cold Spring Harbor Laboratory Press, New York, 1999).
- Matsuura, M. *et al.* A bacterial group II intron encoding reverse transcriptase, maturase, and DNA endonuclease activities: biochemical demonstration of maturase activity and insertion of new genetic information within the intron. *Genes Dev.* **11**, 2910–2924 (1997).
- Cousineau, B. *et al.* Retrohoming of a bacterial group II intron: mobility via complete reverse splicing, independent of homologous DNA recombination. *Cell* **94**, 451–462 (1998).
- Kuipers, O. P., Beerthuyzen, M. M., de Ruyter, P. G., Luesink, E. J. & de Vos, W. M. Autoregulation of nisin biosynthesis in *Lactococcus lactis* by signal transduction. *J. Biol. Chem.* **270**, 27299–27304 (1995).
- Biswas, I., Gruss, A., Ehrlich, S. D. & Maguin, E. High-efficiency gene inactivation and replacement system for gram-positive bacteria. *J. Bacteriol.* **175**, 3628–3625 (1993).
- Mueller, M. W., Allmaier, M., Eskes, R. & Schwyen, R. J. Transposition of group II intron *all* in yeast and invasion of mitochondrial genes at new locations. *Nature* **366**, 174–176 (1993).
- Sellem, C. H., Lecellier, G. & Belcour, L. Transposition of a group II intron. *Nature* **366**, 176–178 (1993).
- Mohr, G., Smith, D., Belfort, M. & Lambowitz, A. M. Rules for DNA target site recognition by a *Lactococcus* intron enable retargeting of the intron to specific DNA sequences. *Genes Dev.* **14**, 559–573 (2000).
- Roman, J. & Woodson, S. A. Reverse splicing of the *Tetrahymena* IVS: Evidence for multiple reaction sites in the 23S tRNA. *RNA* **1**, 478–490 (1995).
- Woodson, S. A. & Cech, T. R. Reverse self-splicing of the *Tetrahymena* group I intron: Implication for the directionality of splicing and for intron transposition. *Cell* **57**, 335–345 (1989).
- Augustin, S., Muller, M. W. & Schwyen, R. J. Reverse self-splicing of group II intron RNAs *in vitro*. *Nature* **343**, 383–386 (1990).
- Mori, M. & Schmelzer, C. Integration of group II intron *bII* into a foreign RNA by reversal of the self-splicing reaction *in vitro*. *Cell* **60**, 629–636 (1990).
- Mohr, G. & Lambowitz, A. M. Integration of a group I intron into a ribosomal RNA sequence promoted by a tyrosyl-tRNA synthetase. *Nature* **354**, 164–167 (1991).
- Thompson, A. J. & Herrin, D. L. A chloroplast group I intron undergoes the first step of reverse splicing into host cytoplasmic 5.8 S rRNA. *J. Mol. Biol.* **236**, 455–468 (1994).
- Roman, J., Rubin, M. N. & Woodson, S. A. Sequence specificity of *in vivo* reverse splicing of the *Tetrahymena* group I intron. *RNA* **5**, 1–13 (1999).
- Eickbush, T. H. Mobile introns: retrohoming by complete reverse splicing. *Curr. Biol.* **9**, R11–R14 (1999).
- Curcio, M. J. & Belfort, M. Retrohoming: cDNA-mediated mobility of group II introns requires a catalytic RNA. *Cell* **84**, 9–12 (1996).
- Ferat, J. & Michel, F. Group II self-splicing introns in bacteria. *Nature* **364**, 358–361 (1993).
- Xiong, Y. & Eickbush, T. H. Origin and evolution of retroelements based upon their reverse transcriptase sequences. *EMBO J.* **9**, 3353–3362 (1990).
- Zimmerly, S., Guo, H., Perlman, P. S. & Lambowitz, A. M. Group II intron mobility occurs by target DNA-primed reverse transcription. *Cell* **82**, 545–554 (1995).
- Smit, A. F. The origin of interspersed repeats in the human genome. *Curr. Opin. Genet. Dev.* **6**, 743–748 (1996).

Acknowledgements

We thank J. San Filippo and A. Lambowitz for providing the LtrA Endo⁻ YRT mutant. We also thank D. Manias and G. Dunny for help in constructing pLEIt^dKR⁺; D. Ehrlich and A. Sorokin for information based on the *L. lactis* DNA sequence; J. Curcio, K. Derbyshire, V. Derbyshire, S. Hanes, A. Lambowitz, R. Lease, R. Morse, D. Nag and M. Parker for comments on the manuscript; N. J. Schisler and J. Palmer for estimates of intron composition of the human genome; and M. Carl for manuscript preparation. This work was supported by NIH grants to M.B.

Correspondence and requests for material should be addressed to M.B. (e-mail: belfort@wadsworth.org).