IN BRIEF

TECHNIQUE

Shedding light on disulfide bond formation: engineering a redox switch in green fluorescent protein.

Østergaard, H. *et al. EMBO J.* **20**, 5853–5862 (2001)

To study the formation of disulphide bonds in living cells, the authors have introduced two redox-active cysteines into yellow fluorescent protein (YFP). Under conditions in which this modified YFP forms a disulphide bond, its fluorescence is reduced twofold, allowing direct quantification of the redox state of the probe, and hence of its environment. The authors show that this YFP variant is indeed a sensitive probe of the redox state in *Escherichia coli* cytosol. This novel YFP probe should provide the tool to study disulphide bond formation in yeast and mammalian cells that has been missing so far

APOPTOSIS

Crystal structure of a procaspase-7 zymogen: mechanisms of activation and substrate binding.

Chai, J. *et al. Cell* **107**, 399–407 (2001)

Until now, all reported structural information on caspases has been derived from the inhibitor-bound forms of these enzymes or their procaspase precursors. But, reporting in *Cell*, Yigong Shi and colleagues describe the structures, at 2.6 and 2.7 Å respectively, of an active caspase and its inactive zymogen precursor — without any bound cofactors. The authors generated a non-cleavable procaspase-7 mutant and found that the structure of this differs considerably from that of a previously reported inhibitor-bound caspase-7 in the area surrounding the catalytic cleft. Binding by inhibitors causes the amino terminus of the small subunit to flip by 180°, allowing it to interact with — and hence stabilize — the catalytic cleft. These findings, say the authors, “provide a structural mechanism for caspase activation and substrate/inhibitor binding”.

DEVELOPMENT

Lsh, a member of the SNF2 family, is required for genome-wide methylation.

Dennis, K. *et al. Genes Dev.* **15**, 2940–2944 (2001)

In the mammalian genome, methylation of cytosine residues is known to be crucial for normal development, but the mechanisms that specify particular loci for methylation are not yet fully understood. Dennis *et al.* studied the effect of lymphoid-specific helicase (Lsh), a member of the SNF2 family of chromatin-remodelling proteins, on genome methylation patterns, and found that Lsh is essential for genome-wide methylation in mice. Despite being relatively normal in terms of their morphology and histology, *Lsh*^{−/−} mice were found to be hypomethylated at single-copy genes and repetitive sequences. The mice can survive embryogenesis, but die soon after birth. The results indicate that global DNA methylation is not essential for normal embryogenesis and that chromatin structure affects this methylation. The latter observation is particularly important, as previously only the opposite was thought to be true.

DNA REPAIR

Damage control

How a cell can detect DNA damage and recruit the correct cellular mechanics to repair faults has remained a mystery. But, reporting in *Science*, Katsunori Sugimoto and colleagues have shed more light on this process by showing how there might be two separate mechanisms involved in recognizing DNA damage.

In response to DNA damage, eukaryotic cells activate checkpoint pathways that arrest cell-cycle progression and induce the expression of genes that are required for DNA repair. This checkpoint machinery consists of proteins that recognize DNA damage and initiate the signalling response. Without this control mechanism, genomic instability can occur, which could lead to cancer.

In *Saccharomyces cerevisiae*, two essential genes, *MEC1* and *RAD53*, form the core of the checkpoint pathway. After DNA damage, Rad53 is phosphorylated and activated by a Mec1-dependent mechanism; and this phosphorylation is also dependent on other genes such as *RAD9*, *DDC1*, *MEC3*, *RAD17* and *RAD24*.

To detect the recruitment mechanism of these gene products, Sugimoto and colleagues created DNA lesions at a specific site and tested whether checkpoint proteins bound to that region. These lesions were generated by continuously expressing homothallic switching (HO) endonuclease, which creates a site-specific double-strand break (DSB) in the genome at the mating-type (*MAT*) locus.

The authors analysed the status of Rad53 phosphorylation in cells in which HO expression was driven from the galactose-inducible *GAL10* promoter. This confirmed that Rad53 is phosphorylated in response to DSB formation and that there is a lag between these two events.

Analysis of the Rad53 phosphorylation status in mutants showed this is dependent on *MEC1*, *DDC1*, *MEC3*, *RAD9*, *RAD17* and *RAD24*, and chromatin immunoprecipitation experiments revealed that the Ddc1 protein associates with a region near

the *MAT* locus after HO expression. Further mutant analysis showed that the Ddc1 association was dependent on Rad24 but not on Mec1 or Rad9.

But if this is the case, then what is the function of *MEC1*, which had been thought to be crucial to this process? Analysis of the interaction of Mec1 with the HO-induced DSB lesions in modified cells showed that Mec1 does bind to a region near the cleavage site after HO expression, but that this association is independent of Ddc1, Rad9 and Rad24.

So, the authors propose that Mec1 and Ddc1 are recruited independently to sites of DSBs, suggesting there might be two separate mechanisms involved in recognition of DNA damage. And as Mec1 acts upstream of Rad53, it seems clear that the checkpoint pathway is a well-defined signal transduction pathway. The authors also say that homologues of these genes have been found in other species, including humans, so it seems likely that this checkpoint control mechanism is conserved in all eukaryotic cells. But what remains to be identified is how DNA damage is recognized to initiate the checkpoint signalling.

Simon Frantz

References and links

ORIGINAL RESEARCH PAPER Kondo, T., Wakayama, T., Naiki, T., Matsumoto, K. & Sugimoto, K. Recruitment of Mec1 and Ddc1 to double-strand breaks through distinct mechanisms. *Science* **294**, 867–870 (2001)

FURTHER READING Melo, J. A., Cohen, J. & Toczyski, D. P. Two checkpoint complexes are independently recruited to sites of DNA damage *in vivo*. *Genes Dev.* **15**, 2809–2821 (2001)