10^{11} eV. It seems to us that, at least below 10^{14} eV, first-order Fermi acceleration may dominate, this process occurring in supernova remnants.

In conclusion, second-order Fermi acceleration may work at low energies ($<10^{11} - 10^{14}$ eV) and contribute a small fraction of the Galaxy cosmic rays there, but it requires unphysical values for the parameters of the Galaxy at much higher energies. The problem of the origin of ultra-high-energy cosmic rays therefore remains.

X. Chi

C. Dahanayake*

J. Wdowczyk

A. W. Wolfendale

Department of Physics,
University of Durham,
Durham DH1 3LE, UK

* Permanent address: University of Kelanya, Sri Lanka.

SIR — Bryant *et al.*² have given an alternative treatment of Fermi acceleration based on an analogy between collisions of cosmic rays with clouds and coin tosses. We show elsewhere⁶ how this treatment may be extended to include the bias for head-on versus overtaking collisions that is an essential feature in Fermi's original treatment. When the bias is included, the result of Bryant *et al.* reduces to a one-dimensional version of a result that is well known⁷⁻⁹.

The suggestion by Bryant *et al.* (see also the accompanying News and Views by Zank¹⁰) that 'high flyers' in Fermi acceleration can account for cosmic rays up to $\sim 10^{20}$ eV is inconsistent with known results for Fermi acceleration. In the conventional stochastic treatment of Fermi acceleration, monoenergetic particles with initial energy E_0 experience a systematic acceleration and a diffusion in energy. After N collisions, systematic acceleration leads to a mean energy $E = E_0 \exp(\frac{1}{2}\beta^2 N)$, where the factor $\frac{1}{2}$ is a feature of the one-dimensional treatment, and the diffusion causes a spread with half-width $\Delta E \approx E\beta N^{1/2}$ about this mean. The 'high flyers' correspond to the diffusive high-energy tail.

Bryant *et al.* drew an incorrect conclusion about the effectiveness of the acceleration, by which they clearly meant the time for acceleration: by integrating over N , which corresponds to evaluating the asymptotic spectrum by integrating over time, they lost all information on how long it takes to set up the asymptotic spectrum. In fact, after a fixed number of collisions, the spectrum of 'high flyers' falls off exponentially, faster than any power law, and so necessarily faster than the asymptotic power law spectrum. The characteristic time for the asymptotic power-law spectrum to be set up at a given energy is the time required for systematic acceleration to that energy. The 'high flyers' can play no

significant role independent of the systematic acceleration.

Bryant *et al.* provide new insight into Fermi acceleration, but they report no new physics and cannot resolve the long-standing difficulties in the application of the Fermi mechanism to cosmic rays.

Lewis Ball

Don Melrose

Colin Norman*

Research Centre for Theoretical

Astrophysics,

University of Sydney,

New South Wales 2006, Australia

* Permanent address: Department of Physics and Astronomy, Johns Hopkins University and Space Telescope Science Institute.

BRYANT *ET AL.* REPLY — In our initially purely mathematical re-evaluation of the capabilities of Fermi acceleration², the key parameters were all derived, using essentially only the observed spectral slope; none was freely adopted, as appears to have been assumed by Chi *et al.* Preliminary results of a separate exercise to refine the *prima facie* conclusions in the light of astrophysical measurements are reported here.

We omitted the usual statistical bias favouring head-on, or energy-gaining collisions, because a preliminary Monte Carlo numerical study had shown it not to be essential to the formation of a power-law distribution. We readily find from a formulation⁶ including a systematic term that the value of $N_0\beta^2$ required to give the observed spectral slope is $\frac{1}{3}$, as opposed to $\frac{1}{2}$ with no bias. This increases E_{lim} , but only by 7 per cent. Deductions based on 'a fixed number of collisions' (to quote Ball *et al.*) seem to be invalidated by the finite, well-defined probabilities² for any number of collisions and for any combination of energy gains and energy losses. As the maximum number of collisions is limited, in principle, only by the age of the system, there seems to be no fundamental reason why the asymptotic power-law spectrum should not be reached.

Let us now consider the possible constraints imposed on the derived optimal or critical magnetic fields, H_{crit} , by observed Faraday rotation of pulsar radiation. For an average Faraday rotation to correspond to B_0 , space would need to be filled with magnetic fields of this magnitude. So, as the filling factor is $(\%a)^3$, the average Faraday rotation, and apparent field, will be reduced by this factor. Random orientations within a distance scale $l \gg d$ will lead to a further reduction, which from simple considerations may be estimated as $\approx 1/\sqrt{3} (\%a)^{3/2}$. The apparent field on the l scale is thus $B_l \approx k (\%a/\sqrt{4\pi})^3 B_0$, where $k = 1/\sqrt{3}$ for $l \gg d$, and $k \rightarrow 1$ as $l \rightarrow d$. On the measurement scale of 50 pc ($1.5 \times$

10^{20} cm), the values of H_{crit} derived for $\beta = 10^{-1}$, 10^{-2} , 10^{-3} and 10^{-4} would appear as approximately 90, 6, 0.4 and 0.03 μG , respectively. (We have used here true values, rather than the rounded ones of our Table 1.) The last three are clearly permitted within the observational estimate of 5 μG (ref. 4). Further, the single-cell approximation used to estimate the field is considered to be too simplistic to give meaningful results, and the random field is actually more complex⁴, so even the $\beta = 10^{-1}$ model seems not too extreme. It may be prudent, though, for this case, to curtail the field strength by a factor of ten, and thus limit the maximum energy to 3×10^{19} eV. From the $\beta = 10^{-2}$ example, it is clear that proton energies up to 5×10^{19} eV are permitted by current knowledge of galactic magnetic fields. For heavy nuclei the limiting energy is higher, permitted values for iron, for example, extending up to $\sim 10^{21}$ eV.

To conclude, we respond individually to the numbered comments of Chi *et al.* (1) The energy densities of H_{crit} do, indeed, range from 3×10^4 eV cm^{-3} to 3×10^{-7} eV cm^{-3} , when the true (pre-rounding) filling factor of 3×10^{-3} is taken into account. If, as is implied, 1 eV cm^{-3} is taken as a criterion of acceptability, scattering-centre fields of ~ 100 μG would (with the above filling factor) be permitted. Taking up our earlier qualification² to the effect that, if the magnetic field strength is known to be less than H_{crit} , then the limiting energy is reduced in proportion, our four models yield maximum energies for protons of 4×10^{17} , 4×10^{16} , 4×10^{15} and 4×10^{14} eV. For iron nuclei these maxima become approximately 10^{19} , 10^{18} , 10^{17} and 10^{16} eV. These figures are encouragingly close to the highest observed values. (2) We have shown here that even the critical fields resulting automatically from the analysis (H_{crit}) are consistent with observation when the filling factor, $(\%a)^3$, rather than $\%a$, is used, and random orientation is allowed for. (3) The product of aH_{crit} is indeed, for the $\beta = 10^{-1}$ case, as quoted. But for fields of 100 μG , as already discussed, the product ranges from 3×10^{-4} mGpc to 0.3 mGpc for our four models. Such values seem to correspond, as far as we have been able to ascertain, to those of diffuse clouds and dark clouds. (4,5) The scattering centres referred to in our paper were not necessarily gas clouds, nor were they necessarily stable on cosmic timescales. It is possible that the inhomogeneities are, in fact, wavelike in nature, reflecting a degree of turbulence in the interstellar plasma. The scattering-centre treatment requires only that the inhomogeneities remain sensibly unchanged while being traversed by a cosmic ray. Stability timescales range,