

providing a strong positive feedback. As reported at the meeting, our model experiments⁸ can reproduce the extreme high-latitude temperature amplification of the reconstructions only by increasing ocean heat transport, a process by which sea ice becomes the prime positive feedback. The increased ocean heat transport might have come about through altered surface wind stress attributable to a change in topography, or through increases in deep-water production, perhaps associated with oceanic sill changes. Raymo noted that there is now some evidence that North Atlantic Deep Water production was greater during the late Pliocene⁹.

Where does this leave us? The participants generally agreed that although gradual uplift undoubtedly occurred in some regions, perhaps leading to increased erosion, the effects of these processes on climate are still debatable. We do not know what the CO₂ levels were in the Tertiary, nor whether higher CO₂ was in fact responsible for the warmth of those times. The question of the applicability of Tertiary climates as analogues for future trace gas-induced warming remains an open one. □

David Rind is at the Goddard Space Flight Center, Institute for Space Studies, New York, New York 10023, USA.

EVOLUTIONARY BIOLOGY

What the sperm count costs

Linda Partridge and Paul H. Harvey

DISSECTING the sex life of the nematode worm *Caenorhabditis elegans* has already provided surprises for biologists interested in life-history theory. In a report on page 456 of this issue¹, Van Voorhies throws another spanner in the works by demonstrating that the costs of producing sperm are not as negligible as we might have thought.

C. elegans individuals are normally self-fertilizing hermaphrodites, although the occasional male crops up from time to time. Last year's big surprise was that individual hermaphrodites usually produce more eggs than sperm, and that fecundity is limited by sperm production². Mutants that produced more sperm and fewer eggs left more offspring during their lifetime. The reason such mutants are selected against is that sperm are produced before eggs, so that the generation time for an individual that produces fewer sperm is shorter. In a growing population, which is where a reproducing *C. elegans* is likely to find itself, short generation time can be favoured over higher lifetime fecundity, and so the wild type can win against the more fecund mutants^{2,3}. Van Voorhies also uses mutants that are expected to change life-history schedules in predictable ways, and then sees if they do.

Sex differences

Biologists, who usually define males and females by the relative size of their gametes, believe that many other sex differences follow from this fundamental one. The argument generally runs along the following lines. If we assume that males and females have roughly the same amount of resources to devote to gamete production, then males will be able to produce more gametes than

females. Now make the further assumption that gametes are the only resources donated to the offspring, and it follows that female gametes will become a limiting resource for male reproduction. Males should compete to gain access for their sperm (or pollen) to eggs (or ovules)⁴. Such competition among males has led to the sexual selection of characters producing high male mating success, and hence the elaborate ornamentation and weaponry of the males of many species⁵.

Resources devoted to reproduction cannot be used for other vital processes, such as repair and maintenance, which are important for survival. Van Voorhies' experiments use decreases in rates of survival, or lifespan, to compare the costs of reproduction for both males and females. It is well established that such costs of reproduction can be important in practice⁶. If eggs are more expensive to produce than sperm, when a fixed number of eggs is added to a clutch we should expect to see a greater drop in survival for the female than for a male adding the same number of sperm. When hermaphrodites were mated to males (hermaphrodites will not mate with each other), the egg output went up two to threefold, yet lifespan in the mated hermaphrodites was no lower than that of unmated controls. In contrast, mating did reduce the lifespan of males and it is also known to increase their sperm production. Further evidence that the cost of mating was a consequence of making sperm, rather than the physical act of copulation, came from the finding that lifespan was increased relative to wildtype controls in both males and hermaphrodites that carried a mutant making them defective in spermatogenesis.

In the face of this evidence, the conclusion that each sperm costs more than each egg may seem hard to resist. However, males produce far more sperm (3,000+) than are needed to fertilize a hermaphrodite's eggs. Perhaps some of those sperm are metabolized into eggs after a hermaphrodite mates, thereby reducing the apparent cost of egg production. Nevertheless, high costs of spermatogenesis seem inescapable. Costs of reproduction for males are not new⁷, but the finding that spermatogenesis can so effectively curtail survival certainly is. On the basis of the data from the males alone, it could be argued that males are different from hermaphrodites in respects other than gamete production, and they also produce more sperm, and that is why males show such large reproductive costs.

Limiting factors

Sperm are qualitatively as well as quantitatively different from eggs, which may help explain why they are so costly to produce. Some support for this idea comes from a comparison of the rates of oogenesis versus spermatogenesis in *C. elegans*. Instead of the 1:500 ratio predicted by the size difference, sperm volume is produced at only 12.5 times the rate of egg volume. Sperm may contain a limiting nutrient that is present in larger quantities than in eggs, their synthesis may require more energy, or the rate of meiosis may be limiting. Whatever the explanation, *C. elegans* sperm are clearly limiting for female reproduction, contrary to the general expectation. Under such circumstances, hermaphrodites would be expected to compete for matings with males. Van Voorhies' unexplained finding that the act of mating itself appears to increase male lifespan would select further for a more than usually sex-mad male phenotype. Indeed, we are left wondering why selection has not succeeded in producing more males. □

Linda Partridge is in the Institute of Cell, Animal and Population Biology, Division of Biological Sciences, University of Edinburgh, West Mains Road, Edinburgh EH9 3JT, UK, and Paul H. Harvey is in the Department of Zoology, University of Oxford, South Parks Road, Oxford OX1 3PS, UK.

1. Van Voorhies, W. A. *Nature* **360**, 456–458 (1992).
2. Hodgkin, J. & Barnes, T. M. *Proc. R. Soc. B* **246**, 19–24 (1991).
3. Godfray, H. C. J. & Harvey, P. H. *Nature* **354**, 190–191 (1991).
4. Bateman, A. J. *Heredity* **2**, 349–368 (1946).
5. Trivers, R. L. in *Sexual Selection and the Descent of Man* (ed. Campbell, B.) 136–179 (Aldine, Chicago, 1972).
6. Bell, G. & Koufopanou, V. *Oxf. Surv. evol. Biol.* **3**, 83–131 (1986).
7. Partridge, L. & Farquhar, M. *Nature* **294**, 580–582 (1981).