Running out of control

Simon M. Hughes

THERE is a Chinese proverb that says: "When the dust settles you will see if you are riding a horse or an ass". The dust was in the air when mutations of *myoD* and *myf-5*, two genes thought to control muscle formation, gave rise to mice with surprisingly mild muscle deficits¹⁻³. But that was eight months ago. Now, two papers on pages 501 and 532 of this issue^{4,5} begin to clear the air. They show that disruption of the *myogenin* gene, which encodes the third, and in some ways the most exciting, member of the MyoD family of myogenic helix-loop-helix (HLH) DNA-binding proteins, yields mice that have little skeletal muscle. So, unlike *myoD* and *myf-5*, *myogenin* seems to be required for efficient muscle formation.

Myogenin is exciting because, like the rest of its family, it can convert other cell types to muscle and is expressed early in myogenesis *in vivo* but, uniquely in the family, there is a strong correlation between myogenin expression and myoblast differentiation. Muscle is formed by a multi-step process in which proliferating mononucleate myoblasts first arise from the somites; a proportion of these myoblasts then migrate to limb or ventral body-wall regions where they cease division and fuse with one another to form multinucleate muscle fibres. Analysis of limb muscle from one *myogenin* mutant line shows greatly reduced numbers of apparently normal fibres that express myosin surrounded by masses of mononucleate cells⁴. These mononucleate cells may be undifferentiated myoblasts because limb tissue from the other mutant line will generate large numbers of myogenic cells when put in culture⁵. So the mutants seem to have a deficiency not of myoblast generation, but of differentiation.

If myogenin functions in differentiation, how can one explain the sparse formation of fibres? Myogenin may not be required for differentiation of all subpopulations of myoblasts (and there is strong evidence for myoblast diversity from tissue culture work). Alternatively, it may simply increase the efficiency of myoblast differentiation. Why myoblasts without myogenin seem to differentiate better in a dish than in the animal is puzzling. The absence of myogenin in the few muscle fibres that form could cause those fibres to inhibit differentiation of the surrounding myoblasts, an inhibition that can occur in mature muscle⁶. Or

perhaps attention should focus on the ability of transcription factors like myogenin to prevent cells entering a new cell cycle, in the manner of tumour suppressor proteins. When cultured in the absence of growth factors, even *myogenin*-deficient cells may be forced to cease division, and then differentiation may follow by default. But *in vivo*, where cells may have access to a rich mixture of growth factors and may therefore be much closer to re-initiating a cell cycle, the lack of myogenin may be sufficient to tip the balance in favour

During G1 a myoblast decides between at least three possible routes. Contrary to the classic model of a bimodal switch between cell division and differentiation, a myoblast in the animal may also have the option to become a quiescent satellite cell, which is thought to be the inactive stem cell within muscle. A complete explanation of myoblast behaviour will take this trichotomy into account.

of proliferation.

Such a scheme fits both with the independent actions of myogenic HLH family members in cell division and differentiation^{7,8}, and with the finding⁹ that MyoD and myogenin binding to the unphosphorylated retinoblastoma gene product (pRB) may enhance both the tumour-suppressing abilities of pRB and the capacity of myogenin to activate muscle differentiation-specific genes. Moreover, the differentiation of myoblasts can be regulated by growth factor-dependent phosphorylation of myogenin¹⁰, an observation that strongly suggests that myogenin is involved in cell-cycle control. As the details of the mechanism by which the G1 phase of the cell cycle is regulated become clearer¹¹, it seems likely that muscle, through the myogenic HLH proteins, will be at the centre of studies of how cells make the decision between proliferation, quiescence and differentiation (see figure).

Although the *myogenin* mutant mice

generated in both studies^{4,5} do have some differentiated muscle fibres, the two papers diverge about the severity of the deficit in different muscles. Nabeshima *et al.*⁵ find more extensive muscle fibre formation in back and intercostal muscle (which may derive directly from the somites, where muscle differentiation occurs before myogenin expression) than in limb and ventral body-wall musculature (which derives from somites indirectly through a population of migratory precursor cells, that only initiate detectable myogenin expression as they differentiate)¹². Hasty *et al.*⁴ do not report these regional differences. The discrepancy, if confirmed, could arise either from differences in the genetic background of the two mouse strains or, more worryingly, from minor differences in the constructs, which would imply that one (or both) animals lack true null alleles.

The *myoD*, *myf-5* and *myogenin* mutant mice have several features in common. In each case myoblasts are formed and can differentiate into multinucleate muscle fibres, so none of the genes is absolutely required either for commitment to the muscle lineage or for terminal differentiation into post-mitotic contractile muscle. These animals provide a means of examining the interactions of myogenic HLH family members by breeding for double mutants and by analysing the behaviour of cultured cells that are missing particular genes. The finding that MyoD can induce fibroblasts from the *myogenin* mutant mice to differentiate as muscle⁵, is likely to be a first step to the dissection of the regulatory loops controlling myogenesis. It seems that as the dust settles around myogenin and its kin, we may find a race of thoroughbreds that will run and run. □

Simon M. Hughes is in the MRC Muscle and Cell Motility Unit and the Developmental Biology Research Centre, King's College London, 26-29 Drury Lane, London WC2B 5RL, UK.

1. Rudnicki, M. A., Braun, T., Hinuma, S. & Jaenisch, R. *Cell* **71**, 383-390 (1992).
2. Braun, T., Rudnicki, M. A., Arnold, H. H. & Jaenisch, R. *Cell* **71**, 369-382 (1992).
3. Hughes, S. M. *Nature* **360**, 536-537 (1992).
4. Hasty, P., Bradley, A., Morris, J. H., Edmondson, D. G., Venuti, J. M., Olson, E. N. & Klein, W. H. *Nature* **364**, 501-506 (1993).
5. Nabeshima, Y. *et al.* *Nature* **364**, 532-535 (1993).
6. Bischoff, R. *Development* **109**, 943-952 (1990).
7. Crescenzi, M., Fleming, T. P., Lassar, A. B., Weintraub, H. & Aaronson, S. A. *Proc. natn. Acad. Sci. U.S.A.* **87**, 8442-8446 (1990).
8. Sorrentino, V., Pepperkok, R., Davis, R. L., Ansorge, W. & Philipson, L. *Nature* **345**, 813-815 (1990).
9. Gu, W. *et al.* *Cell* **72**, 309-324 (1993).
10. Li, L. *et al.* *Cell* **71**, 1181-1194 (1992).
11. Sherr, C. J. *Cell* **73**, 1059-1065 (1993).
12. Cusella De Angelis, M. G. *et al.* *J. Cell Biol.* **116**, 1243-1255 (1992).