ting idea that these new alleles arose in response to selection pressures from disease encountered by Indian populations as they migrated southwards into the tropics. Disease resistance is also the most likely explanation for the significant excess of MHC heterozygotes among the Havasupai, a small tribe of about 600 people living in the splendid isolation of a side-gorge off the Grand Canyon (P. Hedrick, Arizona State University). Heterozygotes at the MHC seem to have a selective advantage over homozygotes because they can present more distinct peptide sequences to the immune system.

The identification of the HLA-B locus as particularly responsive to selection pressure is not confined to the Americas, or to people. Thus whereas the human class I HLA-A alleles A3/A1/A11 have close relatives in the common and pygmy chimpanzees *Pan troglodytes* and *P. paniscus*, there are considerable differences among class I HLA-B alleles in these species, particularly in the peptide-binding region.

What might be the mechanisms of production of new allelic variants in the human MHC? Gene conversion (exchanges of short segments of nucleotides between closely related alleles), it seems, is at least part of the answer. The human class I allele HLA-B*76, for example, found in Thais, has probably arisen by the substitution of just two amino acids at positions 166 and 167 (Parham). Most presumed conversions in MHC class I appear to involve the region coding for the peptide-binding site, but it is not known whether this is due to specificity in the conversion mechanism or to subsequent selection by disease.

Of the human class II alleles, diversification has occurred at all three of the main loci, DR, DP and DQ. For unknown reasons, different DP alleles seem to be of strikingly high frequency in different ethnic groups: the HLA-DP0501 allele, for example, is most prevalent in Japanese and south-east Asian communities (A. Begovitch, Roche, Alameda). So DP variations hold promise as ethnic or population-specific markers.

MHC alleles can be grouped according to sequence similarity into allelic lineages, which presumably reveal a common ancestry. Thus alleles of class II DRB1 and DRB6 genes are shared between humans, gorillas and chimpanzees. But there are new allelic variants in each species, showing that the DRB1 and DRB6 lineages are more than 25 million years old (R. Bontrop, University Hospital, Leiden). Furthermore, it seems that these molecules are functionally conserved in the peptide-binding region — antigen-presenting cells taken from rhesus macaques, for example, can present a peptide of the 65K heat-shock protein of *Mycobacterium leprae* to human T cells,

where both species share MHC DRB1*03 lineage molecules.

Going further back in time, there is the issue of whether New World monkeys (platyrrhines) and Old World simians (catarrhines) have a common ancestry or diverged independently from prosimian primates. As with humans, monkey MHC alleles generally differ at peptide-binding sites, but platyrrhine MHC genes have peculiar sequence 'motifs' in class II genes DRB1, 3 and 5 (W. Mayer, Max-Planck-Institut, Tübingen). It seems clear that these genes evolved more than 37 million years ago, probably before the divergence of platyrrhines and catarrhines (although fossil evidence tends to put divergences between primate groups even further back³).

Pseudogenes

Analysis of platyrrhine MHC genes (which are mostly non-functional pseudogenes) also reveals how new genes can be created from the ruins of the old (Mayer). The pseudogene DRB6, for example, is no longer active in platyrrhines, but is transcribed in catarrhines and humans. It seems that a functional promoter resembling that of mouse mammary tumour virus is responsible for gene expression, suggesting that a retroviral insertion occurred in the ancestral DRB6 gene more than 23 million years ago.

The advent of the prosimians lies even further back in time (more than 85 million years ago). Human-like class II DRB-1, 3 and 6 genes are found in extant examples of this group, such as bushbabies and lemurs, and phylogenetic trees show similarities and consistency with primates more generally (F. Figueroa, Max-Planck-Institut, Tübingen).

The persistence of alleles and allelic lineages through speciation events is not only evident in mammals. Analysis of a few of the hundreds of species of cichlid fishes in Africa's Lake Malawi and tributary rivers has shown that the founder population must have been highly polymorphic, and much of its allelic diversity has been passed down to the descendant species, all of which have arisen within the past two million years (Jan Klein, Max-Planck-Institut, Tübingen). So far, there is no evidence for the evolution of clusters of new species-specific alleles, such as have arisen in the class I HLA-B genes of the South American Indians. The inference to be made here is that the lake has provided a relatively stable environment, at least in terms of disease, compared to that encountered by migrating peoples⁴. □

Julie Clayton and Henry Gee are assistant editors at Nature.

1. Belich, M. P. *et al.* *Nature* **357**, 326–329 (1992).
2. Watkins, D. I. *et al.* *Nature* **357**, 329–333 (1992).
3. Martin, R. D. *Nature* **263**, 223–234 (1993).
4. Klein, D. *et al.* *Nature* **364**, 330–334 (1993).

Sparkling water

LAST week Daedalus decided that seals and dolphins were protected from the 'bends' by a bubble-inhibitor in their blood. He planned to identify and synthesize this material, and use it to load water with a vast amount of air which nonetheless could not bubble out. A swimming bath doped with DREADCO's 'Nobub' could hold so much oxygen that swimmers could breathe the water.

Daedalus now wants to try this trick on its obvious beneficiaries — fish. They are seriously restricted by the low solubility of oxygen in water. An aquarium supersaturated with oxygen and stabilized with Nobub would, for the first time, give them a copious oxygen supply. They should become much more active, and grow with wonderful rapidity. By breeding fish selectively in such an aquarium, DREADCO's biologists hope to produce strains adapted to exploit this novel environment: energetic, fast-growing and highly nutritious. The ultimate goal is a supersaturated fish farm with vastly improved yields.

The farm, of course, will generate a lot of excreta and general water pollution. Daedalus argues that this will be rapidly biodegraded in the oxygen-rich water. Similarly, he reckons that Nobub could help to clean up polluted rivers and lakes. A pressurization-unit could pump many atmospheres of oxygen into a Nobub-treated river, where it would accelerate natural purification. The most extreme biological oxygen demands would be effortlessly met. Industry and sewage and refuse-disposal services could go back to their old, cheap, simple policy of just dumping their wastes in the nearest river, while trusting Nobub and compressed air to clean up the mess. Nobub itself, having evolved to stabilize the blood-oxygen of aquatic mammals, will of course resist oxidation.

Small-scale uses for Nobub could also appear. Many housewives believe in the cleaning ability of detergent foam. DREADCO's Nobub detergent seems foam-free, despite being supersaturated with oxygen. But when a special Nobub-antagonist is added, the resulting foamy blast dislodges the most obdurate stains. Daedalus also began to devise a Nobub-doped beer, which would not foam crazily out of the freshly opened can or bottle. It would only seem fizz-free, however. As its content of Nobub was digested in the drinker's stomach, it would release its concealed gas, blasting him with almost terminal flatulence. So Daedalus now plans to counter fizzy beer by serving it in glasses lined with a Nobub-antagonist, to get rid of the wretched gas before it enters the drinker.

David Jones