higher visual centres in night vision; thus any two light stimuli causing equal rates of rhodopsin isomerization produce indistinguishable neural signals¹⁰. An explanation for the monogeusia analogous to that accounting for scotopic monochromacy is that the three sugars bind reversibly to a single class of membrane receptor in taste cells. On the assumptions that such binding is first-order, and that the psychophysical indiscriminability relation is transitive, our data imply that at the tongue

surface the apparent relative affinities for sucrose, fructose and glucose are 1, 2/3 and 1/3, respectively.

P. A. S. Breslin

S. Kemp*

G. K. Beauchamp

*Monell Chemical Senses Center,
Philadelphia,
Pennsylvania 19104,
USA*

*Present address: Givaudan-Roure, Teaneck, New Jersey 07666, USA.

Climate effects on mountain plants

SIR — Temperature-limited environments such as boreal regions, arctic regions and high mountains are thought to be very sensitive to greenhouse warming¹. As a result of such warming, for example, plant species and communities in high mountains will be pushed upwards in elevation and may be eliminated if already at mountain summits^{2,3}. If evidence were to be found of a continuing change of summit floras, this would indicate that global warming is already having a significant ecological impact. Although preliminary observations have recently been presented⁴, no conclusive evidence has

yet been reported.

During the summer of 1992, we collected data on the state of the flora at 26 summits exceeding 3,000 m in the middle part of the Alps (western Austria, eastern Switzerland) and compared the actual records on cover and abundance of vascular plant species with historical records^{5–10}. Species richness has increased during the past few decades (see figure), and is more pronounced at lower altitudes. However, upward movement of the alpine-nival flora is an overall trend, as indicated by the comparison of the fitted regression lines for species-richness with

altitude. Although the presented data are not corrected for age of the historical record or for the expansion of the considered belt, the difference between the two regression lines is obvious. The exponential decrease of species richness with altitude is a general feature of the nival vegetation above the closed alpine grasslands.

On the basis of 12 very precise historical records^{8–10}, we calculated moving rates for nine typical nival plant species. The maximum values approached 4 m per decade, although most of the values were below 1 m per decade. The old records used for this calculation were collected about 70–90 years ago. According to the records from meteorological stations in Austria, the mean annual temperature has increased since that time by 0.7 °C (ref. 11). Taking into account an average decrease of 0.5 °C per 100 m of increasing altitude, this warming should theoretically lead to a shift in the altitudinal vegetation belts at a rate of 8–10 m per decade. The empirical

values are clearly far lower than this.

In conclusion, there is no doubt that even moderate warming induces migration processes, and that this process is under way. The example from the limits of plant life at high alpine summits is of general importance and suggests that global warming is already having a significant effect on alpine plant ecology. Even in situations where plants must move upwards, the warming is sufficient to stimulate migration, and may cause disastrous extinctions in these environments.

Georg Grabherr, Michael Gottfried

Harald Paul

Department of Vegetation

Ecology and Conservation Biology,

University of Vienna, PO Box 285,

A1091 Vienna, Austria

1. Markham, S., Dudley, N. & Stoltson, S. *Some Like it Hot* (WWF International, Gland, 1993).
2. Holten, J. I. in *Impacts of Climate Change on Natural Ecosystems* (eds Holten, J. I. et al.) 84–105 (Norwegian Institute for Nature Research, Trondheim, 1993).
3. Nilsson, S. & Pitt, D. *Mountain World in Danger* (Earthscan, London, 1991).
4. Hofer, H. R. *Ber. Geobot. Inst. ETH, Stiftung Rübel* **58**, 39–54 (1992).
5. Klebelsberg, R. *Das Vordringen der Hochgebirgsvegetation in den Tiroler Alpen*. *Österr. Bot. Zeitschr. Jahrg.* 177–186; 241–254 (1913).
6. Rübel, E. *Pflanzengeographische Monographie des Berninagebietes*. *Bot. Jahrbuch Bd.* 47/1–4 (1911).
7. Reisinger, H. & Pitschmann, H. *Obere Grenzen von Flora und Vegetation in der Nivalstufe der Zentralen Ötztaler Alpen (Tirol)*. *Vegetatio*, **VIII**, 93–129 (1958).
8. Braun-Blanquet, J. *Ein Jahrhundert Florenwandel am Piz Linard (3414 m)* (Bull. Jard. Bot., Bruxelles, 1957).
9. Braun, J. *Die Vegetationsverhältnisse in der Schneestufe der Rätisch-Lepontischen Alpen* **48**, 156–307 (Neue Denkschr. Schweiz. Naturforsch. Ges., 1913).
10. Braun-Blanquet, J. *Über die obersten Grenzen Pflanzlichen Lebens im Gipfelbereich des Schweizerischen Nationalparks* **6**, 119–142 (Kommission der Schweizerischen Naturforsch. Ges. zur Wiss. Erforsch. des Nationalparks, 1958).
11. Auer, R. et al. in *Bestandsaufnahme anthropogene Klimaänderungen mögliche Auswirkungen auf Österreich — mögliche Massnahmen Österreich* (Österr. Akademie Wissenschaft, 1992).

Species richness of historical and our present-day records at nival summits in the Alps plotted against altitude. Rare species are downweighted (0.25), species of moderate abundance were given the weight 0.5, frequent species the weight 1.0. Rare species were downweighted due to the comparatively high probability that these were overlooked by the original authors. Displayed are 24 summits with siliceous bedrock. Age of the historical records is 40 to 90 years (circles, 1895–1918; squares, 1947–53; triangles, present-day). Arrows, increase in species richness, which is pronounced at lower altitudes and low to moderate at higher altitudes. Most summits considered are climbed only occasionally, so the effect of humans is negligible.

Scaling and brain connectivity

SIR — From published information describing the connections between areas of macaque monkey visual cortex, Young has applied non-metric multidimensional scaling (NMDS)^{1–4} to construct a two-dimensional map of 30 areas⁵. Areas are represented as points and were found to be arranged on a circle, which Young took as evidence for the 'two-streams' hypothesis of visual processing.

NMDS is designed for ordinal data. From a matrix of similarities between a given set of objects, a representation of the objects in a geometric space is sought such that the rank order of the distances between objects in the space corresponds to the reverse rank order of the similarities. In Young's case, similarity encodes the degree of connectivity for each pair of areas, with the connectivity specified as either reciprocal, unidirectional, or absent. We first explored the fit of his