The morphological and histological evidence presented here demonstrates that the remarkable mandibular filter-feeding apparatus of *Pterodaustro* is composed of true dental structures. Elaborate filter-feeding mechanisms are rare among adult tetrapods. The filter-feeding apparatus of *Pterodaustro* is not comparable to that of baleen whales⁶ or filter-feeding birds⁷, which are equipped with non-dental, epidermally derived sieving structures. Modified teeth for filtering are known for mammals (such as the crab-eater seal, *Lobodon carcinophagus*⁸), the Permian mesosaurs⁹, and other pterosaurs (such as *Ctenochasma* and *Gnathosaurus*³), although none of these shows the extraordinary specialization of *Pterodaustro*.

L. M. Chiappe

Department of Ornithology,
American Museum of Natural History,
New York, New York 10024, USA

A. Chinsamy

Earth Sciences Division,
South African Museum, PO Box 61,
Cape Town 8000, South Africa

Social stress and dominance

SIR—It has recently been shown that social dominance can carry subtle costs^{1–5}. Although it is generally thought that social stress falls more heavily on subordinates than on dominants, few studies have examined how stress relates to rank in the wild. We have investigated the levels of glucocorticoid stress hormones in two carnivores that live in complex societies, African wild dogs (*Lycaon pictus*) and dwarf mongooses (*Helogale parvula*). For these species, chronic stress is apparently a cost of social dominance rather than of subordination.

The link between social subordination and elevated glucocorticoids has primarily been shown in captivity. Although wild subordinate baboons have chronically elevated glucocorticoid levels⁶, data from other free-living species do not confirm that social subordinates experience greater stress than dominants^{7–9}. In the wild, subordinates can often avoid aggressive interactions, while dominants initiate aggression to reinforce their social status. Consequently, dominant dwarf mongooses and wild dogs engage in aggressive interactions more often than do subordinates, particularly during mating periods (wild dog females: $t_{33}=2.30$ (Student's *t*-test), $P=0.015$; wild dog males: $t_{78}=2.81$, $P=0.003$; dwarf mongooses, see ref. 10). High rates of aggression may be stressful, regardless of whether the encounters are won. If so, dominants would suffer greater social stress than subordinates.

We measured the urinary cortisol levels

c.v. was 3.4%. The inter-assay c.v. was 18.4%. HPLC showed a single peak of immunoreactivity in mongoose urine, which coeluted with free cortisol. Error bars show 1 s.e.

of dwarf mongooses (Serengeti National Park, Tanzania; 97 females, 82 males, 740 samples) and the faecal corticosterone levels of wild dogs (Selous Game Reserve, Tanzania; 22 females, 34 males, 216 samples). In both studies, samples were collected noninvasively from known individuals whose social rank was determined from aggressive interactions¹⁰. For African wild dogs, dominants had higher basal corticosterone levels than subordinates (*a* in the figure; partial $F_{1,207}=6.76$, $P=0.01$, controlling for gender and reproductive condition). Because we collected samples from unhandled animals, we have no data on corticosteroid responses to acute stress in wild dogs. For female dwarf mongooses, basal cortisol levels were higher in dominants than in subordinates (*b* in the figure; $t_{99}=6.37$, $P<0.001$). Nonetheless, dominant females produced higher cortisol levels than subordinates in response to the stress of trapping (*b* in the figure: $t_{74}=3.06$, $P=0.003$), indicating that their acute stress response was not compromised by chronically elevated cortisol. In male mongooses, basal cortisol levels did not correlate with rank, but dominant males showed low peak cortisol levels in response to trap-stress ($t_{91}=3.01$, $P=0.003$), indicating that their acute stress response was compromised.

Because chronically elevated glucocorticoids can inhibit reproduction^{6,7}, it is intriguing that dominants are typically the only breeders within packs of both species (for example, dwarf mongooses, ref. 11; wild dogs, S.C. and N.M.C., unpublished data). Dominant wild dogs and dwarf mongooses apparently experience greater social stress than do subordinates. This result is predicted if aggression itself is stressful, because dominant dwarf

Glucocorticoid levels in African wild dogs and dwarf mongooses. *a*, Dominant African wild dogs (blue bars) show higher basal corticosterone levels than subordinates (red bars) ($F=6.76$; $P=0.01$). Corticosterone values are expressed as ng per g dry faeces, measured with a double-antibody ¹²⁵I-corticosterone radioimmunoassay (ICN Biomedicals). Assay sensitivity was 25 ng corticosterone per ml. The inter-assay coefficient of variation (c.v.) was 8.0% for a high-corticosterone control and 13.7% for a low-corticosterone control; the intra-assay c.v. was <10% for both controls. HPLC showed a single peak of corticosterone immunoreactivity in wild dog faeces, with polarity intermediate between cortisol and corticosterone. *b*, Dominant female dwarf mongooses show higher basal cortisol levels than subordinates ($t=6.37$; $P<0.001$) and higher peak cortisol levels in response to stress ($t=3.06$; $P=0.003$). The effect of rank on cortisol levels remains significant after controlling for the effect of diurnal variation. Reproductive condition (mating/non-mating) did not affect cortisol levels. Cortisol levels are expressed as ng of hormone per mg of urinary creatinine to control for urine concentration. Urinary cortisol was assayed at 1:250 dilution (0.5 ml) using a ³H-cortisol RIA kit (RSL Immuchem). The intra-assay

mongooses and wild dogs often assert their dominance through aggression. In contrast to our findings, dominant baboons rarely assert their rank⁴, and social stress falls on subordinates.

Social stress may vary with social conditions, so that dominants are stressed (rather than subordinates) if they are highly aggressive. In captivity, stress may fall on subordinates because they cannot avoid aggressive dominants. Broader data are needed to resolve the differences among species (including humans) and social contexts. Nonetheless, our findings show that social stress can be a cost that offsets the benefits of dominance.

Scott Creel

Nancy Marusha Creel

Field Research Center for Ecology
and Ethology,
Rockefeller University,
Millbrook, New York 12545, USA

Steven L. Monfort

Department of Reproductive Physiology,
Conservation and Research Center,
National Zoological Park,
Front Royal, Virginia 22630, USA

1. Packer, C. et al. *Nature* **373**, 60–63 (1995).
2. Wasser, S. *Nature* **376**, 219–220 (1995).
3. Packer, C. *Nature* **377**, 689–690 (1995).
4. Altmann, J. et al. *Nature* **377**, 688–689 (1995).
5. Frank, L. G. et al. *Nature* **377**, 584–585 (1995).
6. Sapolsky, R. M. *Endocrinology* **116**, 2273–2278 (1985).
7. Wingfield, J. C. et al. *J. Zool., Lond.* **225**, 43–58 (1991).
8. Schoech, R. L. et al. *Condor* **93**, 354–364 (1991).
9. Mays, N. A. et al. *Auk* **108**, 619–637 (1991).
10. Creel, S. R. et al. *Anim. Behav.* **43**, 231–245 (1992).
11. Keane, B. et al. *Anim. Behav.* **47**, 65–75 (1994).

Scientific Correspondence

Scientific Correspondence is intended to provide a forum in which readers may raise points of a scientific character. Priority will be given to letters of fewer than 500 words.