homeodomain is sufficient for the binding to *cad* mRNA; other regions of the protein, or other homeodomain proteins such as *cad*, orthodenticle and the homeodomain of antennapedia, did not bind. That the *bcd* homeodomain interacts with *cad* mRNA *in vitro* is consistent with the *in vivo* observation, suggesting that the homeodomain of *bcd* is necessary for the formation of the *cad* gradient (Fig. 1f,g). The failure of other homeodomain proteins to bind *cad* mRNA does not rule out the possibility that they possess RNA-binding properties as preceded by zinc-finger-type transcription factors¹⁰.

To assess the functional significance of the *in vitro* interaction between *bcd* and *cad* mRNA, we performed co-transfection experiments with *Drosophila* Schneider cells¹¹. We used chloramphenicol acetyl transferase (CAT) reporter gene constructs¹¹ containing different 3' UTR sequences (Fig. 3). The reporter gene constructs were set under the control of the *bcd*-responsive, *cis*-acting element of the zygotic *hunchback* promoter¹². After co-transfection with *bcd*, we determined the stimulated transcription of CAT mRNA by reverse transcription-polymerase chain reaction (RT-PCR)¹³, and the translation of the CAT mRNAs with the different 3' UTRs by assaying CAT activity¹¹.

CAT mRNAs containing the *Drosophila cad* 3' UTR sequences or the regions including the BBR produced about sevenfold less CAT activity than CAT mRNA containing the *cad* 3' UTR sequences of *Clogmia* or the 3' half of the 3' UTR of *Drosophila cad* (Fig. 3a-c). CAT mRNA containing the BBR within the simian virus 40 (SV40) 3' UTR¹¹ were also translated less efficiently than those containing only the SV40 3' UTR. These results suggest that *bcd* may not only act as a transcriptional activator in the assay system, but also reduce the translation efficiency of the BBR-containing mRNAs.

To show that *bcd* does indeed suppress the translation of BBR-containing mRNA, we examined reporter gene constructs containing the SV40 3' UTR with or without the BBR that were transcriptionally activated by the yeast GAL4 transcriptional activator¹⁴. In the absence of *bcd*, the GAL4-stimulated CAT mRNAs were translated with similar efficiency (Fig. 4). In the presence of *bcd*, however, the translation of the BBR-containing CAT mRNA was significantly reduced, but the translation of CAT mRNA without the BBR was not significantly affected. This shows that *bcd* suppresses the translation of BBR-containing mRNA.

Our results are consistent with the argument that the anterior determinant *bcd* acts not only as a transcriptional activator but also functions in a manner analogous to the key components of the posterior maternal pattern-organizer system¹⁵⁻²⁰, which acts by translational repression through the activities of *nanos* and *pumilio*¹⁵⁻²⁰. After binding the *pumilio* protein and a 55K protein to the *nanos*-response element within the 3' UTR of *hunchback* mRNA^{17,20}, the *nanos* protein is thought to interact with this 'landing pad' to provide region-specific translational repression of the evenly distributed maternal *hunchback* mRNA¹⁵⁻¹⁹. This allows for the activation of the posterior gap genes *knirps* and *giant* in response to *cad*⁵. Thus, while in the posterior region of the embryo translational and transcriptional control are conducted through separate components, the anterior determinant *bcd* can exert both regulatory functions. Our findings also imply that the asymmetric distribution of the key components of the anterior and posterior maternal pattern-forming systems, which was thought to be set up independently^{1,2}, is linked through the dual function of *bcd*. □

Received 5 January; accepted 26 January 1996.

1. St Johnston, R. D. & Nüsslein-Volhard, C. *Cell* **68**, 201-219 (1992).
2. Pankratz, M. J. & Jäckle, H. in *The Development of Drosophila melanogaster* (eds Bate, M. & Martinez Arias, A.) 467-516 (Cold Spring Harbor Laboratory Press, NY, 1993).
3. Driever, W. & Nüsslein-Volhard, C. *Cell* **54**, 83-93 (1988).
4. Struhl, G., Struhl, K. & Macdonald, P. M. *Cell* **57**, 1259-1263 (1989).
5. Rivera-Pomar, R., Lu, X., Perrimon, N., Taubert, H. & Jäckle, H. *Nature* **376**, 253-256 (1995).
6. Mlodzik, M., Fjose, A. & Gehring, W. J. *EMBO J.* **4**, 2961-2969 (1985).
7. Macdonald, P. & Struhl, G. *Nature* **324**, 537-545 (1986).
8. Mlodzik, M. & Gehring, W. J. *Development* **101**, 421-435 (1987).
9. Mlodzik, M. & Gehring, W. J. *Cell* **48**, 465-478 (1987).
10. Pieler, T. & Theunissen, O. *Trends biochem. Sci.* **18**, 226-230 (1993).
11. Sauer, F. & Jäckle, H. *Nature* **353**, 563-566 (1991).
12. Driever, W. & Nüsslein-Volhard, C. *Nature* **337**, 138-143 (1989).

13. Ruden, D. & Jäckle, H. *Development* **121**, 63-73 (1995).
14. Sauer, F. & Jäckle, H. M. *Nature* **364**, 454-457 (1993).
15. Lehmann, R. & Nüsslein-Volhard, C. *Nature* **329**, 167-170 (1987).
16. Tautz, D. *Nature* **332**, 281-284 (1988).
17. Wharton, R. & Struhl, G. *Cell* **67**, 955-967 (1991).
18. Struhl, G. *Nature* **338**, 741-744 (1989).
19. Hülskamp, M., Schröder, C., Pfeifle, C., Jäckle, H. & Tautz, D. *Nature* **338**, 629-632 (1988).
20. Murata, Y. & Wharton, R. *Cell* **80**, 747-756 (1995).
21. Berlieth, T. et al. *EMBO J.* **7**, 1749-1756 (1988).
22. Driever, W. thesis, Eberhard-Karls-Univ. Tübingen (1989).
23. Simpson-Brose, M., Treisman, J. & Desplan, C. *Cell* **78**, 855-865 (1994).
24. Tautz, D. & Pfeifle, C. *Chromosoma* **98**, 81-85 (1989).
25. Rebagliati, M. *Cell* **58**, 231-232 (1989).
26. Müller, M. et al. *EMBO J.* **7**, 4299-4304 (1988).
27. Hess, M. A. & Duncan, R. J. *biol. Chem.* **269**, 10913-10922 (1994).
28. Soeller, W., Poole, S. & Kornberg, T. *Genes Dev.* **2**, 68-81 (1988).
29. Kerrigan, L. A., Croston, G. E., Lira, L. M. & Kadonaga, J. T. *J. biol. Chem.* **266**, 574-582 (1991).

ACKNOWLEDGEMENTS. We thank W. Driever for fly stocks; G. Dowe, M. Gemkow, T. Häder, C. Hartmann, G. Humbert Lan and M. Pankratz for their contributions; and G. Struhl for discussion on the roles of *bcd* and *cad*. This work was supported by the Max Planck Gesellschaft and the Fonds der chemischen Industrie (H.J.).

ADDENDUM

Identification of the breast cancer susceptibility gene *BRCA2*

Richard Wooster, Graham Bignell, Jonathan Lancaster, Sally Swift, Sheila Seal, Jonathan Mangion, Nadine Collins, Simon Gregory, Curtis Gumbs, Gos Michlem, Rita Barfoot, Rifat Hamoudi, Sandeep Patel, Catherine Rice, Patrick Biggs, Yasmin Hashim, Amanda Smith, Frances Connor, Adelgeir Arason, Julius Gudmundsson, David Ficenc, David Kelsell, Deborah Ford, Patricia Tonin, D. Timothy Bishop, Nigel K. Spurr, Bruce A. J. Ponder, Rosalind Eeles, Julian Peto, Peter Devilee, Cees Cornelisse, Henry Lynch, Steven Narod, Gilbert Lenoir, Valdgardur Egilsson, Rosa Bjork Barkadottir, Douglas F. Easton, David R. Bentley, P. Andrew Futreal, Alan Ashworth & Michael R. Stratton

Nature **378**, 789-792 (1995)

THE sequence for the *BRCA2* gene is now available at the web page of the Institute of Cancer Research, address <http://www.icr.ac.uk/molcarc/brca2.htm>, as well as from Genbank. □

ERRATUM

Mechanosensory signalling in *C. elegans* mediated by the GLR-1 glutamate receptor

Andres V. Maricq, Erin Peckol, Monica Driscoll & Cornelia I. Bargmann

Nature **378**, 78-81 (1995)

FIGURES 3 and 4 of this Letter were inadvertently transposed during the production process; the legend to Fig. 3 therefore describes the figure published as Fig. 4 and vice versa. □