for this activity, and named it agrin. Agrins are made by motor neurons and transported to the periphery where they become highly concentrated in the basal lamina at the neuromuscular junction. Crucially, for the hypothesis that agrin mediates nerve-induced AChR aggregation¹, they showed that the formation of AChR clusters during neuromuscular junction formation *in vitro* is prevented by agrin-binding antibodies⁵.

One way of testing the importance of agrin *in vivo* is to knock out the gene concerned. Gautam *et al.*¹ have now done this, with striking results. Mice lacking agrin are immobile when born and die within hours. Although AChRs are present on the surface of the muscle fibre, they are not aggregated into clusters at sites of nerve-muscle contact. Although motor nerve fibres branch exuberantly within the muscles, they do not form recognizable neuromuscular junctions. Furthermore, the muscle cell nuclei underlying the nerve branches, which normally preferentially express the genes that encode the AChR, fail to do so in the absence of agrin. These studies establish the essential role of agrin in synapse formation and add support to the view that its effects go beyond those on AChR aggregation⁶.

If agrin is an essential signal for neuromuscular junction formation, what mediates the muscle's response to it? Considerable excitement accompanied the finding that α -dystroglycan, part of a complex that links dystrophin to the cell surface, is a major agrin-binding protein. But it has since been shown that the region of the agrin molecule by which it binds to α -dystroglycan is not essential for AChR aggregating activity and that isoforms of agrin made by muscle cells bind well to α -dystroglycan but do not cause AChR aggregation⁷. So although α -dystroglycan binds agrin, it seems that it is not the receptor that mediates AChR clustering.

It has been known for some time that agrin activity requires protein phosphorylation⁸, suggesting that agrin activates a protein kinase. Last year, Valenzuela *et al.*⁹ identified a receptor tyrosine kinase specific for skeletal muscle, which they named MuSK and which has some particularly intriguing properties. Like AChRs, MuSK is uniformly distributed over the surface of immature and inactive muscle fibres, and becomes restricted to the region of the neuromuscular junction during development. Now the same group² reports that mice in which the MuSK gene has been knocked out have a phenotype that is virtually identical to that of agrin-deficient mice and a strikingly similar and equally comprehensive impairment of the development of neuromuscular junctions. Testing the notion that MuSK might itself be the agrin receptor, Glass *et al.*³ found that AChRs on MuSK-deficient muscle

cells grown in culture did not aggregate in response to agrin.

Receptor tyrosine kinases respond to the binding of extracellular signalling molecules such as growth factors by phosphorylating proteins, which may include themselves. Fragments of agrin that are adequate to cause AChR aggregation also cause rapid autophosphorylation of MuSK, whereas a number of other growth factors do not³. However, agrin does not bind directly to the extracellular domain of MuSK on its own, either *in vitro* or when MuSK expression is induced in non-muscle cells. On the other hand, expression of MuSK in immature myotubes induces agrin binding to the cell surface. This suggests that, as with a number of other receptor tyrosine kinases, agrin binds to a complex of MuSK with at least one other component which is present specifically on muscle cells.

It has been a long haul to these genetic knockout studies, and it is highly satisfying that the results are now in: they provide strong support for McMahan's 'agrin hypothesis'⁴, and do much to establish that agrin is an essential signal for AChR aggregation at the neuromuscular junction. Although the evidence that MuSK is the receptor that mediates these effects of agrin is not yet unequivocal, it is clear that it plays an essential part in the muscle's response to agrin. Thus signal-activated protein phosphorylation, a widespread component of growth control systems, also plays a key part in synapse formation.

In addition, these papers and another in press¹⁰ imply that the consequences of agrin signalling extend beyond AChR aggregation to other aspects of the development of neuromuscular junctions, including the control of nerve growth and AChR gene expression. Because agrin and/or its gene is expressed in both cholinergic and non-cholinergic neurons in the central nervous system¹¹, it is likely that there, too, agrin provides essential signals for synapse formation. □

Clarke R. Slater, of the Department of Neurobiology, University of Newcastle upon Tyne, is in the Muscular Dystrophy Group Research Laboratories, Newcastle General Hospital, Westgate Road, Newcastle upon Tyne NE4 6BE, UK.

- Gautam, M. *et al.* *Cell* **85**, 525–535 (1996).
- DeChiara, T. M. *et al.* *Cell* **85**, 501–512 (1996).
- Glass, D. J. *et al.* *Cell* **85**, 513–523 (1996).
- McMahan, U. J. *Cold Spring Harb. Symp. quant. Biol.* **60**, 407–418 (1990).
- Reist, N. E., Werle, M. J. & McMahan, U. J. *Neuron* **8**, 865–868 (1992).
- Cohen, I. *et al.* *Soc. Neurosci. Abstr.* **21**, 1748 (1995).
- Meier, T., Gesemann, M., Cavalli, V., Ruegg, M. A. & Wallace, B. G. *EMBO J.* (in the press).
- Wallace, B. G. *J. Cell Biol.* **125**, 661–668 (1994).
- Valenzuela, D. M. *et al.* *Neuron* **15**, 573–584 (1995).
- Jones, G. *et al.* *Proc. natn. Acad. Sci. U.S.A.* (in the press).
- O'Connor, L. T., Lauterborn, J. C., Gall, C. M. & Smith, M. A. *J. Neurosci.* **14**, 1141–1152 (1994).

Electric fire

ALONE among vehicles, the rocket has to carry its entire energy supply — oxygen as well as fuel. In this connection Daedalus recalls the wire-guided missile, steered by commands sent to it along an unreeling cable. In principle, it could get energy from the cable as well. But how to power a rocket electrically? The plasma torch attains a temperature far beyond that of combustion by means of an electric arc struck across its flame. Daedalus reckons that a rocket might similarly double the energy of its fuel by striking a big plasma arc across its exhaust jet. Sadly, the required heavy current would need such thick unreeling cables that they would soon drag the vehicle back to Earth.

But Daedalus does not give up so easily. The plasma torch is hot enough to ionize many of its gas molecules. So he is designing a rocket whose exhaust plume is itself an electrical conductor. For maximum conductivity, he will dope the fuel with a readily ionized element such as caesium. The vehicle will need two separate and widely spaced motors, one for the outward current and one for the return, but this is a minor complication. There will be two flat electrodes on the launch pad, one under each motor, and both connected to a power station. When the motors fire, the circuit will be completed, and a huge current will pass up one plume and down the other. The rocket will roar into the sky, trailing two conducting plumes that carry energy to its motors. The current will heat the whole length of both plumes, keeping them hot and conducting, but wasting a lot of energy in the process. Even so, the heat will be most intense at the point of minimum cross-section — in the rocket nozzles themselves, just where it is needed. In any case, electricity is very cheap compared to rocket fuels. The system could be quite economic.

How high could the rocket climb before its energizing current chokes off? A fluid conductor is compressed and compacted by the magnetic 'pinch effect'. Daedalus argues that this will oppose the diffusive spreading of the plumes, and will keep them compact, hot and conducting for many kilometres into the sky. (On the same principle the two plumes, carrying current in opposite directions, will repel each other, thus preventing a short-circuit between them.) A rocket burns most of its fuel in the early part of its climb; by the time its current is cut off, it will have gained a powerful boost. Furthermore, thrust adjustments and control signals for the tricky early stages of flight could be sent up the conducting plumes. Space launchers will take a new great leap upwards.

David Jones