- L1 consensus primers and RFLP analysis. *Mol Cell Probes* 1992;6:145–52.
52. Jacobs MV, de Roda Husman AM, van den Brule AJC, Snijders PJF, Meijer CJLM, Walboomers JMM. Group-specific differentiation between high and low-risk human papillomavirus genotypes by general primer-mediated PCR and two cocktails of oligonucleotide probes. *J Clin Microbiol* 1996;33:901–5.
 53. Jacobs MV, Snijders PJF, van den Brule AJC, Helmerhorst TJM, Meijer CJLM, Walboomers JMM. A general primer GP5+/GP6+ mediated PCR enzyme immunoassay method for rapid detection of 14 high-risk and 6 low-risk human papillomavirus genotypes in cervical scrapings. *J Clin Microbiol* 1997;35:791–5.
 54. Matsukura T, Sugase M. Molecular cloning of a novel human papillomavirus (type 58) from an invasive cervical carcinoma. *Virology* 1990;177:833–6.
 55. Gorgoulis V, Giatromanolaki A, Karameris A, Tsatsanis C, Aninos D, Ozanne B, *et al.* Epidermal growth factor receptor expression in squamous cell lung carcinomas: an immunohistochemical and gene analysis in formalin-fixed, paraffin-embedded material. *Virchows Arch A Pathol Anat* 1993;423:295–302.
 56. Herrington CS, McGee JO. Diagnostic molecular pathology. A practical approach. Vol. 2, IRL. Oxford: Oxford University Press; 1992. p. 87.
 57. Herrington CS, Burns J, Graham AK, Bhatt B, McGee JO'D. Interphase cytogenetics using biotin and digoxigenin labelled probes: I. Relative sensitivity of both reporter molecules for the detection of HPV 16 in CaSki cells. *J Clin Pathol* 1989;42:591–600.
 58. Park JS, Namkoong SE, Lee HY, Kim SJ, Daniel RW, Shah KV. Detection of human papillomavirus genotypes in cervical neoplasia from Korean women using polymerase chain reaction. *Gynecol Oncol* 1991;41:129–34.
 59. Fouret P, Monceaux G, Temam S, Lacourreye L, Saint-Guily JL. Human papillomavirus in head and neck squamous cell carcinomas in nonsmokers. *Arch Otolaryngol Head Neck Surg* 1997;123:513–6.
 60. Cooper K, Herrington CS, Graham AK, Evans MF, McGee JO. In situ evidence for HPV 16, 18, 33 integration in cervical squamous cell cancer in Britain and South Africa. *J Clin Pathol* 1991;44:406–9.
 61. Liu X, Han S, Baluda MA, Park NH. HPV-16 oncogenes E6 and E7 are mutagenic in normal human oral keratinocytes. *Oncogene* 1997;14:2347–53.

Book Review

Rapley R, editor: *The Nucleic Acid Protocols Handbook*, 1050 pp, Totowa, NJ, Humana Press, 2000 (\$110).

This book contains 120 standard protocols for analyzing nucleic acids. It is divided into 10 parts, including nucleic acid extraction, probe design and synthesis, gene library construction and screening, DNA sequencing, polymerase chain reaction, and techniques for inducing and identifying mutations and for localizing genes. Almost all chapters contain references to Maniatis's 1989 spiral-bound volumes, indicating that the present book is in essence a practice-tested version of that classical laboratory manual. Most protocols refer to widely used techniques, the description of which could be found, in one form or another, in many other manuals or research

papers. The advantage to the potential user is that the techniques are presented in minute details and in a "cookbook-standardized manner." That most contributors are from England or Europe should not discourage anyone because the themes covered are universal. The personal notes inserted by each contributor provide assurance that the techniques would work in any laboratory, as long as the instructions are followed carefully. Having all of those protocols updated and tested in practice and bound in one volume will be attractive to many scientists who deal with molecular biology techniques.

Hazem M. Hallak
*Thomas Jefferson University
 Philadelphia, Pennsylvania*