Editor's note

Is there a role of hypoxemia in penile fibrosis

Pro: Robert B. Moreland Ph.D, *Department of Urology, Boston University School of Medicine*

Con: Jacob Rajfer, M.D., *Department of Urology, UCLA School of Medicine*

In Dr Moreland's very nice presentation and manuscript he presented data showing the effects of hypoxia on human penile smooth muscle cells in tissue culture. The possibility of similar effect *in vivo* in the penis is a theoretical one and based on the assumption that in the man without severe vascular disease the penile corpora suffer from extended hypoxia.

In the contrary view Dr Rajfer suggested that there is no evidence in the absence of severe vascular disease that hypoxia in the penis actually exists. There is no evidence that at low penile blood flow the penile smooth muscle and endothelial cells do not receive sufficient oxygenation to maintain normal metabolic processes. Therefore, the artificial hypoxic environment created in the laboratory does not exist in normal man who may have erectile impairment.

A Melman
Editor

Is there a role of hypoxemia in penile fibrosis: a viewpoint presented to the Society for the Study of Impotence

RB Moreland

Department of Urology, Boston University School of Medicine, Boston, Massachusetts 02118, USA

During erection, oxygen tension changes in the corpus cavernosum penis from 25–40 mm Hg in the flaccid state to 90–100 mm Hg in the erect state.¹ The relationship between corpus cavernosum trabecular structure and erectile function is dependent on a critical balance of smooth muscle to connective tissue for successful veno-occlusion.² In this article, the potential role for transforming growth factor β_1 (TGF- β_1) and prostaglandin E (PGE) in maintaining a functional smooth muscle/connective tissue balance are discussed as well as the importance of oxygen tension in the synthesis of these factors. Correlations between animal models of disease as well as clinical reports are presented in support of a role for hypoxemia in penile fibrosis. A case is presented for a biological basis of nocturnal penile tumescence in the preservation of potency and an overall hypothesis for the molecular pathology of erectile dysfunction is proposed.

Keywords: collagen; cAMP; PGE₁; TGF- β_1 ; vascular smooth muscle; corpus cavernosum; erectile dysfunction; hypoxemia

Introduction

There has been considerable discussion regarding the role of oxygen tension in the modulation of male

erectile function at the level of smooth muscle tone and on connective tissue metabolism. One of the issues that remains to be resolved is whether hypoxemia is associated with penile fibrosis. At flaccidity, the corpus cavernosum trabecular smooth muscle is contracted and the penile blood PO₂ is 25–40 mm Hg.¹ During erection, the trabecular smooth muscle relaxes, allowing an inflow of arterial blood, and the blood PO₂ increases, reaching

Correspondence: Dr RB Moreland, Room W607A, 700 Albany Street, Boston, MA 02118, USA. Email: moreland@bu.edu

a plateau at 90–100 mm Hg.¹ This increase in blood PO₂ precedes the rise in cavernosal pressure which represents functional veno-occlusion as the subtunical draining venules are stretched and occluded. It is our hypothesis that this difference in oxygen tension in the flaccid and erect states results in the regulation of synthesis of cytokines, autoids, growth and vasoactive factors, which play a major role in connective tissue metabolism (Figure 1). Thus, in flaccidity, reduced oxygen tension leads to trabecular connective tissue synthesis and during erection, increased oxygen tension leads to reduced connective tissue synthesis and collagen degradation. Sexual as well as nonsexual (for example, nocturnal penile tumescence) erections may be important in modulating trabecular connective tissue metabolism and to maintain a functional connective tissue/smooth muscle ratio² crucial for erectile function.

The association of hypoxemia with erectile dysfunction would imply that the epidemiology of this condition would be correlated with vascular disease. Vasculogenic erectile dysfunction is primarily associated with vascular insufficiency to the penis.^{3,4} This condition is a complication of vascular disease resulting from diabetes mellitus, cigarette smoking, hypertension, or hypercholesterolemia and atherosclerosis. It may also result from crural⁵ or pudendal⁶ trauma which damages the penile arterial supply.³ Therefore, vascular disease and hence vascular insufficiency are consistent with corpus cavernosum hypoxemia.

If erectile dysfunction is a result of trabecular structural changes, one would expect to find common pathologic characteristics. The histopathology of penile biopsy specimens from patients with erectile

Figure 1 The seesaw of erectile (dys)function. At flaccidity, corpus cavernosal blood PO₂ is low (25–40 mm Hg¹), the trabecular smooth muscle of the lacunar spaces contracted. It is our hypothesis that these conditions favor the production of growth factors, cytokines and vasoactive factors which favor connective tissue synthesis. During erection, corpus cavernosal blood PO₂ increases and plateaus at 90–100 mm Hg¹ and the trabecular smooth muscle of the lacunar spaces relaxes. We further hypothesize that these conditions favor connective tissue degradation. It is the sum of both processes which lead to the trabecular smooth muscle/connective tissue balance.

dysfunction include a qualitative linkage between arterial insufficiency, corpus cavernosum smooth muscle cell atrophy and excess connective tissue (collagen) accumulation.^{7–11} In a prospective study of men undergoing penile prosthesis insertion, we have found a direct correlation between erectile function and the ratio of trabecular smooth muscle to total erectile tissue.² Erectile function was defined by experimentally determined parameters of flow to maintain erection, resistance to filling, and pressure decay using dynamic infusion corpus cavernosography/cavernosometry (DICC)¹² before penile implant insertion. Postoperatively, penile biopsies were examined by quantitative histomorphometry staining with desmin (a smooth muscle selective marker) immunohistochemistry or Masson's trichrome staining. Biopsies from potent men, penile cancer, crural trauma, and Peyronie's disease patients desiring implants rather than Nesbitt's procedure had a range of smooth muscle from 40–45%; consistent with accepted 'normal' values of percent smooth muscle. Patients with the highest mean percentage of trabecular smooth muscle (41–43%) exhibited a narrow range of normal flow-to-maintain values (5 ml/min), consistent with good veno-occlusion. Patients with intermediate values of mean percentage of trabecular smooth muscle (32–40%) displayed intermediate values of abnormal flow-to-maintain (15–30 ml/min), and some venous leakage while patients with the lowest mean percentage of trabecular smooth muscle (13–30%) revealed a wide range of high abnormal flow-to-maintain values (50–120 ml/min) and substantial cavernous leakage. Similar relationships were found when outflow resistance and pressure decay were correlated with percent trabecular smooth muscle. Therefore, flow to maintain erection increases due to venous leakage; resistance falls as veno-occlusion become less successful; and pressure decay, a measure of both inflow and outflow increases with decreased veno-occlusion. These results suggest a critical balance of trabecular smooth muscle and connective tissue is necessary for successful veno-occlusion and that excessive diffuse connective tissue is correlated with a higher likelihood of venous leakage and erectile failure.² A defined understanding of the pathophysiology of erectile dysfunction requires a knowledge of the molecular and biochemical regulation of connective tissue synthesis and degradation in the corpus cavernosum.

It is well accepted that the trabecular smooth muscle cell, the predominant mesenchymal cell type in the corpus cavernosum, regulates vasoconstriction and vasodilation, thus providing the state of erection of the corpora.¹³ What is less well known is that vascular smooth muscle cells synthesize collagen and connective tissue proteins.¹⁴ Human corpus cavernosum smooth muscle cells in culture express type I, III, and V/XI fibrillar collagens.¹⁴ Type IV collagen plays an important role in basement membrane assembly and has been reported in

corpus cavernosum biopsies at the smooth muscle-endothelial cell boundary of the trabeculae as well as in between smooth muscle cells.¹¹ Collagen accumulation can be modulated by degradative enzymes (for example collagenases and gelatinases) secreted by the surrounding smooth muscle cells. The matrix metalloproteinases are a family of secreted zinc proteinases which degrade collagen and gelatin (denatured collagen).¹⁵ These proteins are expressed as zymogens and activated by proteolysis. The normal processes of remodeling and deposition of extracellular matrix as well as events leading to excess matrix accumulation remain to be fully characterized in the corpus cavernosum.

In this article, a series of critical questions will be addressed pertaining to the possible role of hypoxemia in penile fibrosis. In particular, there will be a focus on the role of transforming growth factor β_1 expression induced by hypoxemia, TGF- β_1 induction of collagen synthesis in corpus cavernosum, the inhibition of prostaglandin E synthesis in corpus cavernosum by hypoxemia, role of PGE in the suppression of TGF- β_1 -induced collagen synthesis in corpus cavernosum, and whether or not hypogonadism is the same as hypoxemia.

What molecular factors regulate the processes of collagen synthesis and degradation in the corpus cavernosum?

Initial studies from our laboratory have chosen to focus on two factors: transforming growth factor- β_1 (TGF- β_1) as a 'fibrogenic cytokine' and prostaglandin E as an autacoid which can inhibit connective tissue synthesis. TGF- β_1 is a pleiotropic cytokine that induces connective tissue synthesis^{16,17} and inhibits growth of vascular smooth muscle cells.¹⁸ These are the two main changes observed in corpus cavernosum histopathology (for example, increases in connective tissue and atrophy and death of smooth muscle cells).^{2,9-11} TGF- β_1 alters the composition of extracellular matrix surrounding the cells by inducing expression of collagens, fibronectin and proteoglycans, while inhibiting the activity and expression of collagenases and other proteases.^{16,17} This results in a net collagen accumulation in the extracellular matrix. It has been demonstrated in normal rat kidney cells in culture that the growth inhibitory effects of TGF- β_1 are mediated in part through fibrillar collagen accumulation.¹⁹ In studies on pulmonary, hepatic and renal fibrosis, severity of disease was correlated with increased expression of TGF- β_1 .^{15,16} In the vasculature, increased expression of TGF- β_1 has been correlated with diffuse intimal thickening in human coronary arteries and human restenosis lesions.¹⁶ Transfection of the TGF- β_1 gene

into porcine femoral arteries results in a fibrocellular hyperplasia²⁰ with histologic similarities to corpus cavernosum biopsies from diseased patients.^{2,9-11} Thus, TGF- β_1 is a good candidate for a fibrogenic cytokine in the corpus cavernosum.

What factors potentially regulate the TGF- β_1 response? It is known that prostaglandin E relaxes corporal smooth muscle strips in organ baths; while PGF₂ α , thromboxane A₂ and prostacyclin elicit corpus cavernosal contractions.¹³ Interestingly, relaxation may not be the only effect of PGE. Previous studies have shown that PGE₁²⁵ and PGE₂²⁶ suppress collagen synthesis in primary cultures of human lung fibroblasts. Treatment with PGE₁ resulted in a 47% reduction in collagen biosynthesis with little or no effect on other secreted proteins.²⁵ TGF- β_1 -induced collagen synthesis in fibroblasts is suppressed by PGE₂.²⁶ Consistent with this observation, cultured lung fibroblasts from patients suffering from idiopathic pulmonary fibrosis have a diminished capacity to synthesize PGE₂.²⁷ In order to establish a cause and effect relationship of PGE in prophylaxis of lung fibrosis, prostaglandin G/H synthase (PGHS, a key enzyme in PGE synthesis) was transfected into rabbits. Animals were treated with pulmonary dosages of bacterial lipopolysaccharide which results in pulmonary hypertension and fibrosis. PGHS-transfected rabbit lungs had an increased capacity to synthesize PGE₂ and were protected from endotoxin-induced pulmonary hypertension.²⁸ Finally, PGE, which is derived from twenty carbon, polyunsaturated fatty acids (dihomo- γ -linoleic acid for PGE₁ and arachadonic acid for PGE₂), is synthesized by a series of reactions in which these fatty acids are converted into prostaglandin H by prostaglandin G/H synthase in an oxygen dependent reaction then to PGE by PGE epimerase. It is the oxygen dependence of PGE synthesis which may underlie its role in hypoxemia and penile fibrosis.

What is the evidence that TGF- β_1 is synthesized by corpus cavernosum smooth muscle cells and its synthesis regulated by oxygen tension?

We have demonstrated the expression of TGF- β_1 mRNA in human corpus cavernosum by Northern blot analysis,¹⁴ reverse-transcribed polymerase chain reaction (RT-PCR) of archival surgical specimens,² and TGF- β protein by immunohistochemical staining in corpus cavernosum biopsies.²¹ Further, oxygen tensions consistent with flaccid blood PO₂ (30 mm Hg) induce TGF- β_1 mRNA expression approximately two-fold in eighteen hours and three-fold in twenty four hours.²¹ What is the evidence that PGE is synthesized by corpus cavernosum

smooth muscle cells and its synthesis regulated by oxygen tension? Both human²⁹ and rabbit³⁰ corpus cavernosum in organ culture synthesize PGE; in the rabbit,³⁰ oxygen tensions consistent with flaccid blood PO₂ (30 mm Hg) inhibit this production. We find that human corpus cavernosum smooth muscle cells in culture also synthesize PGE and that this process is inhibited by PO₂ of 30 mm Hg.²¹ Thus, human corpus cavernosum smooth muscle cells synthesize both TGF- β_1 and PGE; and oxygen tensions consistent with flaccid blood PO₂ induce TGF- β_1 expression and inhibit PGE synthesis.

What is the evidence that TGF- β_1 and PGE play a role in human corpus cavernosum smooth muscle collagen synthesis?

TGF- β_1 induces a 2–4-fold increase in collagen synthesis in human corpus cavernosum smooth

muscle cells in culture.^{2,14} The actions of TGF- β_1 are dependent on three different subclasses of cell surface receptors.^{22,23} Human corpus cavernosum smooth muscle cells express all three types of TGF- β receptors as determined by crosslinking to [¹²⁵I]-TGF- β_1 .^{2,14} The type II receptor is expressed more than the type I, consistent with a smooth muscle phenotype.²⁴ The dose of TGF- β_1 which results in half-maximal stimulation of collagen synthesis in human corpus cavernosum smooth muscle cells is consistent with the reported affinity constants for the high affinity receptors (Kd ~ 8 pM).¹⁴ Cells derived from each of three groups in the study above comparing DICC results to percentage trabecular smooth muscle² all express TGF- β_1 mRNA, and TGF- β_1 autoinduces its own mRNA as well as all three types of TGF- β receptors. Thus, TGF- β_1 can induce itself and its receptors, perhaps perpetuating a fibrotic response; consistent with what Border and Ruoslahti termed 'the dark side of fibrosis'.¹⁶ We have reported that PGE₁, which elevates cyclic AMP

Figure 2 Working hypothesis of regulation of collagen synthesis in human corpus cavernosum smooth muscle. Left panel depicts activity of PGHS1/2 in the synthesis of PGE precursors and finally PGE at high oxygen tension associated with the erect state. PGE binds to specific EP2 and/or EP4 receptors which, through a Gs protein-coupled signal, activate adenylate cyclases to synthesize cAMP. cAMP inhibits collagen synthesis. Phosphodiesterases (PDE) terminate the cAMP signal by hydrolysis to AMP; a process inhibited by PDE inhibitors such as papaverine, or the type 3 PDE inhibitor, milrinone. Right panel depicts at low oxygen tension associated with the flaccid state, TGF- β_1 expression is increased, TGF- β_1 is secreted, activated and binds to high affinity type I and type II receptors (TGF- β RI and TGF- β RII). This increases TGF- β_1 expression, expression of TGF- β receptors and induces collagen and connective tissue synthesis. Both pathways at high and low oxygen tension work together to yield the final connective tissue/smooth muscle balance which may be altered in diseased states.²

in human corpus cavernosum smooth muscle cells,²² suppresses TGF- β_1 -induced collagen synthesis.^{14,22} PGE₁ can also suppress TGF- β_1 -induced TGF- β_1 mRNA autoinduction, and to some degree, TGF- β_1 induced increase in TGF- β receptors.¹⁴ Thus, PGE synthesized by smooth muscle cells, may suppress the actions of endogenously synthesized TGF- β_1 . Therefore, the interaction between PGE and TGF- β_1 pathways may key in determining connective tissue/smooth muscle balance in the corpus cavernosum (Figure 2).

All of the data in support of a role of hypoxemia in penile fibrosis discussed so far is from tissue culture and *in vitro* models.

Is there any evidence in animal models of disease that supports this hypothesis?

Three such models will be discussed. A rabbit model of atherosclerosis has been developed in which a high cholesterol diet in combination with iliac artery balloon injury leads to corpus cavernosum ischemia and erectile dysfunction.^{31,32} In this model, impaired erectile tissue function and decreased corporal blood flow increased with severity of disease.^{31,33} When penile biopsy specimens were examined for each of the three rabbit groups (control, hypercholesterolemic, and atherosclerotic, ischemic), it was found that the ratio of trabecular smooth muscle to total tissue area decreased with increasing severity of disease (45.4% in control, 39.1% in hypercholesterolemic, and 33.9% in atherosclerotic, ischemic).³⁴ Finally, in the atherosclerotic animals, an increase in corpus cavernosum tissue TGF- β_1 and decreased capacity for PGE synthesis in organ culture was noted when compared to control animals.³⁵

In an alloxan-induced diabetic rabbit model, it was found that diabetes mellitus impaired both neurogenic and endothelium-dependent relaxation of rabbit corpus cavernosum smooth muscle.³⁶ When penile biopsy specimens were examined for each of the three rabbit groups control (blood glucose 111 ± 2 mg/dl), mild diabetic (blood glucose 133 ± 4 mg/dl but decreased blood insulin levels), and severe diabetic (low blood insulin, blood glucose 416 ± 20 mg/dl), it was found that the ratio of trabecular smooth muscle to total tissue area decreased with increasing severity of disease (42.5% in control, 39.5% in mild diabetic, and 35.4% in severe diabetic)³⁷ similar to the results in the atherosclerotic rabbit model.³⁴ While direct correlations to TGF- β_1 and PGE levels are yet to be described in this model, the severity of disease is correlated with structural changes in the rabbit corpus cavernosum.

Both of these animal models of disease show trabecular structural changes which correlate with the severity of disease, and in one instance there is an increase in TGF- β_1 and a decrease in PGE.³⁵ What is the evidence that TGF- β_1 is necessary and sufficient as an agent of corpus cavernosum fibrosis? In order to establish an *in vivo* model for TGF- β_1 -induced corporal fibrosis, TGF- β_1 -impregnated slow release pellets were introduced into the rabbit corpus cavernosum.³⁸ Controls included placebo time release pellets and untreated controls. A significant increase in connective tissue was observed compared to placebo by 72 h of treatment (35.5% smooth muscle per erectile tissue area vs 40.0% for placebo and 43.5% for untreated controls).³⁸ This induction of corporal fibrosis is comparable to that observed in 16 week atherosclerotic, ischemic rabbits.³⁴ The expression of TGF- β_1 mRNA and protein in human corpus cavernosum, as well as induction of collagen synthesis in cell culture and connective tissue in an animal model suggests a role for TGF- β_1 in corpus cavernosum connective tissue synthesis.

Is hypogonadism associated with hypoxemia?

This is an interesting point to address as there are anecdotal reports of eugonadal and hypogonadal men with erectile dysfunction who have been successfully treated and recovered function by administration of androgen supplements. A recent review of androgens and penile erection stated that 'a critical examination of the literature fails to establish a clear association between erectile function in men, and blood levels of testosterone, dihydrotestosterone, or other androgens'.³⁹ This may be due in part to the dual sites of action of androgens (central and peripheral), the difficulty in constructing definitive clinical studies, and the fact that even in men with low blood levels of testosterone, potency is reported.⁴⁰ The discrepancies in the literature are illustrated by the following two studies. In one study, clomiphene citrate was administered to elevate blood testosterone levels in impotent men with secondary hypogonadism. Despite the return to normal testosterone blood levels, no overall improvement in erectile function was noted in any group; and limited improvement in nocturnal penile tumescence was observed only in young and healthy men.⁴¹ On the other hand in another recent report, testosterone undecanoate oral supplements improved self-reported erectile function in 61% hypogonadal males in the study.⁴² In humans, it is therefore difficult to assess the benefits of androgen therapy for hypogonadism; much less whether or not hypogonadism is associated with hypoxemia. Erection in the rat as well as neural nitric oxide synthase (nNOS) is regulated by androgens.^{39,43} In a recent report 40% of nNOS and 20% of vasoactive intestinal peptide containing nerves co-stained with androgen

receptor in the major pelvic ganglion of the adult rat penis,⁴⁴ suggesting a role of androgen in regulation of these neurons. When aged rats are compared to young rats, the erectile response remains intact to electrical and peripheral stimulation, but nNOS containing nerves as well as percent trabecular smooth muscle decreases in the aged group compared to the young group.⁴⁵ In contrast, an nNOS gene knockout in mice results in a strain which is not only potent but reproduces aggressively and is prone to violence with its litter mates.⁴⁶ These results suggest that while androgen regulation of neural nitric oxide synthase is an important component of erectile response in the rat, other components (endothelial nitric oxide synthase) can compensate for loss of this gene. They further fail to show an association between hypogonadism and hypoxemia.

We have examined TGF- β_1 and PGE as two possible factors that may regulate corpus cavernosum collagen metabolism, shown that both substances are both synthesized in the corpus cavernosum and that their synthesis is regulated by oxygen tension.

How might these substances interact to preserve a functional smooth muscle: connective tissue balance in the corpus cavernosum?

Nocturnal penile tumescence (NPT) was first reported in infants by Halverson.⁴⁷ Most men have three to five nocturnal episodes, each lasting anywhere from 30–45 min and coincident with rapid eye movement (REM) sleep.⁴⁹ A recent report of term fetuses found a 22% incidence of penile tumescence using ultrasound imaging.⁴⁸ Therefore, NPT begins as human males are born and continues in a circadian fashion throughout life, declining in older age.⁵¹ As these episodes occur daily, they provide a brief period (anywhere from 1.5–3 h/d⁴⁹) of increased blood PO₂ in the corpus cavernosum and hence the potential to synthesize substances such as prostanoids or nitric oxide whose synthesis is favored by the higher oxygen tensions. However, since this amounts to only 6–12% of the day, the corpus cavernosum is at the lower blood PO₂ in the absence of sexual stimulation for most of the day. If this hypothesis is valid, why is there not fibrosis of the corpora? In studies in cell culture, one application of PGE₁ was sufficient, despite the predicted short half-life of PGE₁ to suppress TGF- β_1 -induced collagen synthesis.¹⁴ Thus, the brief elevation of cAMP levels in response to PGE₁ binding may be sufficient in human corpus cavernosum smooth muscle cells²² as has been observed in human fetal lung fibroblasts^{25,26} to suppress TGF- β_1 -induced collagen synthesis. This dampening effect may be key in the maintenance of a functional trabecular smooth muscle/connective tissue balance.²

What clinical correlates exist to the importance of NPT or erections and potency?

Erectile dysfunction has been reported in patients in which REM sleep and NPT are absent (sleep apnea⁵¹) or disrupted (narcolepsy⁵² and epilepsy⁵³). Of particular interest is one study in which one third of the patients with sleep apnea and erectile dysfunction, continuous positive airway pressure improved erectile function as well as nocturnal penile tumescence.⁵⁴ While further studies are needed, it appears that 'erections may be good for preserving erections.' Radical retropubic prostatectomy, even when performed by nerve sparing techniques, often results in either a temporary or permanent loss of erectile function. A recent prospective study of nerve-sparing radical prostatectomy patients in Italy showed that those patients on postoperative prophylactic PGE₁ were more likely to recover erectile function than those on placebo.⁵⁵ Finally, while psychogenic effects cannot be ruled out, up to 28% of patients in one study using intracorporal PGE₁ achieved improved spontaneous erections and less frequent need for intracorporal injections.⁵⁶ Taken together, these studies suggest a role of hypoxemia in penile fibrosis and a role for and in preserving a functional smooth muscle/connective tissue balance in the corpus cavernosum.

Conclusions

While the study of the molecular pathology and etiology of corpus cavernosum fibrosis and vasculogenic impotence have just begun within the last ten years, basic research is needed to address which other factors regulate a functional smooth muscle/connective tissue ratio. How is connective tissue turnover regulated in the corpus cavernosum? Can it be accelerated? What is the molecular basis of preservation of potency? Is nocturnal penile tumescence a key in maintaining potency, and how can it be enhanced? If detected early, can treatment 'reverse' the condition? The issue of prophylaxis (for example, preservation of potency) both after radical retropubic prostatectomy and with aging is one which awaits new studies. Finally, a working knowledge of the basic science of these processes promises to lead to rational design of effective therapeutic strategies for the treatment of vasculogenic impotence.

Acknowledgements

The author is indebted to Drs RJ Krane, and I Goldstein, for their support, encouragement, and

insight. The helpful comments, suggestions and discussion of Drs A Nehra, A Traish, MT Watkins, and S Gupta, were key in the preparation of this talk and manuscript. This paper was presented at the Third Annual Meeting of the Society for the Study of Impotence, April 12, 1997, in New Orleans, Louisiana and was supported by grants DK47950, DK39080, DK40025 and DK40487 from the National Institutes of Health and in part, by a grant from Upjohn-Pharmacia.

References

- 1 Kim N *et al.* Oxygen tension regulates the nitric oxide pathway: Physiological role in penile erection. *J Clin Invest* 1993; **91**: 437–443.
- 2 Nehra A *et al.* Mechanisms of venous leakage: a prospective clinicopathological correlation of corporeal function and structure. *J Urol* 1996; **156**: 1320–1329.
- 3 Benet AE, Melman A. The epidemiology of erectile dysfunction. *Urol Clin N Am* 1995; **22**: 699–709.
- 4 Feldman HA *et al.* Impotence and its medical and psychosocial correlates: results of the Massachusetts Male Aging Study. *J Urol* 1994; **151**: 54–61.
- 5 Munarriz RM *et al.* Blunt trauma: the pathophysiology of hemodynamic injury leading to erectile dysfunction. *J Urol* 1995; **153**: 1831–1840.
- 6 Shafik A. Pudendal artery syndrome with erectile dysfunction: treatment by pudendal canal decompression. *Arch Androl* 1995; **34**: 83–94.
- 7 Persson C *et al.* Correlation of altered penile ultrastructure with clinical arterial evaluation. *J Urol* 1989; **142**: 1462–1468.
- 8 Jevtich M, Khawand NY, Vidic B. Clinical significance of ultrastructural findings in the corpus cavernosa of normal and impotent men. *J Urol* 1990; **143**: 289–293.
- 9 Mersdorf A *et al.* Ultrastructural changes in impotent penile tissue: a comparison of 65 patients *J Urol* 1991; **145**: 749–758.
- 10 Wespes E *et al.* Computerized analysis of smooth muscle fibers in potent and impotent patients. *J Urol* 1991; **146**: 1015–1017.
- 11 Luangkhot R *et al.* Collagen alterations in the corpus cavernosum of men with sexual dysfunction. *J Urol* 1992; **148**: 467–471.
- 12 Hatzichristou DG *et al.* *In vivo* assessment of trabecular smooth muscle tone, its application in pharmacocavernosometry and analysis of intracavernosal pressure determinants. *J Urol* 1995; **153**: 1126–1131.
- 13 Andersson K-E, Wagner G. Physiology of erection. *Physiol Rev* 1995; **75**: 191–236.
- 14 Moreland RB *et al.* PGE₁ suppresses the induction of collagen synthesis by transforming growth factor- β_1 in human corpus cavernosum smooth muscle. *J Urol* 1995; **153**: 826–834.
- 15 Woessner JF. Matrix metalloproteinases and their inhibitors in connective tissue remodeling. *FASEB J* 1991; **5**: 2145–2154.
- 16 Border WA, Ruoslahti E. Transforming growth factor- β in disease: the dark side of tissue repair. *J Clin Inv* 1992; **90**: 1–7.
- 17 Border WA, Noble NA. Transforming growth factor β in tissue fibrosis. *New Engl J Med* 1994; **331**: 1286–1291.
- 18 Majack RA. Beta-type transforming growth factor specifies organizational behavior in vascular smooth muscle cell cultures. *J Cell Biol* 1987; **105**: 465–471.
- 19 Nugent MA, Newman MJ. Inhibition of normal rat kidney cell growth by transforming growth factor- β_1 is mediated by collagen. *J Biol Chem* 1989; **264**: 18069–18077.
- 20 Nabel EG *et al.* Direct transfer of transforming growth factor beta 1 gene into arteries stimulates fibrocellular hyperplasia. *Proc Natl Acad Sci USA* 1993; **90**: 10759–10763.
- 21 Moreland RB, Watkins MT, Nehra A, Huang Y-H, Munarriz R, Salimpour P, Goldstein J and Traish AM. Oxygen tension modulates transforming growth factor β_1 expression and PGE production in human corpus cavernosum smooth muscle cells. *Mol Urol* 1998; **2**: 41–47.
- 22 Moreland RB *et al.* Cyclic AMP modulates TGF β_1 -induced fibrillar collagen synthesis in cultured human corpus cavernosum smooth muscle cells. *Int J Imp Res* 1998, in press.
- 23 Massague J, Attisano L, Wrana JL. The TGF- β family and its composite receptors. *Trends in Cell Biol* 1994; **4**: 172–178.
- 24 Goodman LV, Majack RA. Vascular smooth muscle cells express distinct transforming growth factor- β receptor phenotypes as a function of cell density. *J Biol Chem* 1989; **264**: 5241–5244.
- 25 Baum BJ *et al.* Effect of cyclic AMP on the intracellular degradation of newly synthesized collagen. *J Biol Chem* 1980; **255**: 2843–2847.
- 26 Fine A *et al.* The differential effect of prostaglandin E₂ on transforming growth factor- β and insulin-induced collagen formation in lung fibroblasts. *J Biol Chem* 1989; **264**: 16988–16991.
- 27 Wilborn J *et al.* Cultured lung fibroblasts isolated from patients with idiopathic pulmonary fibrosis have a diminished capacity to synthesize prostaglandin E₂ and to express cyclooxygenase-2. *J Clin Inv* 1995; **95**: 1861–1868.
- 28 Conary JT *et al.* Protection of rabbit lungs from endotoxin injury by *in vivo* hyperexpression of the prostaglandin G/H synthase gene. *J Clin Inv* 1994; **93**: 1834–1840.
- 29 Roy AC, Tan SM, Kottegoda SR, Ratnam SS. Ability of human corpus cavernosum to generate prostaglandins and thromboxanes *in vitro*. *IRCS Med Sci* 1984; **12**: 608.
- 30 Daley JT *et al.* Prostanoid production in rabbit corpus cavernosum I. Regulation by oxygen tension. *J Urol* 1996; **155**: 1482–1487.
- 31 Azadzi KM, Siroky MB, Goldstein I. Study of etiologic relationship of arterial atherosclerosis to corporal veno-occlusive dysfunction in the rabbit. *J Urol* 1996; **155**: 1795–1800.
- 32 Kim JH *et al.* Experimental hypercholesterolemia in rabbits induces cavernosal atherosclerosis with endothelial and smooth muscle cell dysfunction. *J Urol* 1994; **151**: 198–205.
- 33 Azadzi KM *et al.* Relationship between cavernosal ischemia and corporal veno-occlusive dysfunction in an animal model. *J Urol* 1997; **157**: 1011–1017.
- 34 Nehra A *et al.* Cavernosal expandability is an erectile tissue mechanical property which predicts trabecular histology in an animal model of vasculogenic erectile dysfunction. *J Urol* 1998, in press.
- 35 Azadzi KM *et al.* Ischemia-induced TGF- β_1 mediated cavernosal fibrosis is modulated by eicosanoids. *Int J Imp Res* 1996; **8**: 108 (Abstract A36).
- 36 Azadzi KM, Saenz de Tejada I. Diabetes mellitus impairs neurogenic and endothelium-dependent relaxation of rabbit corpus cavernosum smooth muscle. *J Urol* 1992; **148**: 1587–1591.
- 37 Gupta S *et al.* Diabetes-induced structural changes in rabbit corpus cavernosum. *Int J Imp Res* 1996; **8**: 136 (Abstract D55).
- 38 Nehra A *et al.* An *in vivo* model for transforming growth factor β_1 induced corporal fibrosis: implications for penile ischemia-assisted fibrosis. *J Urol* 1996; **155**: 622A (Abstract 1243).
- 39 Mills TM, Reilly CM, Lewis RW. Androgens and penile erection: a review. *J Androl* 1996; **17**: 633–638.
- 40 Buena F *et al.* Sexual function does not change when serum testosterone levels are pharmacologically varied within the normal male range. *Fertil Steril* 1995; **59**: 1118–1123.
- 41 Guay AT, Bansal S, Heatley GJ. Effect of raising endogenous testosterone levels in impotent men with secondary hypergonadism: double blind placebo-controlled trial with clomiphene citrate. *Clin Endocrinol Metab* 1995; **80**: 3546–3552.
- 42 Morales A, Johnston B, Heaton JP, Lundie M. Testosterone supplementation for hypogonadal impotence: assessment of biochemical measures and therapeutic outcomes. *J Urol* 1997; **157**: 849–854.

- 43 Penson DF *et al.* Androgen and pituitary control of penile nitric oxide synthase and erectile function in the rat. *Biol Reprod* 1996; **55**: 567–574.
- 44 Schirar A, Chang C, Rousseau JP. Localization of androgen receptor in nitric oxide synthase- and vasoactive intestinal peptide-containing neurons of the major pelvic ganglion innervating the rat penis. *J Neuroendocrinol* 1997; **9**: 141–150.
- 45 Carrier S *et al.* Age decreases nitric oxide synthase-containing nerve fibers in the rat penis. *J Urol* 1997; **157**: 1088–1092.
- 46 Burnett AL *et al.* Nitric oxide-dependent penile erection in mice lacking neuronal nitric oxide synthase. *Mol Med* 1996; **2**: 288–296.
- 47 Halverson HM. Genital and sphincter behavior of the male infant. *J Gen Psychol* 1940; **56**: 95.
- 48 Shirozu H *et al.* Penile tumescence in the human fetus at term—a preliminary report. *Early Hum Dev* 1995; **41**: 159–166.
- 49 Fischer C, Gross J, Zuch J. Cycle of penile erections synchronous with dreaming (REM) sleep: preliminary report. *Arch Gen Psychiatry* 1965; **12**: 29–45.
- 50 Schiavi RC *et al.* Hormones and nocturnal penile tumescence in healthy aging men. *Arch Sex Behav* 1993; **22**: 207–215.
- 51 Hirshkowitz M *et al.* Prevalence of sleep apnea in men with erectile dysfunction. *Urology* 1990; **36**: 232–234.
- 52 Karacan I. Erectile dysfunction in narcoleptic patients. *Sleep* 1986; **9**: 227–231.
- 53 Guldner GT, Morrell MJ. Nocturnal penile tumescence and rigidity evaluation in men with epilepsy. *Epilepsia* 1996; **37**: 1211–1214.
- 54 Karacan I, Karatas M. Erectile dysfunction in sleep apnea and response to CPAP. *J Sex and Marital Therapy* 1995; **21**: 239–247.
- 55 Montorsi F *et al.* Recovery of spontaneous erectile function after nerve-sparing radical retropubic prostatectomy with and without early intracavernous injections of Alprostadil: Results of a prospective randomized trial. *J Urol* 1997; **158**: 1408–1410.
- 56 Basile G *et al.* Can self-injection therapy “cure” impotence? In: Goldstein I and Lue TF, eds. *The Role of Alprostadil in the Diagnosis and Treatment of Erectile Dysfunction*. Excerpta Medica: Princeton, 1993, pp 109–116.