GUEST EDITORIAL

Association studies in psychiatric genetics

The term 'allelic association' was coined by Edwards¹ to describe lack of independence in a population between alleles at two closely linked loci. This simple descriptive term was proposed to replace several terms such as linkage inequality, gametic disequilibrium and linkage disequilibrium by which the phenomenon had previously been described.² The goal of allelic association studies of disease is to demonstrate a significantly different distribution of allelic variants in affected and unaffected individuals. There are essentially three possible explanations for this, two of which are scientifically interesting. The not so interesting explanation is that the disease association is actually due to population stratification. This occurs when recent admixtures of populations result in both the disease and the marker allele occurring at a higher frequency in one component of the population admixture than in another. Provided that stratification can be excluded, the causes of an association between a marker and a disease are likely to be either pleiotropy or the presence of very tight linkage between the marker and a susceptibility locus. Pleiotropy is the phenomenon whereby a marker has two (or more) properties consisting of: (a) being detectable as a polymorphism; and (b) having a direct effect on a trait such as susceptibility to a disease. Linkage detected in families does not usually result in association detected in populations. However, tight linkage can result in allelic association with a disease when the marker and a susceptibility locus are so close together that an association with a particular allele is maintained over many generations of recombination. In theory a fourth cause of allelic association is the operation of a common selective process at both loci. In practice few examples of this are known in man, but one is the association between G6PD deficiency, thalassaemia and resistance to malaria in certain areas of the Mediterranean.³

The physical distance over which detectable allelic association is maintained depends upon a number of factors, including the recombination frequency, the number of generations since the second mutational event occurred and attributes of the population such as selection, admixture and rate of population growth, as well as factors relating to the study such as the size of the sample and the informativity of the markers. This is an area that requires more work, but published data on RFLPs suggest that allelic association is detectable over distances up to 500 kb⁴ whereas recent stud-

ies with highly polymorphic microsatellite markers suggest that associations can often be detected over distances of one megabase or more.^{5,6} This fits well with fairly simple methods of estimating how long it takes an association to decay. For example, the 'half-life' of an association between alleles at loci 1 cM apart is about 69 generations or around 2000 years.

Association studies have considerable theoretical appeal in the analysis of common diseases since they can detect genes that contribute only a relatively small proportion of the overall liability of developing a disorder. For example, the association between blood group O and duodenal ulcer accounts for little over 1% of the variance in liability.⁷ Such a small effect could only be detected by linkage studies in prohibitively large samples. Risch and Merikangas⁸ show that well over 2000 families are needed to detect a gene with a genotype relative risk of 2 or less using allele-sharing linkage methods. In contrast, the required sample size for an association study is vastly less and is generally fewer than a thousand which is well within practical limits. The power of association studies to detect genes of small effect is particularly important since it seems increasingly likely that the majority of psychiatric disorders reflect the combined action of several or many genes of moderate to small effect (ie oligogenic or polygenic inheritance).

There are, however, a number of problems with association studies. First, they are limited by the fact the actual disease-causing polymorphism, or one very tightly linked to it, must be tested. Thus at the present time such studies are usually conducted using markers close to or within candidate genes. Unfortunately because we understand very little about the neurobiological substrates of most psychiatric disorders, it is rarely possible to specify highly plausible candidates (eg APP for familial Alzheimer's disease and PrP for familial prion diseases) and the prior odds against true association in any experiment are therefore substantial. This is even worse when a non-functional (so-called anonymous) polymorphism is studied which is not known to reside in close proximity to a candidate gene. A further problem is that of multiple testing which arises when several polymorphisms are tested and/or each polymorphism has multiple alleles. This means that if say 20 patient-control comparisons are made, on average, one 'significant' allelic association will be found even if no true association exists. A conservative approach is to apply the so-called Bonferroni correction and multiply the threshold for statistical significance by the number of tests, though this in turn runs the risk of increasing type II errors. Clearly some attempt must be made to reduce the probability of false

positives, but setting a very stringent test criterion may result in true positives being missed. Therefore it seems preferable to regard the occurrence of some false positives as inevitable and to rely upon replication to identify true positive findings. A partial solution is for individual groups to attempt their own replication by splitting their sample into two. All markers are tested in the first part but only those satisfying a not too stringent alpha level (say 0.01) are tested in the second.⁹

In addition to the study of candidate genes, association studies are likely to be used increasingly to narrow down regions of interest after preliminary localisation of a gene by linkage studies. Here it is appropriate to use anonymous markers, usually closely-spaced, highly polymorphic microsatellites, but due attention must be given to the issue of multiple testing. The prior odds of association in this situation are more difficult to gauge, since the presence of linkage does not necessarily imply the presence of association though it will increase the prior odds of association over those in an unlinked region. Once again replication will be required to be sure of an association and individual groups might like to rely upon the split-sample design.

Another important problem with traditional case-control association studies comes when there is inadequate matching between the patient and control groups. In this situation it is possible for stratification effects to produce either spurious positive or negative results and this is exemplified in publications on classical genetic markers in schizophrenia.¹⁰ Other than by ensuring that studies are carried out on homogeneous populations with good ethnic matching between patients and controls, association studies can be carried out on family material consisting of affected individuals and both their parents. In such parent-offspring trios, comparing the frequency of transmitted with untransmitted alleles provides an internal control that is not affected by stratification.^{11,12} Such 'intra-familial association studies' are becoming increasingly popular in psychiatric genetics.

With these issues in mind how do we interpret data from disease association studies and apparent discrepancies between studies? From the foregoing it should be clear that positive findings should be treated with a fair degree of caution, at least until a consistent pattern of replication has emerged. Certainly more credence should be given to findings involving polymorphisms that are likely to alter structure or expression of plausible candidate genes, particularly when a functional alteration in the predicted direction has actually been demonstrated. Similarly a finding that is strongly statistically significant ($P < 0.001$) even after correction for multiple testing is more likely to be robust. However, even under these ideal circumstances the false positive rate is likely to be high.¹³ Positive studies should also be scrutinised carefully for the possibility that the findings have resulted from population stratification. The use of an intra-familial design is particularly reassuring in this regard, though it should be remembered that studies of this sort do not distinguish between linkage and association unless only a single affected individual

from each family has been studied.¹⁴ If a case control design has been used then evidence of careful ethnic matching should be sought.

On the other hand failures to replicate and negative studies should be scrutinised carefully too, particularly as regards the issue of power. Many small negative studies are reported which actually have very little power to detect moderate or small effect sizes. We show in Table 1 the approximate sample sizes required to have 80% power of detecting associations with alleles conferring a range of different relative risks. It can be seen that for relative risks of less than two, sample sizes of several hundred or more are often required depending upon the frequency of the disease-associated allele. For example, an association between schizophrenia and the T102C polymorphism in the 5HT2A receptor gene was recently observed in a large multicentre European collaborative study.¹⁵ A number of small 'negative' reports have been published, but we can estimate that the sample needed to replicate this finding at the 5% significance level with a power of >0.80 is about 1000. Indeed meta-analysis of all published data which included over 3000 subjects supports the original finding and suggests that publication bias is unlikely.¹⁶

From the foregoing we can begin to see how we should respond to conflicting reports where some studies show a significant association between a specific polymorphism and a disease whereas others do not. Broadly speaking we need to consider both procedural and statistical issues. As far as procedures are concerned particular attention should be paid to the possibility of population stratification and also the question

Table 1 Sample sizes needed for 80% power (5% significance level) for range of relative risks (RR) associated with a biallelic marker

	RR	Attributable risk					
		0.05	0.1	0.2	0.3	0.4	0.5
Dominant	1.25	1921	1528				
	1.5	905	547	499	1875		
	1.75	628	354	249	308	1528	
	2	499	273	174	170	273	
	2.5	374	199	116	96	104	176
	3	310	164	92	71	69	84
	4	249	128	69	51	45	45
	5	219	112	58	41	35	33
Recessive	1.25	2671	1528				
	1.5	1546	700	444	931		
	1.75	1179	503	259	239	649	
	2	986	409	194	152	180	
	2.5	781	316	140	96	86	108
	3	669	267	114	75	60	62
	4	541	213	88	56	41	37
	5	472	186	75	47	35	30

Testing of allele numbers in cases vs controls via 2×2 contingency tables. NB The controls are assumed to be disease-free and to have population allele frequencies.

of whether there are differences in diagnostic practices, sample ascertainment and ethnicity. Of course discrepant results may be real and reflect heterogeneity with a genetic risk factor playing a quantitatively or qualitatively different role in different diagnostic or ethnic groups. Also, where the association is based on close linkage, ethnic differences may result from differences in allelic association in different populations. This raises the question of what exactly should be classed as a 'failure to replicate'. In our view this term should be reserved for studies of the same populations using similar ascertainment and diagnostic practices. The first statistical issue to consider is how statistically significant are the positive findings? Some appreciation of the prior probability of association is important here, though for many psychiatric disorders this is likely to be difficult and based on little more than intuition. The power of the negative studies also needs to be considered. In making these judgements greater credence should be given to the results of larger studies and there is certainly a place for meta-analysis. This should include methods such as funnel plot analysis¹⁷ in order to determine whether publication bias in the form of selective non-publication of small negative studies is likely to be operating.

In an earlier issue the Editor asks what might be the meaning of a statistically significant association, in which 60% of individuals with a common psychiatric disorder have a specific polymorphism in a plausible candidate gene in contrast to an overall instance of 35% in the general population.¹⁸ Assuming that such a finding is robust (and the stated P value of $<10^{-5}$ is promising!), what, he asks, is the value of such a finding? In this example the relative risk of disease associated with the polymorphism is approximately 2.8. This suggests either that the polymorphism itself confers an approximately three-fold increased risk of developing the disorder or that it is closely linked to a polymorphism which confers a larger risk. In order to resolve this issue further polymorphisms need to be sought in the area and further association and linkage studies performed. There may be difficulties in distinguishing the pathogenic variant from other 'hitch-hiking' polymorphisms that show allelic association with it¹⁹ but once this has been achieved it will provide strong evidence that the specific gene is indeed involved in conferring susceptibility to the disorder. Given a reasonably modest relative risk such as this, it is unlikely that detection of this association alone will greatly facilitate diagnosis. However, the position might alter as other risk factors are identified. Moreover, it is possible that the risk allele might be associated with a particular course or outcome or with response to treatment.

It is important to note that a genetic factor that is common in the population can be associated with a high attributable risk even though the relative risk may be low. This means that some therapy targeting the immediate effects of the risk factor can have quite a large impact on the disorder at a population level in spite of the small relative risk. This is intuitively easier to grasp when the notion of liability-threshold models

is considered. Thus a risk factor may be neither necessary nor sufficient to cause illness but, if common, result in a substantial number of potential cases passing beyond the threshold for developing the disorder. The message here then is that the identification of genes of small effect might directly be of therapeutic importance as well as being essential if we are to build up a complete picture of the pathophysiology of the disorder, itself a prerequisite for rational therapy.

What recommendations can we make for the conduct of association studies in psychiatric genetics? First and most importantly, we must study samples of adequate size. Power to detect and replicate modest or small effects of the size that are likely to be operating in the majority of psychiatric disorders requires samples of the order of several hundred to a thousand (Table 1). We believe that inclusion of appropriate power calculations should be a mandatory prerequisite for the publication of 'negative' association studies. However it is also important that publication bias does not operate to prevent the dissemination of negative findings. In this regard we welcome the announcement that *Molecular Psychiatry* is to introduce a Scientific Correspondence section to allow the publication of brief research reports of complete studies.* There is also a need to identify epidemiologically meaningful samples to allow the size of gene effects to be measured at a population level and also to allow potential interactions with other genes and with environmental factors to be studied.

Studies of candidate genes should focus upon polymorphisms that lie within the coding or regulatory regions of the gene and are likely to result in altered structure or expression of the protein.²⁰ Where association studies are being conducted to achieve greater localisation within a linked region then highly polymorphic microsatellites are the markers of choice. These should be closely spaced at an interval of 1 cM or less.

Strenuous efforts should be made to reduce the possibility of spurious results due to population stratification. In case control studies the groups should be carefully matched for ethnicity and be drawn from the same ethnically homogeneous population. In late onset disorders the controls and patients should be matched for age since some genes influence longevity. In populations such as the USA where admixture is a much greater problem than say in Northern Europe, intra-familial association studies offer a solution. However, they do not overcome the problem of ethnic differences in disease aetiology or allelic association due to tight linkage. It may therefore be prudent even when an intra-familial design is used, to draw samples from as ethnically a homogeneous population as is possible. Furthermore, in adult onset disorders parent-offspring

*Editor's note: *Molecular Psychiatry* maintains its editorial policy of not publishing pilot studies, preliminary data, or case reports. Scientific correspondence is a new section for brief reports: for more information, please refer to our revised notes for contributors, in this issue of *Molecular Psychiatry*.

Table 2 Sample sizes needed for 80% power when a proportion of the 'control' population has the disease

	Proportion of 'controls' affected (%)	Attributable risk				
		0.05	0.1	0.2	0.3	0.4
Dominant	0	499	273	174	170	273
	1	511	279	176	174	279
	5	561	306	194	188	302
	10	637	346	219	211	338
	15	724	393	247	237	378
	20	830	450	281	271	427
Recessive	0	986	409	194	152	180
	1	1008	419	197	152	184
	5	1096	454	213	166	199
	10	1224	507	237	184	219
	15	1376	571	267	205	243
	20	1558	645	302	231	271

Assumes a relative risk of 2 and 5% significance level.

trios are hard to find; in our experience both parents can be sampled in only about 30% of cases of schizophrenia or bipolar disorder. Clearly there is a possibility that by focusing only upon trios, one is introducing unknown biases.

For relatively uncommon disorders such as schizophrenia and bipolar disorder with lifetime risks of 1% or less, it is not necessary to screen controls for psychiatric disorder since the inadvertent inclusion of a few affected cases will have little effect on power and does not warrant the time and expense of interviewing controls (Table 2). In addition, there are problems with using 'superclean' controls from which all cases with a history of psychiatric disorder have been excluded since allele frequency differences between groups might relate to the exclusion of a common disorder from the controls rather than the disorder being studied. For more common disorders such as major depression, where the rates of illness in the controls are likely to be higher, screening of cases for the disorder being studied is probably prudent as the loss of power becomes greater (Table 2). Finally, it goes without saying that diagnostic standards need to be high and there needs to be a clear description of how the samples and controls are ascertained.

As the human genome project progresses, more and more potential candidate genes will be identified. Indeed we can look forward to the time in the next 10–15 years when not only have all genes been identified but all common intragenic variation also.^{8,21} This together with advances in technology means that we can begin to envisage a time when genome-wide association studies of common diseases will be possible. These will require large, well-characterised samples and will be subject to many of the same problems that we have considered here. However we believe that these difficulties can be overcome if due

attention is paid to procedural and statistical issues and then the rewards for psychiatry are likely to be great.

MJ Owen, P Holmans and P McGuffin
University of Wales College of Medicine
Cardiff, UK

References

- 1 Edwards JH. Allelic association in man. In: Eriksson AW (ed). *Population Structure and Genetic Disorders*. Academic Press: London, 1980.
- 2 Edwards JH. The formal problems of linkage. In: McGuffin P, Murray R (eds). *The New Genetics of Mental Illness*. Butterworth Heinemann: Oxford, 1991.
- 3 Cavalli-Sforza LL, Bodmer WF. *The Genetics of Human Populations*. WH Freeman: San Francisco, 1971.
- 4 Jorde LB, Watkins WS, Carlson M, Groden J, Albertsen H, Thliveris A, Leppert M. Linkage disequilibrium predicts physical distance in the adenomatous polyposis coli region. *Am J Hum Genet* 1994; **54**: 884–898.
- 5 Peterson AC, Di Rienzo A, Lehesjoki A-E, de la Chapelle A, Slatkin M, Freimer NB. The distribution of linkage disequilibrium over anonymous genome regions. *Hum Mol Genet* 1995; **4**: 887–894.
- 6 Petrukhin K, Fischer SG, Pirastu M, Tanzi RF, Chernov I, Devoto M, Brzustowicz LM, Cayanis E, Vitale E, Russo JJ, Matseoane D, Boukhgalter B, Wasco W, Figus AL, Loudianos J, Cao A, Sternlieb I, Evgrafov O, Parano E, Pavone L, Warburton D, Ott J, Penchaszadeh GK, Scheinberg IH, Gilliam TC. Mapping, cloning and genetic characterization of the region containing in Wilson disease gene. *Nature Genet* 1993; **5**: 338.
- 7 Edwards JH. The meaning of the association between blood groups and disease. *Ann Hum Genet* 1965; **29**: 77–83.
- 8 Risch N, Merikangas K. The future of genetic studies of complex human diseases. *Science* 1996; **273**: 1516–1517.
- 9 Owen MJ, McGuffin P. Association and linkage: complementary strategies for complex disorders. *J Med Genet* 1993; **30**: 638–639.
- 10 McGuffin P, Sturt E. Genetic markers in schizophrenia. *Hum Hered* 1986; **26**: 461–465.
- 11 Falk CT, Rubinstein P. Haplotype relative risks: easy reliable way to construct a proper control sample for risk calculations. *Ann Hum Genet* 1987; **51**: 227–233.
- 12 Schaid DJ, Sommer SS. Comparison of statistics for candidate-gene association studies using cases and parents. *Am J Hum Genet* 1994; **55**: 402–409.
- 13 Crowe RR. Candidate genes in psychiatry: an epidemiological perspective. *Am J Med Genet (Neuropsych Genet)* 1993; **48**: 74–77.
- 14 Ewens WJ, Spielman RS. The transmission/disequilibrium test: history, subdivision and admixture. *Am J Hum Genet* 1995; **57**: 455–464.
- 15 Williams J, Spurlock G, McGuffin P, Mallet J, Nothen MN, Gill M, Aschauer H, Nylander P-O, Maciardi F, Owen MJ. Association between schizophrenia and the T102C polymorphism of 5-hydroxytryptamine type 2a receptor gene. *Lancet* 1996; **347**: 1294–1296.
- 16 Williams J, McGuffin P, Nothen M, Owen MJ, EMAS Collaborative Group. A meta-analysis of association between the 5HT2A receptor T102C polymorphism and schizophrenia. *Lancet* 1997; **349**: 1221.
- 17 Egger M, Davey Smith G. Misleading meta-analysis. *Br Med J* 1995; **310**: 752–754.
- 18 Licinio J. Molecular psychiatry: does a new field need to examine strategy? *Mol Psychiatry* 1997; **2**: 3–4.
- 19 Craddock N, Owen MJ. Candidate gene association studies in psychiatric genetics: a SERTain future? *Mol Psychiatry* 1996; **1**: 434–436.
- 20 Sobell JL, Heston LL, Sommer SS. Delineation of genetic predisposition to multifactorial disease: a general approach on the threshold of feasibility. *Genomics* 1992; **12**: 1–6.
- 21 Lander ES. The new genomics: global views of biology. *Science* 1996; **274**: 540–545.