synthesized by PCR. DNase I footprinting reactions were processed as described²⁴.

Reporter gene construction. Substitution of a single *Bgl*II site for the quadrant enhancer Mad binding-site sequence GCTGCCGTCGCG was done by assembling two fragments generated by PCR with *Pfu* polymerase. Expansion of the single *Bgl*II site to two tandem sites matching the Q^m sequence was done by insertion of an oligonucleotide containing *Bbs*I internal sites that allowed for subsequent cleavage and ligation to generate the tandem *Bgl*II sites. Both sequences were confirmed by sequencing and ligated into the *Kpn*I/*Not*I sites of the Hsp *lacZ*-CaSpeR *Drosophila* transformation vector for the generation of transgenic lines²⁴.

Received 20 January; accepted 21 May 1997.

- Graff, J. M., Bansal, A. & Melton, D. A. *Xenopus* Mad proteins transduce distinct subsets of signals for the TGF β superfamily. *Cell* **85**, 479–487 (1996).
- Hoodless, P. A. *et al.* MADR1, a MAD-related protein that functions in BMP2 signaling pathways. *Cell* **85**, 489–500 (1996).
- Liu, F. *et al.* A human Mad protein acting as a BMP-regulated transcriptional activator. *Nature* **381**, 622–623 (1996).
- Eppert, K. *et al.* MADR2 maps to 18q21 and encodes a TGF β -regulated MAD-related protein that is functionally mutated in colorectal carcinoma. *Cell* **86**, 543–552 (1996).
- Zhang, Y., Feng, X.-H., Wu, R.-Y. & Derynck, R. Receptor-associated Mad homologues synergize as effectors of the TGF- β response. *Nature* **383**, 168–172 (1996).
- Lagna, G., Hata, A., Hemmati-Brivnlou, A. & Massagué, J. Partnership between DPC4 and SMAD proteins in TGF- β signalling pathways. *Nature* **383**, 832–836 (1996).
- Savage, C. *et al.* *C. elegans* genes *smo-2*, *smo-3*, and *smo-4* define a conserved family of TGF- β pathway components. *Proc. Natl Acad. Sci. USA* **93**, 790–794 (1996).
- Macias-Silva, M. *et al.* MADR2 is a substrate of the TGF- β receptor and its phosphorylation is required for nuclear accumulation and signaling. *Cell* **87**, 1215–1224 (1996).
- Derynck, R. & Zhang, Y. Intracellular signalling: The Mad way to do it. *Curr. Biol.* **6**, 1226–1229 (1996).
- Chen, X., Rubock, M. J. & Whitman, M. A transcriptional partner for MAD proteins in TGF- β signalling. *Nature* **383**, 691–696 (1996).
- Rafferty, L., Twombly, V., Wharton, K. & Gelbart, W. Genetic screens to identify elements of the decapentaplegic pathway in *Drosophila*. *Genetics* **139**, 241–254 (1995).
- Sekelsky, J., Newfeld, S., Rafferty, L., Chartoff, E. & Gelbart, W. Genetic characterization and cloning of *Mothers against dpp*, a gene required for decapentaplegic function in *Drosophila melanogaster*. *Genetics* **139**, 1347–1358 (1995).
- Newfeld, S. J., Chartoff, E. H., Graff, J. M., Melton, D. A. & Gelbart, W. M. *Mothers against dpp* encodes a conserved cytoplasmic protein required in DPP/TGF- β responsive cells. *Development* **122**, 2099–2108 (1996).
- Wiersdorff, V., Lecuit, T., Cohen, S. M. & Mlodzik, M. *Mad* acts downstream of Dpp receptors, revealing a differential requirement for *dpp* signaling in initiation and propagation of morphogenesis in the *Drosophila* eye. *Development* **122**, 2153–2162 (1996).
- Blair, S. Compartments and appendage development in *Drosophila*. *BioEssays* **17**, 229–309 (1995).
- Basler, K. & Struhl, G. Compartment boundaries and the control of *Drosophila* limb pattern by hedgehog protein. *Nature* **368**, 208–214 (1994).
- Tabata, T. & Kornberg, T. Hedgehog is a signalling protein with a key role in patterning *Drosophila* imaginal discs. *Cell* **76**, 89–102 (1994).
- Zecca, M., Basler, K. & Struhl, G. Sequential organizing activities of engrailed, hedgehog and decapentaplegic in the *Drosophila* wing. *Development* **121**, 2265–2278 (1995).
- Ingham, P. W. & Fietz, M. J. Quantitative effects of hedgehog and decapentaplegic activity on the patterning of the *Drosophila* wing. *Curr. Biol.* **5**, 432–440 (1995).
- Posakony, L., Rafferty, L. & Gelbart, W. Wing formation in *Drosophila melanogaster* requires decapentaplegic gene function along the anterior–posterior compartment boundary. *Mech. Dev.* **33**, 69–82 (1991).
- Capdevilla, J. & Guerrero, I. Targeted expression of the signalling molecule decapentaplegic induces pattern duplications and growth alterations in *Drosophila* wings. *EMBO J.* **13**, 4459–4468 (1994).
- Nellen, D., Burke, R., Struhl, G. & Basler, K. Direct and long-range action of a Dpp morphogen gradient. *Cell* **85**, 357–368 (1996).
- Lecuit, T. *et al.* Two distinct mechanisms for long-range patterning by Decapentaplegic in the *Drosophila* wing. *Nature* **381**, 387–393 (1996).
- Kim, J. *et al.* Integration of positional signals and regulation of wing formation and identity by *Drosophila* vestigial gene. *Nature* **382**, 133–138 (1996).
- de Celis, J. F., Barrio, R. & Kafatos, F. C. A gene complex acting downstream of *dpp* in *Drosophila* wing morphogenesis. *Nature* **381**, 421–424 (1996).
- Grimm, S. & Pflugfelder, G. O. Control of the gene optomotor-blind in *Drosophila* wing development by decapentaplegic and wingless. *Science* **271**, 1601–1604 (1996).
- Chouinard, S. & Kaufman, T. C. Control of expression of the homeotic *labial* (*lab*) locus of *Drosophila melanogaster*: evidence for both positive and negative autogenous regulation. *Development* **113**, 1267–1280 (1991).
- Tremml, G. & Bienz, M. Induction of *labial* expression in the *Drosophila* endoderm: response elements for *dpp* signalling and for autoregulation. *Development* **116**, 447–456 (1992).
- Thüringer, F., Cohen, S. M. & Bienz, M. Dissection of an indirect autoregulatory response of a homeotic *Drosophila* gene. *EMBO J.* **12**, 2419–2430 (1993).
- Eresh, S., Riese, J., Jackson, D. B., Bohmann, D. & Bienz, M. A CREB-binding site as a target for decapentaplegic signalling during *Drosophila* endoderm induction. *EMBO J.* **16**, 2014–2022 (1997).
- Ausubel, F. M. *et al.* (eds) *Current Protocols in Molecular Biology* supplement 28, pp. 16.6.1–16.7.7 (Wiley, New York, 1994).

Acknowledgements. We thank K. Vorwerk for technical assistance; M. Hoffmann for discussion; Y. J. Chen for the MAD cDNA clone and help with sequencing; V. Wiersdorff, M. Mlodzik and S. Cohen for the Mad¹⁻² stock; S. Paddock for help with microscopy; T. Kaufman for the labial endodermal enhancer; M. Hoffmann and G. Halder for comments on the manuscript; and J. Wilson for manuscript preparation. K. J. was supported by a genetics predoctoral training grant and the Cremer Scholarship Fund, and part of this work was supported by the Howard Hughes Medical Institute (S.B.C.).

Correspondence and requests for materials should be addressed to A.L. (e-mail: alaughon@facstaff.wisc.edu).

errata

Differential activation of transcription factors induced by Ca²⁺ response amplitude and duration

Ricardo E. Dolmetsch, Richard S. Lewis, Christopher C. Goodnow & James I. Healy

Nature **386**, 855–858 (1997)

The graphs in Fig. 2e were accidentally swapped around during the production process. The correct versions are shown here. □

Large-scale tectonic deformation inferred from small earthquakes

Falk Amelung & Geoffrey King

Nature **386**, 702–705 (1997)

Owing to an error in the production process, a length scale was omitted from Fig. 2b and the shading in Fig. 2d did not show up. The corrected figure is shown here. □

synthesized by PCR. DNase I footprinting reactions were processed as described²⁴.

Reporter gene construction. Substitution of a single *Bgl*II site for the quadrant enhancer Mad binding-site sequence GCTGCCGTCGCG was done by assembling two fragments generated by PCR with *Pfu* polymerase. Expansion of the single *Bgl*II site to two tandem sites matching the Q^m sequence was done by insertion of an oligonucleotide containing *Bbs*I internal sites that allowed for subsequent cleavage and ligation to generate the tandem *Bgl*II sites. Both sequences were confirmed by sequencing and ligated into the *Kpn*I/*Not*I sites of the Hsp *lacZ*-CaSpeR *Drosophila* transformation vector for the generation of transgenic lines²⁴.

Received 20 January; accepted 21 May 1997.

- Graff, J. M., Bansal, A. & Melton, D. A. *Xenopus* Mad proteins transduce distinct subsets of signals for the TGF β superfamily. *Cell* **85**, 479–487 (1996).
- Hoodless, P. A. *et al.* MADR1, a MAD-related protein that functions in BMP2 signaling pathways. *Cell* **85**, 489–500 (1996).
- Liu, F. *et al.* A human Mad protein acting as a BMP-regulated transcriptional activator. *Nature* **381**, 622–623 (1996).
- Eppert, K. *et al.* MADR2 maps to 18q21 and encodes a TGF β -regulated MAD-related protein that is functionally mutated in colorectal carcinoma. *Cell* **86**, 543–552 (1996).
- Zhang, Y., Feng, X.-H., Wu, R.-Y. & Derynck, R. Receptor-associated Mad homologues synergize as effectors of the TGF- β response. *Nature* **383**, 168–172 (1996).
- Lagna, G., Hata, A., Hemmati-Brivnlou, A. & Massagué, J. Partnership between DPC4 and SMAD proteins in TGF- β signalling pathways. *Nature* **383**, 832–836 (1996).
- Savage, C. *et al.* *C. elegans* genes *smo-2*, *smo-3*, and *smo-4* define a conserved family of TGF- β pathway components. *Proc. Natl Acad. Sci. USA* **93**, 790–794 (1996).
- Macias-Silva, M. *et al.* MADR2 is a substrate of the TGF- β receptor and its phosphorylation is required for nuclear accumulation and signaling. *Cell* **87**, 1215–1224 (1996).
- Derynck, R. & Zhang, Y. Intracellular signalling: The Mad way to do it. *Curr. Biol.* **6**, 1226–1229 (1996).
- Chen, X., Rubock, M. J. & Whitman, M. A transcriptional partner for MAD proteins in TGF- β signalling. *Nature* **383**, 691–696 (1996).
- Rafferty, L., Twombly, V., Wharton, K. & Gelbart, W. Genetic screens to identify elements of the decapentaplegic pathway in *Drosophila*. *Genetics* **139**, 241–254 (1995).
- Sekelsky, J., Newfeld, S., Rafferty, L., Chartoff, E. & Gelbart, W. Genetic characterization and cloning of *Mothers against dpp*, a gene required for decapentaplegic function in *Drosophila melanogaster*. *Genetics* **139**, 1347–1358 (1995).
- Newfeld, S. J., Chartoff, E. H., Graff, J. M., Melton, D. A. & Gelbart, W. M. *Mothers against dpp* encodes a conserved cytoplasmic protein required in DPP/TGF- β responsive cells. *Development* **122**, 2099–2108 (1996).
- Wiersdorff, V., Lecuit, T., Cohen, S. M. & Mlodzik, M. *Mad* acts downstream of Dpp receptors, revealing a differential requirement for *dpp* signaling in initiation and propagation of morphogenesis in the *Drosophila* eye. *Development* **122**, 2153–2162 (1996).
- Blair, S. Compartments and appendage development in *Drosophila*. *BioEssays* **17**, 229–309 (1995).
- Basler, K. & Struhl, G. Compartment boundaries and the control of *Drosophila* limb pattern by hedgehog protein. *Nature* **368**, 208–214 (1994).
- Tabata, T. & Kornberg, T. Hedgehog is a signalling protein with a key role in patterning *Drosophila* imaginal discs. *Cell* **76**, 89–102 (1994).
- Zecca, M., Basler, K. & Struhl, G. Sequential organizing activities of engrailed, hedgehog and decapentaplegic in the *Drosophila* wing. *Development* **121**, 2265–2278 (1995).
- Ingham, P. W. & Fietz, M. J. Quantitative effects of hedgehog and decapentaplegic activity on the patterning of the *Drosophila* wing. *Curr. Biol.* **5**, 432–440 (1995).
- Posakony, L., Rafferty, L. & Gelbart, W. Wing formation in *Drosophila melanogaster* requires decapentaplegic gene function along the anterior–posterior compartment boundary. *Mech. Dev.* **33**, 69–82 (1991).
- Capdevilla, J. & Guerrero, I. Targeted expression of the signalling molecule decapentaplegic induces pattern duplications and growth alterations in *Drosophila* wings. *EMBO J.* **13**, 4459–4468 (1994).
- Nellen, D., Burke, R., Struhl, G. & Basler, K. Direct and long-range action of a Dpp morphogen gradient. *Cell* **85**, 357–368 (1996).
- Lecuit, T. *et al.* Two distinct mechanisms for long-range patterning by Decapentaplegic in the *Drosophila* wing. *Nature* **381**, 387–393 (1996).
- Kim, J. *et al.* Integration of positional signals and regulation of wing formation and identity by *Drosophila* vestigial gene. *Nature* **382**, 133–138 (1996).
- de Celis, J. F., Barrio, R. & Kafatos, F. C. A gene complex acting downstream of *dpp* in *Drosophila* wing morphogenesis. *Nature* **381**, 421–424 (1996).
- Grimm, S. & Pflugfelder, G. O. Control of the gene optomotor-blind in *Drosophila* wing development by decapentaplegic and wingless. *Science* **271**, 1601–1604 (1996).
- Chouinard, S. & Kaufman, T. C. Control of expression of the homeotic *labial* (*lab*) locus of *Drosophila melanogaster*: evidence for both positive and negative autogenous regulation. *Development* **113**, 1267–1280 (1991).
- Tremml, G. & Bienz, M. Induction of *labial* expression in the *Drosophila* endoderm: response elements for *dpp* signalling and for autoregulation. *Development* **116**, 447–456 (1992).
- Thüringer, F., Cohen, S. M. & Bienz, M. Dissection of an indirect autoregulatory response of a homeotic *Drosophila* gene. *EMBO J.* **12**, 2419–2430 (1993).
- Eresh, S., Riese, J., Jackson, D. B., Bohmann, D. & Bienz, M. A CREB-binding site as a target for decapentaplegic signalling during *Drosophila* endoderm induction. *EMBO J.* **16**, 2014–2022 (1997).
- Ausubel, F. M. *et al.* (eds) *Current Protocols in Molecular Biology* supplement 28, pp. 16.6.1–16.7.7 (Wiley, New York, 1994).

Acknowledgements. We thank K. Vorwerk for technical assistance; M. Hoffmann for discussion; Y. J. Chen for the MAD cDNA clone and help with sequencing; V. Wiersdorff, M. Mlodzik and S. Cohen for the Mad¹⁻² stock; S. Paddock for help with microscopy; T. Kaufman for the labial endodermal enhancer; M. Hoffmann and G. Halder for comments on the manuscript; and J. Wilson for manuscript preparation. K. J. was supported by a genetics predoctoral training grant and the Cremer Scholarship Fund, and part of this work was supported by the Howard Hughes Medical Institute (S.B.C.).

Correspondence and requests for materials should be addressed to A.L. (e-mail: alaughon@facstaff.wisc.edu).

errata

Differential activation of transcription factors induced by Ca²⁺ response amplitude and duration

Ricardo E. Dolmetsch, Richard S. Lewis, Christopher C. Goodnow & James I. Healy

Nature **386**, 855–858 (1997)

The graphs in Fig. 2e were accidentally swapped around during the production process. The correct versions are shown here. □

Large-scale tectonic deformation inferred from small earthquakes

Falk Amelung & Geoffrey King

Nature **386**, 702–705 (1997)

Owing to an error in the production process, a length scale was omitted from Fig. 2b and the shading in Fig. 2d did not show up. The corrected figure is shown here. □

immunostained as described. Quantification was achieved by using Image-1 software (Universal Imaging). Positive controls included sections of adrenal medulla, thyroid, pancreas, intestine, ganglia and nerves.

Antisense studies. 100 nM phosphorothioate oligodeoxynucleotides with C-5 propyne modifications at dC and dU (Midland) were reconstituted with 2.5 $\mu\text{g ml}^{-1}$ GS2888 cytofectin³⁰ (Gilead Sciences) in OptiMEM (Gibco), then incubated with 5×10^5 NCI-H209 or DMS53 cells in 6-well plates for 48 h in growth media. 5×10^4 cells were cytospun per slide. Immunostaining, quantified using MetaMorph software (Universal Imaging), included at least ten cell clusters (over 200 cells) for each marker and oligonucleotide. Experiments were done in triplicate for both cell lines and evaluated in a blind fashion. Oligonucleotides c5p1476 (5'-GGAGCCACUGCUUU-3'), c5p1232 (5'-UCCCAACGCCACUGA-3') and c5p25 (5'-UCCUACUAGGCUGC-3') correspond to positions 1,476–1,491 and 1,232–1,247 in the hASH1 3' UTR and 25–40 in the 5' UTR, respectively; the c5pMiss1 sequence is 5'-GCUACAU-CUGGUCG-3'; c5pMiss2 5'-CACUGAUGCACCUGU-3'.

Received 9 January; accepted 20 February 1997.

1. Campuzano, S. *et al.* Molecular genetics of the achaete-scute gene complex of *D. melanogaster*. *Cell* **40**, 327–338 (1985).
2. Jan, Y. N. & Jan, L. Y. Genetic control of cell fate specification in *Drosophila* peripheral nervous system. *Annu. Rev. Genet.* **28**, 373–393 (1994).
3. Johnson, J. E., Birren, S. J. & Anderson, D. J. Two rat homologues of *Drosophila* *achaete-scute* specifically expressed in neuronal precursors. *Nature* **346**, 858–861 (1990).
4. Lo, L. C. *et al.* Mammalian *achaete-scute* homolog 1 is transiently expressed by spatially restricted subsets of early neuroepithelial and neural crest cells. *Genes Dev.* **5**, 1524–1537 (1991).
5. Sommer, L. *et al.* The cellular function of MASH1 in autonomic neurogenesis. *Neuron* **15**, 1245–1258 (1995).
6. Guillemot, F. *et al.* Mammalian *achaete-scute* homolog 1 is required for the early development of olfactory and autonomic neurons. *Cell* **75**, 463–476 (1993).
7. Baylín, S. B. & Gazdar, A. F. in *Small Cell Lung Cancer* 123–143 (Grune and Stratton, New York, 1981).
8. Gazdar, A. F. *et al.* Expression of neuroendocrine cell markers L-dopa decarboxylase, chromogranin A, and dense core granules in human tumors of endocrine and nonendocrine origin. *Cancer Res.* **48**, 4078–4082 (1988).
9. Gazdar, A. F. *et al.* in *Small Cell Lung Cancer* 145–177 (Grune and Stratton, New York, 1981).
10. Linnoila, R. I. in *Neuropeptides in Respiratory Medicine* 197–224 (Dekker, New York, 1994).
11. Mabry, M. *et al.* Transitions between lung cancer phenotypes—implications for tumor progression. *Cancer Cells* **3**, 53–58 (1991).
12. Ball, D. W. *et al.* Identification of a human *achaete-scute* homolog highly expressed in neuroendocrine tumors. *Proc. Natl Acad. Sci. USA* **90**, 5648–5652 (1993).
13. Linnoila, R. I. *et al.* Neuroendocrine differentiation in endocrine and nonendocrine lung carcinomas. *Am. J. Clin. Pathol.* **90**, 641–652 (1988).
14. Linnoila, R. I., Piantadosi, S. & Ruckdeschel, J. C. Impact of neuroendocrine differentiation in non-small cell lung cancer. *Chest* **106**, 367S–371S (1994).
15. Linnoila, R. I. & Aisner, S. C. in *Lung Cancer* 73–95 (Wiley-Liss, New York, 1995).
16. Carney, D. N. *et al.* Establishment and identification of small cell lung cancer cell lines having classic and variant features. *Cancer Res.* **45**, 2913–2923 (1985).
17. Pettengill, O. S. *et al.* Isolation and growth characteristics of continuous cell lines from small-cell carcinoma of the lung. *Cancer Res.* **45**, 906–918 (1980).
18. Bergh, J. *et al.* Establishment and characterization of a continuous lung squamous cell carcinoma cell line (U-1752). *Anticancer Res.* **1**, 317–322 (1981).
19. Gazdar, A. F. *et al.* Establishment of continuous, clonable cultures of small-cell carcinoma of lung which have amine precursor uptake and decarboxylation cell properties. *Cancer Res.* **40**, 3502–3507 (1980).
20. Gazdar, A. F. *et al.* Characterization of variant subclasses of cell lines derived from small cell lung cancer having distinctive biochemical, morphological, and growth properties. *Cancer Res.* **45**, 2924–2930 (1985).
21. Brower, M. *et al.* Growth of cell lines and clinical specimens of human non-small cell lung cancer in a serum-free defined medium. *Cancer Res.* **46**, 798–806 (1986).
22. Linnoila, R. I. Spectrum of neuroendocrine differentiation in lung cancer cell lines featured by cytomorphology, markers, and their corresponding tumors. *J. Cell. Biochem.* **24** (suppl.), 92–106 (1996).
23. Falco, J. P. *et al.* v-rasH induces non-small cell phenotype, with associated growth factors and receptors, in a small cell lung cancer cell line. *J. Clin. Invest.* **85**, 1740–1745 (1990).
24. H. C. Chen *et al.* Conservation of the *Drosophila* lateral inhibition pathway in human lung cancer: a hairy-related protein (HES-1) directly represses *achaete-scute* homolog-1 expression. *Proc. Natl Acad. Sci. USA* (in the press).
25. Jarriault, S. *et al.* Signalling downstream of activated mammalian notch. *Nature* **377**, 355–358 (1995).
26. Ishibashi, M. *et al.* Targeted disruption of mammalian hairy and enhancer of split homolog-1 (HES-1) leads to up-regulation of neural helix-loop-helix factors, premature neurogenesis, and severe neural tube defects. *Genes Dev.* **9**, 3136–3148 (1995).
27. Gomez-Skarmeta, J. L. *et al.* Araucan and caupolican, two members of the novel iroquois complex, encode homeoproteins that control proneural and vein-forming genes. *Cell* **85**, 95–105 (1996).
28. Nelkin, B. D. *et al.* Transcription factor levels in medullary thyroid carcinoma cells differentiated by Harvey ras oncogene: c-jun is increased. *Biochem. Biophys. Res. Commun.* **170**, 140–146 (1990).
29. Broers, J. L. V. *et al.* Expression of c-myc in progenitor cells of the bronchopulmonary epithelium and in a large number of non-small cell lung cancers. *Am. J. Resp. Cell Mol. Biol.* **9**, 33–43 (1993).
30. Lewis, J. G. *et al.* A serum-resistant cytofectin for cellular delivery of antisense oligodeoxynucleotides and plasmid DNA. *Proc. Natl Acad. Sci. USA* **93**, 3176–3181 (1996).

Acknowledgements. H.J.K.v.d.V. was a visiting research assistant of the National Fund for Scientific Research, Belgium. We thank F. Guillemot and A. Joyner for the MASH1⁺ mouse strain and for whole mouse embryo sections hybridized with MASH1; R. Wagner for GS2888 cytofectin; A. F. Gazdar for cell lines; E. Gabrielson for a human fetal lung specimen; S. M. Steinberg for statistical consultation; and S. Jensen, T. Bunnag and K. Wieman for technical assistance. Animal care procedures were approved by the Johns Hopkins Animal Care and Use Committee. Supported in part by grants from the NIH, the American Cancer Society and the D. Collen Research Foundation.

Correspondence and requests for materials should be addressed to D.W.B. (e-mail: dball@welchlink.welch.jhu.edu).

Differential activation of transcription factors induced by Ca²⁺ response amplitude and duration

Ricardo E. Dolmetsch*, Richard S. Lewis*†, Christopher C. Goodnow‡§ & James I. Healy†‡§

* Department of Molecular and Cellular Physiology and Neurosciences Program, † Program in Immunology, ‡ Department of Microbiology and Immunology and Howard Hughes Medical Institute, Stanford University School of Medicine, Stanford, California 94305, USA

An increase in the intracellular calcium ion concentration ([Ca²⁺]_i) controls a diverse range of cell functions, including adhesion, motility, gene expression and proliferation^{1,2}. Calcium signalling patterns can occur as single transients, repetitive oscillations or sustained plateaux^{2,3}, but it is not known whether these patterns are responsible for encoding the specificity of cellular responses. We report here that the amplitude and duration of calcium signals in B lymphocytes controls differential activation of the pro-inflammatory transcriptional regulators NF- κ B, c-Jun N-terminal kinase (JNK) and NFAT. NF- κ B and JNK are selectively activated by a large transient [Ca²⁺]_i rise, whereas NFAT is activated by a low, sustained Ca²⁺ plateau. Differential activation results from differences in the Ca²⁺ sensitivities and kinetic behaviour of the three pathways. Our results show how downstream effectors can decode information contained in the amplitude and duration of Ca²⁺ signals, revealing a mechanism by which a multifunctional second messenger such as Ca²⁺ can achieve specificity in signalling to the nucleus.

Calcium signalling is important for the growth, death, differentiation and function of immune cells^{4–6}. Several Ca²⁺-sensitive transcriptional regulators, including NF- κ B^{7,8}, JNK^{9,10} and NFAT^{4,6,11–14}, participate in varying combinations to promote the expression of genes that underlie these responses. These genes encode haematopoietic growth factors such as the interleukins IL-2 and IL-4 and GM-CSF, and inflammatory cytokines such as IL-1, IL-6, IL-8 and tumour-necrosis factor (TNF)^{4,7,10,14}. We examined the Ca²⁺ sensitivity and response dynamics of the three transcriptional regulators in B lymphocytes from mice expressing an immunoglobulin transgene specific for the hen-egg lysozyme (HEL) antigen¹⁵. Acute stimulation of B cells with HEL and phorbol ester evoked a biphasic Ca²⁺ response that consisted of an initial rapid rise of [Ca²⁺]_i from a baseline of 76 ± 3 nM to a peak of $1,367 \pm 33$ nM, which subsequently declined over 10 min to a low sustained plateau of 227 ± 5 nM (means $\pm$ s.e.m., $n = 572$ cells) (Fig. 1a). The time course of NF- κ B activation during this response was assayed by the phosphorylation and degradation of its cytoplasmic inhibitor I κ B α , or by the accumulation of the NF- κ B subunit, RelA, in the nucleus^{7,16,17}. NF- κ B activation was first detectable 4 min after stimulation with HEL and reached completion by 8 min (Fig. 1b). Like NF- κ B activation, the phosphorylation of JNK1 and its nuclear substrate ATF-2 (refs 10, 18) was also first apparent within 4 min and reached a peak by 12 min (Fig. 1c). In contrast, activation of NFATc, as determined by its translocation to the nucleus, was much more rapid, being essentially complete within 1 min of stimulation with antigen (Fig. 1d, e). The activation of each of these pathways was inhibited by the immunosuppressant cyclosporin A (CsA)¹⁹ as shown previously^{8,9,20}.

An important question is whether the spike and plateau phases of the Ca²⁺ response activate distinct transcriptional pathways. Stimulation of the antigen receptor activates Ca²⁺-independent as well as Ca²⁺-dependent signalling pathways^{5,21}; we therefore applied the

Ca^{2+} ionophore ionomycin (in the presence of phorbol ester) to identify the specific impact of each phase of the Ca^{2+} response. The Ca^{2+} spike was generated in isolation by exposing the cells to ionomycin shortly before chelating extracellular Ca^{2+} with EGTA (Fig. 2a, left). Single-cell Ca^{2+} imaging confirmed that the ionomycin-induced transient had about the same magnitude as the HEL-stimulated spike ($1,301 \pm 46 \text{ nM}$ versus $1,367 \pm 33 \text{ nM}$, respectively; mean $\pm$ s.e.m.) and returned to baseline within 3 min. Furthermore, we verified that the Ca^{2+} spikes produced by HEL

or ionomycin are attributable to Ca^{2+} released from internal stores ($\sim 25\%$) and to Ca^{2+} influx ($\sim 75\%$) (data not shown), indicating that ionomycin and HEL mobilize the same sources of Ca^{2+} to generate the spike. As a positive control, ionomycin was applied without EGTA to create a sustained $[\text{Ca}^{2+}]_i$ increase that fully activated all three pathways (Fig. 2a, right). Stimulation with phorbol ester alone did not raise $[\text{Ca}^{2+}]_i$ (data not shown), nor did it activate NFATc, NF- κ B, or JNK1 (Fig. 1b–d, lanes marked P), confirming that all three pathways require elevated $[\text{Ca}^{2+}]_i$ for activation. In the absence of phorbol ester, Ca^{2+} ionophore alone did not activate NF- κ B or JNK (ref. 19, and data not shown); ionomycin did cause nuclear translocation of NFATc but previous studies have shown that NFAT-dependent transcription requires a second signal that can be provided by protein kinase $\text{C}^{4,14}$.

The high- Ca^{2+} spike evoked prolonged activation of NF- κ B and JNK1/ATF-2. I κ B α became phosphorylated and was degraded within 4–8 min, and RelA accumulated in the nucleus with a parallel time course (Fig. 2b, e). The NF- κ B response was comparable when $[\text{Ca}^{2+}]_i$ was raised continuously (Fig. 2b, e). Likewise, JNK1 and ATF-2 phosphorylation reached a stable maximum 8 min after the initial increase in $[\text{Ca}^{2+}]_i$, regardless of whether the rise was transient or sustained (Fig. 2c, e). Activation of both NF- κ B and

Figure 1 Antigen evokes a biphasic $[\text{Ca}^{2+}]_i$ rise that activates the NF- κ B, JNK1 and NFAT pathways. **a**, Purified splenic B lymphocytes expressing a hen egg lysozyme(HEL)-specific immunoglobulin transgene were analysed by digital video microscopy. Fura-2-loaded cells were stimulated with 500 ng ml^{-1} HEL and 5 ng ml^{-1} phorbol 12,13-dibutyrate (PDBU) (bar). Mean responses of 285 cells (left) and of representative single cells (right) are shown. After stimulation, nuclear and cytoplasmic fractions were analysed by western blotting. **b**, NF- κ B activation is demonstrated by the phosphorylation and degradation of cytoplasmic I κ B α and the accumulation of RelA in nuclear lysates. Phosphorylated I κ B α , first detectable at 4 min, has a slightly lower mobility than the dephosphorylated form. **c**, Cytoplasmic JNK1 and nuclear ATF-2 are phosphorylated following stimulation. Phosphorylation of JNK1 and ATF-2 causes a reduction in electrophoretic mobility¹⁸ that is evident at 4–18 min. **d**, NFATc translocates rapidly from the cytoplasm (cyto) to the nucleus (nuc). NFATc in its phosphorylated state is present as several bands migrating with an apparent M_r of 160K–190K (filled arrows). Nuclear NFATc is dephosphorylated and migrates more rapidly (apparent M_r 80K–150K; open arrows). In **b–d**, lane P shows lysates from cells stimulated with 5 ng ml^{-1} PDBU alone for 12 min as a control. Time in min is indicated above each lane and the M_r values (in K) are shown on the right. Filled and open arrows indicate the positions of phosphorylated and unphosphorylated species, respectively. **e**, Time course of I κ B α degradation ($\blacktriangle$), JNK1 phosphorylation ($\blacksquare$) and NFATc translocation ($\bullet$) as determined by densitometry of the immunoblots in **b–d**.

Figure 2 A brief Ca^{2+} spike induces persistent activation of NF- κ B, JNK1 and ATF-2, but transient translocation of NFATc. **a**, Average Ca^{2+} response of >250 cells treated with 950 nM ionomycin and 5 ng ml^{-1} PDBU (black bar). Left, 3 mM EGTA was added 40 s later (white bar) to terminate the spike. **b**, The Ca^{2+} spike (left lanes) and the continuous $[\text{Ca}^{2+}]_i$ rise (right lanes) activate NF- κ B similarly, as shown by I κ B α phosphorylation and degradation and RelA translocation. **c**, JNK1 and ATF-2 phosphorylation are also sustained following an isolated Ca^{2+} spike or a prolonged $[\text{Ca}^{2+}]_i$ increase. **d**, Cytoplasmic NFATc (cyto) is dephosphorylated and enters the nucleus (nuc) but returns to the cytoplasm after $[\text{Ca}^{2+}]_i$ declines to baseline (left lanes). Continuously elevated $[\text{Ca}^{2+}]_i$ maintains NFATc in the nucleus (right lanes). **e**, Time course of I κ B α degradation, JNK1 phosphorylation and NFATc translocation following the $[\text{Ca}^{2+}]_i$ spike (filled symbols) or the sustained $[\text{Ca}^{2+}]_i$ rise (open symbols). Data were from the experiment shown in **a–d**.

JNK1 was sustained for at least 16 min after termination of the $[Ca^{2+}]_i$ spike, indicating that both pathways retain a memory of transient $[Ca^{2+}]_i$ rises that outlasts the Ca^{2+} signal itself. By contrast, the Ca^{2+} spike evoked only a transient nuclear translocation of NFATc, with one half of the nuclear NFATc returning to the cytoplasm in its phosphorylated form 8 min after $[Ca^{2+}]_i$ returned to baseline (Fig. 2d, e). The reversal of nuclear translocation was

caused by termination of the Ca^{2+} spike, because nuclear NFATc persisted in cells with constant elevated $[Ca^{2+}]_i$ (Fig. 2a, d). These results confirm and extend findings showing that NFAT requires a continuous $[Ca^{2+}]_i$ rise to remain localized in the nucleus^{12,13}. They also demonstrate that differences in deactivation kinetics contribute to the differential activation of multiple transcriptional pathways by Ca^{2+} . Similar results were obtained when HEL followed by EGTA was used to generate an isolated Ca^{2+} spike (data not shown).

We next investigated whether the low- Ca^{2+} plateau triggered by antigen (Fig. 1) can evoke selective activation of transcriptional pathways. Small amounts of ionomycin were added stepwise in order to mimic the antigen-induced plateau without creating an initial spike (Fig. 3a). Low concentrations of ionomycin elicit Ca^{2+} influx in lymphocytes predominantly by activating store-operated Ca^{2+} channels that underlie the antigen-evoked $[Ca^{2+}]_i$ plateau⁶; thus ionomycin seems to mimic the Ca^{2+} -mobilizing action of antigen during the plateau as well as the spike. Stimulation of NF- κ B or JNK1 was not detectable in response to the low- Ca^{2+} plateau (Fig. 3b), whereas NFATc was activated by 70% within 12 min, suggesting that the former are less Ca^{2+} responsive than NFATc. To examine the Ca^{2+} sensitivity of each pathway in more detail, we measured the activation of NF- κ B, JNK1, NFATc and NFATp¹⁴ following a 10-min stimulation with increasing concentrations of ionomycin. Consistent with the results shown in Fig. 3, nuclear translocation of NFATc/p was induced by lower concentrations of ionomycin (79 nM; Fig. 4a) than are required to activate I κ B α degradation or JNK1/ATF-2 phosphorylation (400–600 nM; Fig. 4a). Figure 4c shows the translocation of NFATc/p, degradation of I κ B α , and phosphorylation of JNK1 and ATF-2 as a function of the peak $[Ca^{2+}]_i$ evoked by different doses of ionomycin (Fig. 4b). The results identify a range of $[Ca^{2+}]_i$ levels (200–400 nM) that promote NFATc/p translocation without activating NF- κ B or JNK1. In three experiments, the $[Ca^{2+}]_i$ needed to half-maximally translocate NFATc/p (413 ± 114 nM) was substantially lower than the level needed to attain half-maximal degradation of I κ B α (612 ± 121 nM) or phosphorylation of JNK1 and ATF-2

Figure 3 Selective activation of NFATc by a low $[Ca^{2+}]_i$ plateau. **a**, Cells were stimulated in the constant presence of 5 ng ml^{-1} PDBU with 24 nM ionomycin (IO) for 4 min, followed by addition of 37 nM ionomycin. The steady-state $[Ca^{2+}]_i$ plateau produced under these conditions simulates that induced by HEL stimulation (Fig. 1). **b**, Time course of NFATc translocation (●), I κ B α degradation (▲), and JNK1 phosphorylation (■) for the experiment shown in **a**.

Figure 4 Differential Ca^{2+} sensitivity of NFATc/p, I κ B α /RelA and JNK1/ATF-2. **a**, Cells were treated with increasing concentrations of ionomycin in the presence of 5 ng ml^{-1} PDBU and analysed by western blotting 10 min after stimulation. The concentration of ionomycin in nM is indicated above each lane (U, unstimulated). **b**, Average $[Ca^{2+}]_i$ responses of >250 cells stimulated with increasing doses of ionomycin (23, 35, 52 nM, in addition to the concentrations shown in **a**). The same batch of cells and incubation conditions were used as in **a**. **c**, Degradation of I κ B α (+), phosphorylation of JNK1 (●) and ATF-2 (■), and translocation of NFATc (○) and

NFATp (○) are shown as a function of peak $[Ca^{2+}]_i$ for each dose of ionomycin applied in **b**. Peak $[Ca^{2+}]_i$ is most relevant for NF- κ B and JNK1 because they respond in a prolonged fashion to a transient $[Ca^{2+}]_i$ spike (Fig. 2). Because NFATc/p responds reversibly to increases in $[Ca^{2+}]_i$ (Fig. 2), this analysis may slightly underestimate the true Ca^{2+} sensitivity of NFATc/p. The graph identifies a range of $[Ca^{2+}]_i$ that selectively activates NFATc/p without appreciably stimulating NF- κ B or JNK1/ATF-2.

($1,035 \pm 263$ nM; means $\pm$ s.e.m.). These measurements of the Ca^{2+} sensitivity of NFAT translocation agree with the reported sensitivity of an NFAT/AP1-dependent reporter gene^{6,11}. The Ca^{2+} dependence of NF- κ B and JNK was previously inferred only qualitatively from their stimulation by Ca^{2+} ionophore⁹ or from inhibition by drugs like CsA that inhibit calcineurin⁸. Our results indicate that NFATc/p is significantly more Ca^{2+} -sensitive than NF- κ B and JNK1/ATF-2, and that this enhanced Ca^{2+} sensitivity enables NFAT to be selectively activated by the low- Ca^{2+} plateau.

The ability of cells to react appropriately to a wide variety of environmental stimuli requires a high degree of specificity in signalling from the plasma membrane to the nucleus. It is remarkable that such specificity is achieved, given the relatively small number of second messenger pathways that exist. Many mechanisms have been proposed to explain transcriptional selectivity²², but the contribution of different Ca^{2+} signals to specificity has not been previously recognized. Our results show that the amplitude and duration of dynamic Ca^{2+} signals contribute to transcriptional specificity, and that this is a direct consequence of the differing Ca^{2+} sensitivities and kinetic behaviour of NF- κ B, JNK and NFAT. These differences may also predict frequency-specific effects of $[\text{Ca}^{2+}]_i$ oscillations on gene transcription, which have been implicated in neuronal differentiation²³.

Differential signalling by Ca^{2+} is likely to be important in the regulation of immunity and inflammation. In self-tolerant B cells, basal $[\text{Ca}^{2+}]_i$ is moderately raised and NFAT is activated, but acute stimulation with antigen fails to generate a Ca^{2+} spike or activate the NF- κ B or JNK1 pathways¹⁹. Similarly, elevation of basal $[\text{Ca}^{2+}]_i$ and the absence of an antigen-induced Ca^{2+} spike has been reported in T cells rendered anergic by an anti-CD3 monoclonal antibody²⁴ and in T cells that have differentiated to the anti-inflammatory T_H2 state^{24,25}. Low-amplitude Ca^{2+} responses have also been implicated in the induction of T-cell anergy by altered peptide ligands²⁶. Conversely, a high Ca^{2+} spike without a lower plateau is triggered during the inhibition of B-cell responses by the binding of immune complexes to the Fc receptor^{27,28}. Our findings suggest that the functional consequences of these isolated Ca^{2+} spikes and plateaux, as well as of other Ca^{2+} signalling patterns such as oscillations and waves, may arise from the selective activation of transcriptional regulators. The importance of second messenger amplitude and duration in discriminating between different response pathways may prove to be a common theme in many cells. □

Methods

Calcium imaging. For single-cell calcium imaging, purified B cells²⁹ were incubated at 37° with $1 \mu\text{M}$ Fura-2/AM (Molecular Probes) for 15 min in phenol-red-deficient RPMI supplemented with 3% fetal bovine serum (FBS). After loading, cells were washed and maintained at room temperature in a 5% CO_2 atmosphere for up to 1 hour. Cells were plated on clean glass coverslips and imaged at 37°C in phenol-red-deficient RPMI with 1% FBS. Ratiometric imaging was done on the heated stage of a Zeiss Axiovert 35 microscope using a $40\times$ oil-immersion objective (Zeiss Achrostat, NA 1.3), an intensified CCD camera (Hamamatsu) and a Videoprobe image processor (ETM Systems) essentially as described³⁰.

Quantitation of western blots. Cells were stimulated at a density of $\sim 3 \times 10^7$ cells per ml in phenol-red-deficient RPMI with 1% FBS. Stimulations were stopped on ice, centrifuged and resuspended in hypotonic buffer (5 mM NaCl, 20 mM HEPES, pH 7.5) containing protease (2.5 mM PMSE, $40 \mu\text{g ml}^{-1}$ each of aprotinin and leupeptin, 2 mM EDTA) and phosphatase inhibitors (1 mM NaVO_4 , 6 mM pNPP, 10 mM NaF) and 0.4% NP-40. Nuclei were separated by centrifugation at $600g$ at 3°C and cytoplasmic extracts (supernatant) were added to boiling SDS-PAGE reducing sample buffer. Nuclei were rinsed in hypotonic buffer with inhibitors and lysed in boiling sample buffer. For Figs 1–3, chromatin was removed from nuclear extracts by centrifugation at $70,000g$ in a Beckman airfuge. For Fig. 4, chromatin was sheared by passing it repeatedly through a 26-gauge needle. Extracts were resolved by SDS-PAGE, transferred to nitrocellulose and analysed by immunoblot and ECL (Amer-

sham). Anti-NFATc(7A6) and anti-NFATp(4G10G5) were obtained from L. Timmerman and G. Crabtree, anti-I κ B α and anti-RelA were from Santa Cruz Biotechnology, anti-JNK1 was from Pharmingen and Santa Cruz Biotechnology, and anti-ATF was from J. Hoeffler and M. Green. Autoradiograms were analysed with a Molecular Dynamics computing densitometer using the volume-integrate mode. An exposed area of film outside each lane was used as background. Raw density values for NFATc, I κ B α , and RelA were corrected for the loading of each lane; for JNK1 and ATF-2, the density of phosphorylated bands was calculated relative to the summed density of the phosphorylated and unphosphorylated bands. NFATc activation was calculated from the loss of cytoplasmic material (measurements based on accumulation of nuclear material gave similar results). Activation is expressed relative to the values observed in unstimulated (0%) and maximally stimulated (100%) cells. On an absolute scale, maximal NFATc/p translocation ranged from 75–90%, maximal I κ B α degradation was 80–99%, nuclear RelA was induced maximally by fivefold, ATF-2 phosphorylation was 20–30%, and maximum JNK1 phosphorylation ranged from 50–60%.

Received 16 December 1996; accepted 6 March 1997.

- Ghosh, A. & Greenberg, M. E. Calcium signaling in neurons: molecular mechanisms and cellular consequences. *Science* **268**, 239–247 (1995).
- Berridge, M. J. Inositol trisphosphate and calcium signalling. *Nature* **361**, 315–325 (1993).
- Clapham, D. E. Calcium signaling. *Cell* **80**, 259–268 (1995).
- Crabtree, G. R. & Clippstone, N. A. Signal transmission between the plasma membrane and nucleus of T lymphocytes. *Annu. Rev. Biochem.* **63**, 1045–1083 (1994).
- Gold, M. R. & DeFranco, A. L. Biochemistry of B lymphocyte activation. *Adv. Immunol.* **55**, 221–295 (1994).
- Fanger, C. M., Hoth, M., Crabtree, G. R. & Lewis, R. S. Characterization of T cell mutants with defects in capacitative calcium entry: genetic evidence for the physiological roles of CRAC channels. *J. Cell Biol.* **131**, 655–667 (1995).
- Bauerle, P. A. & Henkel, T. Function and activation of NF- κ B in the immune system. *Annu. Rev. Immunol.* **12**, 141–179 (1994).
- Frantz, B. et al. Calcineurin acts in synergy with PMA to inactivate I κ B/MAD3, an inhibitor of NF- κ B. *EMBO J.* **13**, 861–870 (1994).
- Su, B. et al. JNK is involved in signal integration during costimulation of T lymphocytes. *Cell* **77**, 727–736 (1994).
- Karin, M. & Hunter, T. Transcriptional control by protein phosphorylation: signal transmission from the cell surface to the nucleus. *Curr. Biol.* **5**, 747–757 (1995).
- Negulescu, P. A., Shastri, N. & Cahalan, M. D. Intracellular calcium dependence of gene expression in single T lymphocytes. *Proc. Natl Acad. Sci. USA* **91**, 2873–2877 (1994).
- Timmerman, L. A., Clippstone, N. A., Ho, S. N., Northrop, J. P. & Crabtree, G. R. Rapid shuttling of NF-AT in discrimination of Ca^{2+} signals and immunosuppression. *Nature* **383**, 837–840 (1996).
- Shibasaki, F., Price, E. R., Milan, D. & McKeon, F. Role of kinases and the phosphatase calcineurin in the nuclear shuttling of transcription factor NF-AT4. *Nature* **382**, 370–373 (1996).
- Rao, A. NF-AT: a transcription factor required for the coordinate induction of several cytokine genes. *Immunol. Today* **15**, 274–281 (1994).
- Goodnow, C. C. et al. Altered immunoglobulin expression and functional silencing of self-reactive B lymphocytes in transgenic mice. *Nature* **334**, 676–682 (1988).
- DiDonato, J. A., Mercurio, F. & Karin, M. Phosphorylation of I- κ B α precedes but is not sufficient for its dissociation from NF- κ B. *Mol. Cell Biol.* **15**, 1302–1311 (1995).
- Verma, I. M., Stevenson, J. K., Schwarz, E. M., Van Antwerp, D. & Miyamoto, S. Rel/NF- κ B/I κ B family: intimate tales of association and dissociation. *Genes Dev.* **9**, 2723–2735 (1995).
- Gupta, S., Campbell, D., Dérjard, B. & Davis, R. J. Transcription factor ATF-2 regulation by the JNK signal transduction pathway. *Science* **267**, 389–393 (1995).
- Healy, J. I. et al. Different nuclear signals are activated by the B cell receptor during positive versus negative signaling. *Immunity* (in press).
- Flanagan, W. M., Cortés, B., Bram, R. J. & Crabtree, G. R. Nuclear association of a T-cell transcription factor blocked by FK506 and cyclosporin A. *Nature* **352**, 803–807 (1991).
- Cambier, J. C., Pleiman, C. M. & Clark, M. R. Signal transduction by the B cell antigen receptor and its coreceptors. *Annu. Rev. Immunol.* **12**, 457–486 (1994).
- Ernst, P. & Smale, S. T. Combinatorial regulation of transcription 1: general aspects of transcriptional control. *Immunity* **2**, 311–319 (1995).
- Gu, X. & Spitzer, N. C. Distinct aspects of neuronal differentiation encoded by frequency of spontaneous Ca^{2+} transients. *Nature* **375**, 784–787 (1995).
- Gajewski, T. F., Lancki, D. W., Stack, R. & Fitch, F. W. "Anergy" of T_H0 helper T lymphocytes induces downregulation of T_H1 characteristics and a transition to a T_H2 -like phenotype. *J. Exp. Med.* **179**, 481–491 (1994).
- Gajewski, T. F., Schell, S. R. & Fitch, F. W. Evidence implicating utilization of different T cell receptor-associated signaling pathways by T_H1 and T_H2 clones. *J. Immunol.* **144**, 4110–4120 (1990).
- Sloan-Lancaster, J., Steinberg, T. H. & Allen, P. M. Selective activation of the calcium signaling pathway by altered peptide ligands. *J. Exp. Med.* **184**, 1525–1530 (1996).
- Wilson, H. A. et al. The B lymphocyte calcium response to anti-Ig is diminished by membrane immunoglobulin cross-linkage to the Fc gamma receptor. *J. Immunol.* **138**, 1712–1718 (1987).
- Choquet, D., Partiseti, M., Amigorena, S., Bonnerot, C. & Fridman, W. Cross-linking of IgG receptors inhibits membrane immunoglobulin-stimulated calcium influx in B lymphocytes. *J. Biol. Chem.* **268**, 355–363 (1993).
- Cyster, J. et al. Regulation of B-lymphocyte negative and positive selection by tyrosine phosphatase CD45. *Nature* **381**, 325–328 (1996).
- Dolmetsch, R. & Lewis, R. S. Signaling between intracellular Ca^{2+} stores and depletion-activated Ca^{2+} channels generates $[\text{Ca}^{2+}]_i$ oscillations in T lymphocytes. *J. Gen. Physiol.* **103**, 365–388 (1994).

Acknowledgements. We thank L. Timmerman, G. Crabtree, J. Hoeffler, and M. Green for the generous gifts of antibodies, and M. Hoth, A. Zweifach, and R. Tsien for comments on an earlier version of the manuscript. R.E.D. is supported by an American Heart Association Fellowship, R.S.L. is supported by the NIH, C.C.G. is an investigator of the Howard Hughes Medical Institute, and J.I.H. is a Beckman Scholar supported by the NIGMS Medical Scientist Training Program.

Correspondence and requests for materials should be addressed to R.S.L. (e-mail: rslweis@leland.stanford.edu).

synthesized by PCR. DNase I footprinting reactions were processed as described²⁴.

Reporter gene construction. Substitution of a single *Bgl*II site for the quadrant enhancer Mad binding-site sequence GCTGCCGTCGCG was done by assembling two fragments generated by PCR with *Pfu* polymerase. Expansion of the single *Bgl*II site to two tandem sites matching the Q^m sequence was done by insertion of an oligonucleotide containing *Bbs*I internal sites that allowed for subsequent cleavage and ligation to generate the tandem *Bgl*II sites. Both sequences were confirmed by sequencing and ligated into the *Kpn*I/*Not*I sites of the Hsp *lacZ*-CaSpeR *Drosophila* transformation vector for the generation of transgenic lines²⁴.

Received 20 January; accepted 21 May 1997.

- Graff, J. M., Bansal, A. & Melton, D. A. *Xenopus* Mad proteins transduce distinct subsets of signals for the TGF β superfamily. *Cell* **85**, 479–487 (1996).
- Hoodless, P. A. *et al.* MADR1, a MAD-related protein that functions in BMP2 signaling pathways. *Cell* **85**, 489–500 (1996).
- Liu, F. *et al.* A human Mad protein acting as a BMP-regulated transcriptional activator. *Nature* **381**, 622–623 (1996).
- Eppert, K. *et al.* MADR2 maps to 18q21 and encodes a TGF β -regulated MAD-related protein that is functionally mutated in colorectal carcinoma. *Cell* **86**, 543–552 (1996).
- Zhang, Y., Feng, X.-H., Wu, R.-Y. & Derynck, R. Receptor-associated Mad homologues synergize as effectors of the TGF- β response. *Nature* **383**, 168–172 (1996).
- Lagna, G., Hata, A., Hemmati-Brivnlou, A. & Massagué, J. Partnership between DPC4 and SMAD proteins in TGF- β signalling pathways. *Nature* **383**, 832–836 (1996).
- Savage, C. *et al.* *C. elegans* genes *smad-2*, *smad-3*, and *smad-4* define a conserved family of TGF- β pathway components. *Proc. Natl Acad. Sci. USA* **93**, 790–794 (1996).
- Macias-Silva, M. *et al.* MADR2 is a substrate of the TGF- β receptor and its phosphorylation is required for nuclear accumulation and signaling. *Cell* **87**, 1215–1224 (1996).
- Derynck, R. & Zhang, Y. Intracellular signalling: The Mad way to do it. *Curr. Biol.* **6**, 1226–1229 (1996).
- Chen, X., Rubock, M. J. & Whitman, M. A transcriptional partner for MAD proteins in TGF- β signalling. *Nature* **383**, 691–696 (1996).
- Rafferty, L., Twombly, V., Wharton, K. & Gelbart, W. Genetic screens to identify elements of the decapentaplegic pathway in *Drosophila*. *Genetics* **139**, 241–254 (1995).
- Sekelsky, J., Newfeld, S., Rafferty, L., Chartoff, E. & Gelbart, W. Genetic characterization and cloning of *Mothers against dpp*, a gene required for decapentaplegic function in *Drosophila melanogaster*. *Genetics* **139**, 1347–1358 (1995).
- Newfeld, S. J., Chartoff, E. H., Graff, J. M., Melton, D. A. & Gelbart, W. M. *Mothers against dpp* encodes a conserved cytoplasmic protein required in DPP/TGF- β responsive cells. *Development* **122**, 2099–2108 (1996).
- Wiersdorff, V., Lecuit, T., Cohen, S. M. & Mlodzik, M. *Mad* acts downstream of Dpp receptors, revealing a differential requirement for *dpp* signaling in initiation and propagation of morphogenesis in the *Drosophila* eye. *Development* **122**, 2153–2162 (1996).
- Blair, S. Compartments and appendage development in *Drosophila*. *BioEssays* **17**, 229–309 (1995).
- Basler, K. & Struhl, G. Compartment boundaries and the control of *Drosophila* limb pattern by hedgehog protein. *Nature* **368**, 208–214 (1994).
- Tabata, T. & Kornberg, T. Hedgehog is a signalling protein with a key role in patterning *Drosophila* imaginal discs. *Cell* **76**, 89–102 (1994).
- Zecca, M., Basler, K. & Struhl, G. Sequential organizing activities of engrailed, hedgehog and decapentaplegic in the *Drosophila* wing. *Development* **121**, 2265–2278 (1995).
- Ingham, P. W. & Fietz, M. J. Quantitative effects of hedgehog and decapentaplegic activity on the patterning of the *Drosophila* wing. *Curr. Biol.* **5**, 432–440 (1995).
- Posakony, L., Rafferty, L. & Gelbart, W. Wing formation in *Drosophila melanogaster* requires decapentaplegic gene function along the anterior–posterior compartment boundary. *Mech. Dev.* **33**, 69–82 (1991).
- Capdevilla, J. & Guerrero, I. Targeted expression of the signalling molecule decapentaplegic induces pattern duplications and growth alterations in *Drosophila* wings. *EMBO J.* **13**, 4459–4468 (1994).
- Nellen, D., Burke, R., Struhl, G. & Basler, K. Direct and long-range action of a Dpp morphogen gradient. *Cell* **85**, 357–368 (1996).
- Lecuit, T. *et al.* Two distinct mechanisms for long-range patterning by Decapentaplegic in the *Drosophila* wing. *Nature* **381**, 387–393 (1996).
- Kim, J. *et al.* Integration of positional signals and regulation of wing formation and identity by *Drosophila* vestigial gene. *Nature* **382**, 133–138 (1996).
- de Celis, J. F., Barrio, R. & Kafatos, F. C. A gene complex acting downstream of *dpp* in *Drosophila* wing morphogenesis. *Nature* **381**, 421–424 (1996).
- Grimm, S. & Pflugfelder, G. O. Control of the gene optomotor-blind in *Drosophila* wing development by decapentaplegic and wingless. *Science* **271**, 1601–1604 (1996).
- Chouinard, S. & Kaufman, T. C. Control of expression of the homeotic *labial* (*lab*) locus of *Drosophila melanogaster*: evidence for both positive and negative autogenous regulation. *Development* **113**, 1267–1280 (1991).
- Tremml, G. & Bienz, M. Induction of *labial* expression in the *Drosophila* endoderm: response elements for *dpp* signalling and for autoregulation. *Development* **116**, 447–456 (1992).
- Thüringer, F., Cohen, S. M. & Bienz, M. Dissection of an indirect autoregulatory response of a homeotic *Drosophila* gene. *EMBO J.* **12**, 2419–2430 (1993).
- Eresh, S., Riese, J., Jackson, D. B., Bohmann, D. & Bienz, M. A CREB-binding site as a target for decapentaplegic signalling during *Drosophila* endoderm induction. *EMBO J.* **16**, 2014–2022 (1997).
- Ausubel, F. M. *et al.* (eds) *Current Protocols in Molecular Biology* supplement 28, pp. 16.6.1–16.7.7 (Wiley, New York, 1994).

Acknowledgements. We thank K. Vorwerk for technical assistance; M. Hoffmann for discussion; Y. J. Chen for the MAD cDNA clone and help with sequencing; V. Wiersdorff, M. Mlodzik and S. Cohen for the Mad¹⁻² stock; S. Paddock for help with microscopy; T. Kaufman for the labial endodermal enhancer; M. Hoffmann and G. Halder for comments on the manuscript; and J. Wilson for manuscript preparation. K. J. was supported by a genetics predoctoral training grant and the Cremer Scholarship Fund, and part of this work was supported by the Howard Hughes Medical Institute (S.B.C.).

Correspondence and requests for materials should be addressed to A.L. (e-mail: alaughon@facstaff.wisc.edu).

errata

Differential activation of transcription factors induced by Ca²⁺ response amplitude and duration

Ricardo E. Dolmetsch, Richard S. Lewis, Christopher C. Goodnow & James I. Healy

Nature **386**, 855–858 (1997)

The graphs in Fig. 2e were accidentally swapped around during the production process. The correct versions are shown here. □

Large-scale tectonic deformation inferred from small earthquakes

Falk Amelung & Geoffrey King

Nature **386**, 702–705 (1997)

Owing to an error in the production process, a length scale was omitted from Fig. 2b and the shading in Fig. 2d did not show up. The corrected figure is shown here. □

- Groisman, P. Ya, Karl, T. R. & Knight, T. W. Observed impact of snow cover on the heat balance and the rise of continental spring temperatures. *Science* **263**, 198–200 (1994).
- Tucker, C. J. in *Advances in the Use of NOAA AVHRR Data for Land Applications* (ed. D'Souza, D.) 1–19 (European Economic Union Press, Brussels, 1995).
- James, M. E. & Kalluri, S. N. V. The Pathfinder AVHRR land data set: an improved coarse-resolution data set for terrestrial monitoring. *Int. J. Remote Sens.* **15**, 3347–3364 (1994).
- Tucker, C. J., Fung, I. Y., Keeling, C. D. & Gammon, R. H. Relationship between atmospheric CO₂ variations and a satellite-derived vegetation index. *Nature* **319**, 195–199 (1986).
- Asrar, G., Fuchs, M., Kanemasu, E. T. & Hatfield, J. L. Estimating absorbed photosynthetic radiation and leaf area index from spectral reflectance in wheat. *Agron. J.* **76**, 300–306 (1984).
- Myneni, R. B., Hall, F. G., Sellers, P. J. & Marshak, A. L. the interpretation of spectral vegetation indexes. *IEEE Trans. Geosci. Remote Sens.* **33**, 481–486 (1995).
- Tucker, C. J., Newcomb, W. W. & Dregne, A. E. AVHRR data sets for determination of desert spatial extent. *Int. J. Remote Sens.* **15**, 3547–3566 (1994).
- Rao, C. R. N. & Chen, J. Inter-satellite calibration linkages for the visible and near-infrared channels of the advanced Very High Resolution Radiometer on the NOAA-7, -9, and -11 spacecraft. *Int. J. Remote Sens.* **16**, 1931–1942 (1995).
- Los, S. O. Calibration adjustment of the NOAA AVHRR Normalized Difference Vegetation Index without recourse to component channel 1 and 2 data. *Int. J. Remote Sens.* **14**, 1907–1917 (1993).
- Holben, B. N. Characteristics of maximum value composite images for temporal AVHRR data. *Int. J. Remote Sens.* **7**, 1417–1437 (1986).
- Myneni, R. B., Tucker, C. J., Asrar, G., Keeling, C. D. & Nemani, R. R. Increased vegetation greenness amplitude and growing season duration in northern high latitudes inferred from satellite-sensed vegetation index data from 1981–91. *NASA Tech. Memo. 104638* (NASA Goddard Space Flight Center, Greenbelt, MD, 1996).
- Myneni, R. B., Los, S. & Tucker, C. J. Satellite-based identification of linked vegetation index and sea surface temperature anomaly areas from 1982–1990 for Africa, Australia and South America. *Geophys. Res. Lett.* **23**, 729–732 (1996).
- Keeling, C. D., Whorf, T. P., Wahlen, M. & van der Plicht, J. Interannual extremes in the rate of rise of atmospheric carbon dioxide since 1980. *Nature* **375**, 666–670 (1995).
- Heimann, M., Keeling, C. D. & Tucker, C. J. in *Aspects of Climate Variability in the Pacific and Western Americas* (ed. Peterson, D. H.) 277–303 (Geophys. Monog. Ser., Am. Geophys. Union, Washington DC, 1989).
- Kauppi, P. E., Mielikainen, K. & Kuusela, K. Biomass and carbon budget of European forests from 1971–1990. *Science* **256**, 70–74 (1992).
- Jacoby, G. C., D'Arrigo, R. D. & Davaajamts, T. Mongolian tree rings and 20th-century warming. *Science* **273**, 771–773 (1996).
- Houghton, J. T. et al. (eds) *Climate Change 1995* 1–365 (Cambridge Univ. Press, 1995).
- Jones, P. D., Wigley, T. M. L. & Briffa, K. R. in *Trends '93: A Compendium of Data on Global Change* (eds Boden, T. A., Kaiser, D. P., Sepanski, R. J. & Stoss, F. W.) (ORNL/CDIAC-65, Oak Ridge, TN, 1994).
- Piper, S. C. & Stewart, E. F. A gridded global data set of daily temperature and precipitation for terrestrial biosphere modelling. *Glob. Biogeochem. Cycles* **10**, 757–782 (1996).

Acknowledgements. We thank S. C. Piper and E. F. Stewart for analysis of the station temperature data, S. C. Piper and T. P. Whorf for discussions, and S. Los for help in the calibration of GIMMS NDVI data. This work was supported by the Office of Mission to Planet Earth of NASA. E. F. Stewart's collaboration was made possible by funds from the Electric Power Research Institute and the USNSF.

Correspondence should be addressed to R.B.M. (e-mail: rmyneni@crsa.bu.edu).

Large-scale tectonic deformation inferred from small earthquakes

Falk Amelung* & Geoffrey King

Institut de Physique du Globe, 5, Rue Rene Descartes, F-67084 Strasbourg, France

It is a long-standing question whether the focal mechanisms of small earthquakes can be used to provide information about tectonic deformation on a regional scale. Here we address this question by using a 28-year record of seismicity in the San Francisco Bay area to compare the strain released by small earthquakes with geological, geodetic and plate-tectonic measurements of deformation in this region. We show that on a small spatial scale, the strain released by small earthquakes is closely related to specific geological features. But when averaged over a regional scale, strain release more closely follows the regional pattern of tectonic deformation: this relationship holds for all but the largest earthquakes, indicating that the earthquake strain is self-similar^{1,2} over a broad range of earthquake magnitudes. The lack of self-similarity observed for the largest earthquakes suggests that the time interval studied is not large enough to sample a complete set of events—the fault with the highest probability³ for hosting one such missing event is the Hayward fault.

* Present addresses: Department of Geophysics, Stanford University, Stanford, California 94305, USA (F.A.); Institut de Physique du Globe, Laboratoire de Tectonique, Tour 24, BP89, 4 Place Jussieu, 75252 Paris, France (G.K.).

Following the formulation by Kostrov⁴, the seismic strain of a block in the Earth's crust can be obtained from the geometric moment tensors

$$\epsilon_{ij} = \frac{1}{2V} \sum_{n=1}^N \mathcal{M}_{ij}^{(n)} \quad (1)$$

where $\mathcal{M}_{ij}^{(n)}$ is the geometric moment tensor of the n th earthquake, N is the number of earthquakes and V is the volume of the block. The geometric moment tensor of an earthquake is given by $\mathcal{M}_{ij} = \int_A u dA (u_i n_j + u_j n_i)$, where A is the surface of the fault, dA is a surface element, u is the slip in the earthquake, and u_i and n_i are the unit slip and fault-normal vectors, respectively. The geometric moment tensor $\mathcal{M}_{ij}$, referred to by Ben-Menahem and Singh⁵ as “potency”, is related to the seismic moment tensor M_{ij} used in seismology by $M_{ij} = \mu \mathcal{M}_{ij}$ (ref. 6), where μ is the shear modulus.

The strain in a local region, $\epsilon_{ij}^{(loc)}$, is related to the strain in a larger region, $\epsilon_{ij}^{(reg)}$, by a fourth-order tensor that is a complicated function of time and space. Here we consider special cases in which they are assumed to be linearly related by

$$\epsilon_{ij}^{(loc)} = c \epsilon_{ij}^{(reg)} \quad (2)$$

where c is the amplitude of the local deformation, that is, the shape and the orientation of the local and regional strain tensors are identical but their amplitudes can be different. For homogeneous deformation, $c = 1$. If seismic strain is self-similar and if $\epsilon_{ij}^{(loc)}$ is the seismic strain by earthquakes in one scale, then equation (2) will apply at any scale with different values for c . Some examples are

Figure 1 Homogeneous and inhomogeneous deformation in four different regions subject to pure shear. **a**, Boundary conditions and imposed strain. **b**, The strain tensor is represented by two-dimensional strain rosettes for the horizontal components. White lobes indicate directions of contraction, black lobes indicate directions of extensions. **c**, Homogeneous deformation. **d**, Inhomogeneous deformation due to an individual shear fault. **e**, Inhomogeneous deformation due to two faults. **f**, Quasi-homogeneous deformation due to many shear faults with a variety of orientations.

illustrated in Fig. 1. The (applied) regional strain is pure shear (Fig. 1a). We refer to the orientation and shape of a strain tensor as the strain pattern and use strain rosettes for the representation (Fig. 1b). In the absence of faults the strain is homogeneous and the strain pattern in a local region is identical to the regional strain (Fig. 1c). The strain due to slip on a single fault is in general different from the regional strain (Fig. 1d). Two shear faults are, in principle, sufficient to accommodate any regional shear although strain concentrations can be large (Fig. 1e). The strain pattern in a region that contains

many faults with a variety of orientations and homogeneous spatial distribution tends to equal the regional strain (Fig. 1f). Note that Fig. 1f is identical to Fig. 1c if the faults are small compared to the dimension of the region.

The larger San Francisco Bay area (Fig. 3a) is part of the plate boundary zone between the Pacific and North American plates. The regional deformation is primarily simple shear. South of 37° N nearly all the transform motion is accommodated on the San Andreas fault and the motion is mainly aseismic (creeping section).

Table 1 Seismic deformation in earthquake regions

Region	Direction of principal contraction	principal extensional	Number of events summed
		principal contraction	
Golden Gate	N3°W	2.9	58
San Francisco Peninsula	N31°W	0.29	170
Loma Prieta	N16°W	0.57	885
Hollister	N3°W	0.91	7,117
Calaveras fault	N14°W	0.99	2,303
Hayward fault	N12°W	0.88	341
Livermore	N33°W	1	1,239
San Gregorio fault	N20°W	0.59	75
North Bay	N28°W	0.77	98
Diablo Range	N26°W	0.77	182

Shown here are the directions of principal contraction and ratios between the principal components of the horizontal strain in the regions from Fig. 3, for all events with magnitude ≤ 3.33 .

Figure 2 a, Two-dimensional strain rosettes representing the horizontal seismic strain patterns due to the earthquakes in Fig. 3 in magnitude ranges for which mean fault dimensions differ by factors of 2 (A–J). The numbers of events summed are shown next to the strain rosettes. The fault plane solutions are calculated from the first motion polarities using the computer program FPFIT³². The geometric moments are calculated for the local or duration magnitude, M , using the empirical relations $\log M_0 = 1.5M - 1.5$ for $M > 3.33$ and $\log M_0 = 1.2M - 0.5$ for $M \leq 3.33$ (modified from Bakun's³³ relations for the seismic moment using $\mu = 30$ GPa). In this study only "good" events satisfying some quality criteria are used; with uncertainties in strike, dip and rake less than 30°, 45° and 45°, and with magnitude ≥ 0.3 , depth ≥ 2 km, r.m.s. ≤ 0.3 s; horizontal location error ≤ 2.5 km; vertical location error ≤ 5 km; number of first motions ≥ 30 ; azimuthal gap $\leq 180^\circ$ (see ref. 32 for explanation of these parameters)

$\% \text{rt} \leq 0.4$; $\text{sdr} \geq 0.4$; $\text{fit} \leq 0.4$. **b**, Summed geometric moment within the magnitude intervals. The fault length is for circular faults and displacement-to-length ratios of 3×10^{-4} . The abscissa is not linear in magnitude because of two different moment–magnitude relations. The slip rate contribution is averaged over the observational period of 28 years and assumes that the events fall on a 250-km-long and 15-km-deep plate boundary fault. **c**, Predicted strain pattern for an orientation of the Pacific–North American plate boundary of N38°W, and relative motion of the Pacific plate in N34°W. **d**, Mean geodetic strain pattern from trilateration data¹⁶. The contributions of the San Francisco Bay network and of the Monterey network are summed. It is assumed that they cover three-quarters and one-quarter of the region studied, respectively. The strain rosettes in **c** and **d** represent 5% contraction perpendicular to N38°W.

Figure 3 **a**, Location map showing the seismicity ($M > 1.0$) registered by the Northern Californian Seismic Network (NCSN) between January 1968 and December 1995 in the broader San Francisco Bay area. This study uses 15% of the 95,000 registered events with well constrained moment tensors. **b**, Two-

dimensional strain rosettes representing the horizontal strain due to earthquakes in the magnitude ranges defined in Fig. 2a (denoted by letters A–G). Only strain rosettes resulting from summing >5 events are shown. The numbers of events summed are shown next to the strain rosettes.

Further north, two-thirds of the deformation⁷ occurs on the San Andreas fault in occasional large earthquakes such as the 400-km-long rupture in 1906 or in smaller earthquakes that rupture fault segments 10–40 km long³. The remaining deformation is accommodated on subparallel strike-slip faults. The Hayward fault, for example, has been the source of two earthquakes with magnitudes (M) between 6.5 and 7 in 1836 and 1868 (ref. 8), and the Calaveras fault has ruptured in a sequence of earthquakes with $M > 5.5$ beginning in 1979; the $M = 6.2$ 1984 Morgan Hill earthquake was the largest^{9,10}. The recurrence times of 1906-type San Francisco earthquakes, of events on the Hayward fault and the Calaveras fault earthquakes are of the order of 250 years, 150 years and 80 years, respectively³. The seismic data covers a 28-year time period, which is nearly one-third of the shortest seismic cycle.

The seismic deformation of the San Francisco Bay area for different magnitude intervals is shown in Fig. 2a. The geometric moment tensors are derived from the fault plane solution and the magnitude for each event. The magnitude ranges have been chosen such that the seismic moments vary from one range to the other by a factor of 8. For events that are individually self-similar this corresponds to a factor of 2 change in mean fault dimensions between ranges^{11–13}. It can be seen that the strain patterns in the magnitude intervals below magnitude 3.94 are similar and can be interpreted as right-lateral shear across N38°W and 5% of shortening in perpendicular direction. The strain patterns differ only for the largest earthquakes where few events have been summed. The similarity of the strain patterns, together with the fact that the events occurred in different parts of the Bay area on faults with different orientations, suggests that the small faults slipped such that the ensemble resulted in a homogeneous reduction of elastic strain. If that is true, equation (2) is satisfied and the strain accommodated by small earthquakes

represents the applied deformation. The similarity of the strain patterns also indicates the self-similarity of earthquake strain. As many other aspects of rupture of rock are self-similar phenomena, such a self-similarity is expected to occur. Although the strain patterns for different magnitude ranges are similar, the strain amplitudes vary strongly. This can be seen in Fig. 2b where the summed release of geometric moment is shown. The nearly linear increase of the summed moment with the fault length is characteristic for tectonic environments with a b -value (the slope in the Gutenberg–Richter frequency of earthquake-occurrence relation) of ~ 1.0 (refs 14, 15). Note that the summed moment released by all the earthquakes and that by the 'good' earthquakes (with source parameters satisfying some quality criteria) are similar except in the two lowest magnitude intervals. The appropriateness of the quality criteria has been checked by examining the stability of all the seismic strain patterns obtained in this study using more and less strict criteria.

The seismic strain pattern can be compared with the strain pattern from plate motion and from geodetic¹⁶ information (Fig. 2c, d). The plate motion strain pattern is calculated from the relative velocity between the plates and the plate boundary direction, along which linear deformation is assumed not to occur¹⁷. According to plate motion models, the Pacific plate at the latitude of San Francisco moves N34°W with respect to North America^{18,19}. For a plate boundary direction of N34°W, the San Francisco Bay area would shear right-laterally at N34°W. The average strike of the major strike-slip faults, however, is more westerly. For a plate boundary direction of N38°W the predicted mean deformation contains some boundary normal contraction.

The good agreement between the seismic strain patterns in the lower magnitude intervals ($M = 0.33$ – 3.94), and the strain patterns

from the plate motion and from geodesy clearly shows that the deformation by the small earthquakes is in direction identical to the applied deformation, that is, the seismic (local) deformation and the applied (regional) deformation are linearly related by equation (2). The condition for this to be true appears to be that a sufficient number of earthquakes with comparable size and on faults with a variety of orientations are summed. If not, the deformation will be different because some event types are missing. This is the case for the higher magnitude intervals. In the $M = 3.94$ – 6.34 range, some earthquakes with a significant reverse component are absent. In the $M = 6.34$ – 7.00 range, a strike-slip event is missing. The fault with the highest probability³ for such an earthquake is the Hayward fault.

We note at this stage that the seismic strain pattern, the predicted strain pattern, as well as the geodetic strain pattern imply that, in the San Francisco Bay area, a shortening of $\sim 5\%$ of the shear deformation (2 mm yr^{-1}) occurs normal to the plate boundary (assumed to be oriented $N38^\circ W$). This is indicated by various geological and geophysical evidence^{20–23}, but has been questioned because the geodetic data has been interpreted differently^{16,24}. (Those studies examined the geodetic data for shortening perpendicular to the plate motion direction ($N34^\circ W$) and not perpendicular to the plate boundary direction ($N38^\circ W$).)

So far this study has been concerned with the overall deformation across the entire San Francisco Bay area (Fig. 3a). Figure 3b shows that within the Bay area the seismic deformation varies locally (see also Table 1). In almost all local regions the strain patterns are independent of earthquake size. A comparison with the geodetic strain pattern in these small areas is difficult because the geodetic data lacks sufficient resolution²⁵. The seismic strain patterns, however, are associated with local tectonics and can partly be predicted from the geometry of the major strike-slip faults. In regions where they are oriented in the plate motion direction we expect shear, as observed for example in the Calaveras area. In regions where the trend of the faults is more westerly or more easterly we expect net contraction or net extension. This is observed along the San Andreas fault. In the San Francisco Peninsula, the Loma Prieta and the Hollister regions (although less prominent there), where the fault strikes more westerly than the plate motion, the seismic strain is contractional. Around the Golden Gate, where the fault strikes more easterly, the seismic strain is extensional. These strain patterns are compatible with the uplift of the southern Santa Cruz Mountains and the subsidence of the Golden Gate area, respectively²⁵.

Earthquake focal mechanisms are commonly interpreted with respect to the state of stress in the Earth's crust^{26–28}. Here we have been concerned with the state of strain. In homogeneous, isotropic material the elastic strain results in a stress state with identical principal directions. In the Earth's brittle crust, however, they should be different because the strain localizes on fractures or faults on a range of scales. In determining strain by summing over many events and over a large period of time we average over strain heterogeneities and reproduce the applied deformation. Stress inversion assumes that the earthquakes are driven by a single stress tensor invariant in space and time, and if the same strain and stress directions are obtained from focal mechanism data this is fortuitous. In the Loma Prieta region, for example, the strain directions differ from stress directions previously reported^{29–31}, possibly because the hypothesis of a spatially and temporally uniform stress tensor is not met. □

Received 2 August 1996; accepted 21 February 1997.

1. Mandelbrot, B. B. *The Fractal Geometry of Nature* (Freeman, San Francisco, 1983).
2. Turcotte, D. L. *Fractals and Chaos in Geology and Geophysics* (Cambridge Univ. Press, Cambridge, 1992).
3. Working Group on Californian Earthquake Probabilities *Probabilities of Large Earthquakes in the San Francisco Bay Region, California* (Circ. 1053, US Geol. Survey, 1990).
4. Kostrov, V. V. *Earth Phys.* **1**, 23–40 (1974).
5. Ben-Menahem, A. & Singh, J. *Seismic Waves and Sources* (Springer, New York, 1981).
6. King, G. C. P. *Phil. Trans. R. Soc. Lond. A* **288**, 197–212 (1978).
7. Kelson, K. I., Lettis, W. R. & Lisowski, M. *Proc. 2nd Conf. on Earthquake Hazards in the Eastern San*

Francisco Bay Area 31–38 (Spec. Publ. 113, California Dept. of Conservation, Div. of Mines and Geology, 1992).

8. Ellsworth, W. *Prof. Pap. US Geol. Surv.* **1515**, 153–181 (1990).
9. Oppenheimer, D. H., Reasenber, P. A. & Simpson, R. W. *J. Geophys. Res.* **93**, 9007–9026 (1988).
10. Oppenheimer, D. H., Bakun, W. H. & Lindh, A. G. *J. Geophys. Res.* **95**, 8483–8489 (1990).
11. Bakun, W. H., King, G. C. P. & Cockerham, R. S. in *Earthquake Source Mechanics* 195–207 (Geophys. Monogr. 37, (Maurice Ewing 6), Am. Geophys. Union, 1986).
12. King, G. C. P., Lindh, A. G. & Oppenheimer, D. H. *Geophys. Res. Lett.* **17**, 1449–1452 (1990).
13. King, G. C. P., Oppenheimer, D. & Amelung, F. *Earth Planet. Sci. Lett.* **128**, 55–64 (1994).
14. King, G. C. P. *Pure Appl. Geophys.* **121**, 761–815 (1983).
15. Amelung, F. & King, G. C. P. *Geophys. Res. Lett.* **24**, 507–510 (1997).
16. Lisowski, M., Savage, J. C. & Prescott, W. H. *J. Geophys. Res.* **96**, 8369–8389 (1991).
17. Ekström, G. & England, P. J. *Geophys. Res.* **94**, 10231–10257 (1989).
18. DeMets, C., Gordon, R. G., Argus, D. F. & Stein, S. *Geophys. J. Int.* **101**, 425–478 (1990).
19. DeMets, C., Gordon, R. G., Argus, D. F. & Stein, S. *Geophys. Res. Lett.* **21**, 2191–2194 (1994).
20. Page, B. M. in *Proc. 2nd Conf. on Earthquake Hazards in the Eastern San Francisco Bay Area* 1–7 (Spec. Publ. 113, California Dept. of Conservation, Div. of Mines and Geology, 1992).
21. Page, B. M. & Brocher, T. M. *Geology* **21**, 635–638 (1993).
22. Jones, D. L. *et al. Tectonics* **13**, 561–574 (1994).
23. Zoback, M. D. *et al. Science* **238**, 1105–1111 (1987).
24. Williams, S. D. P. thesis, Univ. Durham (1995).
25. Amelung, F. thesis, Univ. Louis Pasteur, Strasbourg (1996).
26. Gephart, J. W. & D. W. Forsyth *J. Geophys. Res.* **89**, 9305–9320 (1984).
27. Michael, A. J. *J. Geophys. Res.* **89**, 11517–11526 (1984).
28. Rivera, L. & Cisternas, A. *Bull. Seismol. Soc. Am.* **80**, 600–614 (1990).
29. Michael, A. J., Ellsworth, W. L. & Oppenheimer, D. H. *Geophys. Res. Lett.* **17**, 1441–1444 (1990).
30. Beroza, G. C. & Zoback, M. D. *Science* **259**, 210–213 (1993).
31. Zoback, M. D. & Beroza, G. C. *Geology* **21**, 181–185 (1993).
32. Reasenber, P. A. & Oppenheimer, D. H. *US Geol. Surv. Open-File Rep.* **85-739** (1985).
33. Bakun, W. H. *Bull. Seismol. Soc. Am.* **74**, 439–458 (1984).

Acknowledgements. We thank D. Oppenheimer for help with the data. This work was supported by the CNRS. F.A. was supported by the EU EPOCH programme.

Correspondence should be addressed to F.A.

A snake with legs from the marine Cretaceous of the Middle East

Michael W. Caldwell* & Michael S. Y. Lee†

* *Department of Geology, The Field Museum, Roosevelt Road at Lakeshore Drive, Chicago, Illinois 60605, USA and Department of Biological Science, Biological Sciences Center, University of Alberta, Edmonton, Alberta T6G 2E9, Canada*
† *Zoology Building, School of Biological Sciences, The University of Sydney, NSW 2006, Australia*

Although snakes are descended from limbed squamates ('lizards'), all known snakes lack well developed legs and their nearest lizard relatives have yet to be identified^{1–4}. Here we provide compelling evidence that the Cretaceous squamate *Pachyrhachis problematicus*, previously interpreted as a varanoid lizard^{5–7}, is actually a primitive snake with a well developed pelvis and hindlimbs. *Pachyrhachis* is the sister-taxon of all other snakes. The skull exhibits most derived features of modern snakes, and the body is slender and elongated. But unlike other snakes, *Pachyrhachis* retains a well developed sacrum, pelvis and hindlimb (femur, tibia, fibula, tarsals). *Pachyrhachis* was marine, and provides additional support for mosasauroid-snake affinities.

Both specimens of *Pachyrhachis problematicus* were found in limestone quarries at Ein Jabrud (Bed-Meir Formation, lowermost Cenomanian, mid-Cretaceous⁸), 20 km north of Jerusalem, Israel. The first was named *Pachyrhachis problematicus* (H.U.J.-PAL 3569)^{5,6} and the second *Ophiomorphus colberti* (H.U.J.-PAL 3775)⁷. *Ophiomorphus* was pre-occupied, and thus changed to *Estesius*⁹. Some snake-like features were noted in both but it was concluded that they were long-bodied varanoid lizards and probably not closely related to snakes^{6,7}. Our re-study found no significant differences between the two specimens (Figs 1 and 2). The posterior region of *Pachyrhachis* is not preserved, resulting in the reported absence of the hindlimb⁷ (Fig. 3). *Pachyrhachis* is slightly larger and