readable for more than 100 million years in the magnetic, gravitational and topographical characteristics of ridges and their off-spring plates. But for those who depend on direct collection of rock samples for chemical analyses to determine the composition and origins of the rocks, the tape is ruined within tens of kilometres — equivalent to a few hundred thousand years — by sediment cover. A full picture of the processes occurring at mid-ocean ridges requires a '4-D' view of the architecture of the oceanic lithosphere, the cold and hence relatively rigid crust and upper mantle that constitute the tectonic plate. Yet from a chemical perspective, in most circumstances we can only work systematically in one dimension (along the ridge axis at zero age and at the seafloor depth at which the rocks are exposed).

The sequence of rock types to be found on the ocean floor is quite systematic, and each rock carries evidence of its origin in the crust-forming process. The upper mantle is dominated by rocks called peridotites. When plate spreading draws peridotite upwards beneath the ridge, the rock undergoes partial melting. The liquid fraction separates and rises to form the intrusive 'gabbroic' rocks of the lower oceanic crust and the erupted basalts that constitute the upper oceanic crust. The remaining solid material — the residual peridotite — is chemically modified by the loss of the more fusible elements (becoming richer, for example, in chromium relative to aluminium, in titanium relative to zirconium, and in magnesium relative to iron). Some of these residues of melting

Figure 1 Ocean deep, mountain high. Under the waters of the central Atlantic, straddling the axis of the Mid-Atlantic Ridge, is a remarkable geological feature. The Vema transverse ridge is an exposed section of crust and mantle rocks, rising from a depth of more than 5,000 m at its valley floor to only 450 m below the sea surface at its highest point. Mapped by Bonatti *et al.*¹, and viewed here from the north, the ridge is a 20-million-year chronicle of tectonic activity.

eventually end up near the base of the crust and, as they cool, become incorporated into the lithosphere.

Rocks that have undergone melting, together with their residues, are evident elsewhere on Earth, of course — for instance in volcanoes on land. But attempts to use them to investigate Earth's interior are often confounded by the difficulty in distinguishing the source (the composition of the rock

that entered the melting region) from the process that modified it (the extent of melting, the temperatures and pressures involved, and the subsequent segregation, migration, differentiation, and so on). Yet at mid-ocean ridges, the entire crust is constructed in one episode of melting, and so the thickness of the crust provides a key overall constraint on the extent of melting. Given that there is a positive correlation

Astronomy

Sprinkler system

Henize 3-1475 is a planetary nebula — an expanding shell of gas surrounding a star that is entering the last stages of its life (and, in fact, nothing to do with planets at all). The gas shell of such an object may be spherical, dumb-bell-shaped or even completely irregular, but in the case of Henize 3-1475 it is concentrated into two fast-moving jets that emerge with velocities of up to 4 million kilometres per hour. As the system is also rotating, astronomers have nicknamed it the 'garden-sprinkler' nebula.

Angels Riera and colleagues have produced this composite image of Henize 3-1475 using different filters on a wide-field camera that is part of the Hubble Space Telescope. Coupled with observations from

ground-based telescopes, their data reveal that the fast outflow of gas from the central star is not regular. Instead, the gas is ejected in clumps, roughly once every hundred years. How the jets are created is still a mystery, but this clumping effect could be caused by interactions with a nearby companion star, or by magnetic processes in the central star (similar to the 22-year cycle of magnetic activity that creates sunspots on our own star).

The misnomer 'planetary nebula' was coined by William Herschel in 1784: cataloguing these objects, he was struck by how closely they resembled (viewed through a telescope considerably less powerful than Hubble) the planet he had recently discovered, Uranus.

Alison Wright