

Kinetics

# Gaussian statistics in granular gases

A kinetic theory<sup>1,2</sup> of granular media, analogous to that described by Maxwell–Boltzmann statistics for molecular gases and liquids<sup>3,4</sup>, has long been sought in the quest for a continuum theory. Here we present results from a two-layer, vertically vibrated granular system, in which an oscillating plate drives a horizontal layer of heavy grains, which in turn drives an overlying horizontal layer of lighter grains. In the second layer, but not in the first, we find that the velocity distributions are gaussian — a result that is observed for a wide variety of driving and density parameters, indicating that the second layer’s behaviour is much more gas-like than was previously seen in driven granular media. This similarity to molecular gases brings a kinetic theory of granular media a step closer.

Attempts to apply a kinetic theory to granular media have been undermined by their observed non-gaussian velocity distributions<sup>5–12</sup>, which are inconsistent with Maxwell–Boltzmann statistics. We have designed a set-up (Fig. 1a) in which a two-layer granular medium is shaken vertically on a horizontal plate. The plate moves as

$x(t) = A \times \sin(2\pi ft)$ , which corresponds to a dimensionless acceleration of  $\Gamma = (2\pi f)^2/g$  where  $f$  is the oscillation frequency,  $A$  is the oscillation amplitude and  $g$  is the acceleration due to gravity. The plate’s circular sidewall is topped by a Plexiglas plate coated with a conducting film to prevent electrostatic build-up.

The granular medium is composed of two types of grain. The first layer above the oscillating plate is made up of dimeric ‘grains’, each consisting of two 3.2-mm steel spheres connected by a metal rod made from a cut bank chain (as used by banks to prevent patrons from removing their pens); each dimer has a mass ( $m_d$ ) of  $185 \pm 7$  mg. The second layer is composed of monomeric grains that are plastic (Delrin) spheres of diameter 3.175 mm and mass ( $m_m$ )  $22.9 \pm 0.2$  mg. The difference in mass between the grains of the first and second layers keeps the dimers and monomers separated, as the monomers cannot displace the heavier dimers.

The system is prepared by building a single layer of dimers on the plate (Fig. 1b) and placing a second layer of Delrin spheres on top (Fig. 1c). We define the coverage,  $c$ , of the layer to be the proportion of the maximum number of grains that could theoretically be packed into the area if wall effects are ignored. We collected data at  $c = 0.4, 0.6$  and

$0.8, f = 50, 70$  and  $90$  Hz, and at dimensionless accelerations ( $\Gamma$ ) of 1.75, 2.00 and 2.25.

Velocities in the first and second layers can be measured simultaneously by high-speed digital photography, although measurement by this method in the first layer becomes difficult at high coverage. The two layers show very different velocity distributions (Fig. 1d): in the first, the distribution tails are non-gaussian (typical deviation from gaussian flatness is about 1.5, which is consistent with earlier results<sup>5–12</sup>), whereas in the second layer the distribution is much more gaussian (typical deviation from gaussian flatness is less than 0.1).

The gaussian-velocity distributions in the second layer indicate that the grains are more ‘gas-like’ in their behaviour than has previously been found in vibrated granular media. This is probably due to the way in which energy is transferred to the second layer through collisions with the first layer throughout the shaking cycle. By contrast with earlier experiments, the gaussian velocity distribution of the second layer is observed at all coverages, frequencies and accelerations. Our system may therefore be ideally suited to developing a kinetic theory of granular gases.

G. W. Baxter\*, J. S. Olafsen†

\*School of Science, Penn State Erie, The Behrend College, Erie, Pennsylvania 16563, USA

†Department of Physics and Astronomy, University of Kansas, Lawrence, Kansas 66045, USA  
e-mail: jolafsen@ku.edu

- Haff, P. K. *J. Fluid Mech.* **134**, 401–430 (1983).
- Grossman, E. L., Zhou, T. & Ben-Naim, E. *Phys. Rev. E* **55**, 4200–4206 (1997).
- Huang, K. *Statistical Mechanics* (Addison-Wesley, Boston, 1963).
- Batchelor, G. K. (ed.) *An Introduction to Fluid Dynamics* (Cambridge Univ. Press, Cambridge, 1983).
- Olafsen, J. S. & Urbach, J. S. *Phys. Rev. Lett.* **81**, 4369–4372 (1998).
- Olafsen, J. S. & Urbach, J. S. *Phys. Rev. E* **60**, R2468–R2471 (1999).
- Losert, W., Cooper, D. G. W., Delour, J., Kudrolli, A. & Gollub, J. P. *Chaos* **9**, 682–690 (1999).
- Kudrolli, A., Wolpert, M. & Gollub, J. P. *Phys. Rev. Lett.* **78**, 1383–1386 (1997).
- Kudrolli, A. & Henry, I. *Phys. Rev. E* **62**, R1489–R1492 (2000).
- Blair, D. L. & Kudrolli, A. *Phys. Rev. E* **64**, R050301 1–4 (2001).
- Rouyer, F. & Menon, N. *Phys. Rev. Lett.* **85**, 3676–3679 (2000).
- Atwell, J. & Olafsen, J. S. *Phys. Rev. E* (submitted).

Competing financial interests: declared none.


corrigendum

A cloned horse born to its dam twin

Cesare Galli, Irina Lagutina, Gabriella Crotti, Silvia Colleoni, Paola Turini, Nunzia Ponderato, Roberto Duchi, Giovanna Lazzari  
*Nature* **424**, 635 (2003)

It has been drawn to our attention that a successful pregnancy in a goat carrying a genetically identical conceptus has previously been reported<sup>1</sup>; complete immune compatibility was demonstrated between the mother and her kid twin (both generated by splitting a single embryo). These findings also indicate that a maternal immune response is not necessary to support a healthy pregnancy.

- Oppenheim, S. M., Moyer, A. L., Bondurant, R. H., Rowe, J. D. & Anderson, G. B. *Theriogenology* **54**, 629–639 (2000).



**Figure 1** The two-layered granular system. **a**, Geometry of the experimental set-up; the accelerometer is shown at the bottom, on the right. **b**, Photograph of the dimeric layer: diameter of spheres, 3.2 mm; dimer mass, 185 mg. Note the extensive ordered region. **c**, Photograph of white Delrin balls (coverage ( $c$ ) = 0.4) at rest on top of the dimer layer (roughly one-sixth of the cell is shown). **d**, Velocity statistics for the first- and second-layer grains, for dimensionless acceleration  $\Gamma = 2.0$ . The distribution of the first-layer grains is non-gaussian. First layer: crosses,  $f = 70$  Hz,  $c = 0.40$ ; plus symbols,  $f = 50$  Hz,  $c = 0.60$ ; asterisks,  $f = 90$  Hz,  $c = 0.60$ . For the second-layer grains, the renormalized distribution is independent of system parameters within the range studied and is nearly gaussian (solid line) in shape. Second layer: diamonds,  $f = 70$  Hz,  $c = 0.40$ ; squares,  $f = 50$  Hz,  $c = 0.60$ ; circles,  $f = 90$  Hz,  $c = 0.60$ ; triangles,  $f = 70$  Hz,  $c = 0.80$ .