

BRIEF COMMUNICATIONS

Post-spawning egg care by a squid

Spying on a brooding deep-sea squid reveals that it cradles and aerates its eggs while they mature.

Gonatus onyx is one of the most abundant cephalopods in the Pacific and Atlantic Oceans¹ and is an important prey species for a variety of vertebrate predators^{2,3}, but a full understanding of its life history has been hampered because spawning occurs at great depths^{4,5}, where observation is difficult. Here we describe post-spawning egg care, or brooding, in this deep-sea squid. Our finding is unexpected because this behaviour differs from the reproductive habits of all other known squid species.

It was initially assumed that gonatids, like other squids, deposited their eggs on the sea floor and left them to develop on their own¹. A pelagic egg-brooding habit had been proposed for gonatid squids⁶, but could not be verified in the absence of direct observation. It had also been questioned because some aspects of the squids' biology seem to preclude brooding⁷ — principally, the degeneration of musculature following sexual maturation was presumed to limit locomotion and render squids unfit for egg protection^{5,7}.

However, we observed five squids, each holding an egg mass in its arms, at depths between 1,539 and 2,522 m in Monterey Canyon, off California, accessed with the ROV *Tiburon*⁸. Eggs or hatchlings and two adults were collected. The squids used hooks on their arms to hold the egg mass, which consists of two thin membranes in a continuous flat sheet, open at both the distal and proximal ends. The egg mass forms a hollow tube that extends from the mouth to well beyond the end of the arms and contains about 2,000–3,000 eggs (Fig. 1a, b).

Repeated extension of the arms (at intervals of about 30 to 40 s) flushed water through the egg mass: this behaviour probably served to aerate the eggs in the hypoxic midwaters found off California^{9,10}. Aggressive arm movements and escape swimming caused partial disintegration of the more mature egg masses and hatching of the released eggs (Fig. 1c–e). (For movies, see supplementary information.)

The escape responses seen here, considered with the apparent stage of egg development,




Figure 1 | Egg-brooding adult female *Gonatus onyx* (Cephalopoda: Teuthoidea), photographed *in situ*. Mantle length is about 145 mm. **a**, The squid in a horizontal resting position at 2,522 m depth. **b**, Squid holding a tubular egg mass, and **c**, hatchlings being released at 1,539 m depth. **d, e**, Hatched embryos (about 3 mm in length) at **d**, an intermediate stage of development, and **e**, an advanced stage of development. The *in situ* temperature (1.7–3.0 °C) and oxygen concentration (45–90 $\mu\text{mol l}^{-1}$) were measured using a SeaBird SBE-9 conductivity–temperature–depth unit with an oxygen sensor. Additional visual observations (see movie in supplementary information) were recorded with a high-resolution, three-chip video camera transmitting to the *RV Western Flyer* by fibre-optic cable. Assisted by vessel crew and ROV pilots.

indicate that the squid may have undergone a gradual degeneration of locomotory capacity. For example, a specimen bearing undeveloped eggs made a vigorous escape by using fin and mantle contractions, whereas those with advanced embryos (Fig. 1e) showed only respiratory mantle contractions and did not move away; another squid with eggs at an intermediate stage of development retained some locomotory ability (Fig. 1d) (for movies, see supplementary information). Activities of metabolic enzymes in locomotory muscles of spawned female *G. onyx* are lower in specimens that have more advanced embryos⁵.

Low temperature and large eggs will prolong development in *G. onyx*^{7,11}. The abundance of juveniles in near-surface waters peaks seasonally from April through to July¹, which may indicate a yearly cycle with an egg-development period of 6 to 9 months^{5,6}. This estimate is roughly consistent with the timing of our observations. Gonatid squids have sufficient lipid stores to fuel metabolism for a long brooding period, and the gradual decline

in mantle muscle and digestive-gland condition indicates a progressive use of energy stores over a long egg-brooding period⁶. However, two specimens captured within days of each other carried embryos at different stages, so development is apparently not well synchronized.

We have demonstrated active post-spawning egg care in *G. onyx* and this is, to our knowledge, the first case reported in squids. The value of this strategy in the deep sea is supported by strong convergence with distantly related taxa¹² and we expect it to be found in other squids. Despite retaining some capacity for swimming, the relatively immobile brooding squids are found within the usual diving range of whales and elephant seals and so may provide an easy target for such 'mesopelagic' mammals¹³. Gonatid squids and other ontogenetic migrators therefore represent a direct energetic, as well as trophic, link between deep and shallow biomes.

Brad A. Seibel*†, **Bruce H. Robison***, **Steven H. D. Haddock***

*Monterey Bay Aquarium Research Institute, Moss Landing, California 95039, USA
e-mail: seibel@uri.edu

†Present address: Biological Sciences Department, University of Rhode Island, Kingston, Rhode Island 02881, USA

- Okutani, T., Kubodera, T. & Jefferts, K. *Bull. Ocean. Res. Inst. Univ. Tokyo* **26**, 159–192 (1988).
- Clarke, M. R. *Phil. Trans. R. Soc. Lond. B* **351**, 979–983 (1996).
- Hooker, S. K., Whitehead, H. & Gowans, S. *Biol. Conserv.* **104**, 51–48 (2002).
- Arkhipkin, A. I. & Bjorke, H. *Polar Biol.* **22**, 347–365 (1999).
- Hunt, J. C. & Seibel, B. A. *Mar. Biol.* **136**, 543–552 (2000).
- Seibel, B. A., Hochberg, F. G. & Carlini, D. B. *Mar. Biol.* **137**, 519–526 (2000).
- Nesis, K. N. *Russ. J. Mar. Biol.* **25**, 499–506 (1999).
- Robison, B. H. *Mar. Technol. Soc. J.* **26**, 32–39 (1993).
- Strathmann, R. R. & Strathmann, M. F. *J. Mar. Biol. Assoc. UK* **75**, 413–428 (1995).
- Childress, J. J. & Seibel, B. A. *J. Exp. Biol.* **201**, 1223–1232 (1998).
- Boletzky, S. V. *Antarctic Sci.* **6**, 139–142 (1994).
- Childress, J. J. & Price, M. H. *Mar. Biol.* **76**, 165–177 (1983).
- Hochachka, P. W. *Experientia* **48**, 570–574 (1992).

Supplementary information accompanies this communication on Nature's website.

Competing financial interests: declared none.
doi:10.1038/438929a