



## Review

# The 'harmless' release of cytochrome *c*

O Von Ahsen<sup>1,2</sup>, NJ Waterhouse<sup>1,2</sup>, T Kuwana<sup>1,2</sup>,  
DD Newmeyer and DR Green<sup>\*1</sup>

<sup>1</sup> Division of Cellular Immunology, La Jolla Institute for Allergy and Immunology, 10355 Science Center Drive, San Diego, California, CA 92121, USA

<sup>2</sup> The first three authors contributed equally to this work

\* Corresponding author: Tel: 858-558-3500; Fax: 858-558-3526  
E-mail: DGreen5240@aol.com

Received 17.7.00; accepted 14.9.00

Edited by IG Kroemer

## Abstract

**Release of cytochrome *c* from the mitochondria plays an integral role in apoptosis; however, the mechanism by which cytochrome *c* is released remains one of the conundrums that has occupied the field. Recently, evidence has emerged that the commitment to death may be regulated downstream of cytochrome *c* release; therefore the mechanism of release must be subtle enough for the cell to recover from this event. In this review, we discuss the evidence that cytochrome *c* release is mediated by Bcl-2 family proteins in a process that involves only outer membrane permeability but leaves inner membrane energization, protein import function and the ultrastructure of mitochondria intact. *Cell Death and Differentiation* (2000) 7, 1192–1199.**

**Keywords:** cytochrome *c*; mitochondria; apoptosis; cell death; membrane permeability

**Abbreviations:** HSPs, heat shock proteins; DiOC6(3), 3,3'-dihexyloxycarbocyanine; zVAD-fmk, Z-val-ala-asp-fluoromethylketone; UV, ultraviolet light; TMRE, tetramethylrhodamine, ethyl ester; PT, permeability transition; ATR, atractyloside; tBHP, tert-butyl hydroperoxide; CCCP, carbonyl cyanide *m*-chlorophenylhydrazine

## Introduction

Ever since the early discovery that mitochondria enriched fractions are necessary for induction of apoptosis in cell free systems,<sup>1</sup> that Bcl-2 was localized mainly to the outer mitochondrial membrane,<sup>2,3</sup> and that mitochondria can lose membrane potential during apoptosis,<sup>4</sup> mitochondria have been the focus of apoptosis research. The observation that cytochrome *c* was involved in activation of caspases<sup>5,6</sup> and the demonstration that proapoptotic Bcl-2 family members (e.g. Bid, Bax, Bik and Bok) mediate the translocation of cytochrome *c* from mitochondria to cytosol,<sup>7–10</sup> an event that

is blocked by anti apoptotic Bcl-2 family members (e.g. Bcl-2 and Bcl-x<sub>L</sub>),<sup>11–13</sup> firmly established the mitochondrion as a central life and death regulator.

In the cytosol, holocytochrome *c* initiates the formation of a complex known as 'the apoptosome' by binding to an adaptor molecule Apaf-1. This triggers the oligomerization of Apaf-1 followed by recruitment and activation of procaspase-9.<sup>14,15</sup> Activated caspase-9 then cleaves and activates the downstream effectors of apoptotic cell death, caspases 3,6 and 7.<sup>16</sup> Recently, evidence has emerged that the key components of this pathway have been widely conserved during evolution.<sup>14,17–19</sup> Therefore, the release of cytochrome *c* from mitochondria to cytosol can now be regarded as a key regulatory event in apoptosis induced by a wide variety of stimuli. It appears however, that apoptosis induced by ligation of death receptors such as Fas may not require mitochondrial involvement in certain cell types.<sup>20</sup>

For a while, the release of cytochrome *c* was considered by many to be a point of no return and it did not matter if mitochondria were damaged in the process of cytochrome *c* release (since caspase activation would ensure fast and ordered destruction of the dying cell anyway). Recently IAP proteins have been shown to inhibit caspase activation downstream of cytochrome *c* release,<sup>21,22</sup> HSPs have been shown to modulate caspase activation<sup>23–25</sup> and some cells can recover after cytochrome *c* release.<sup>26–28</sup> This hints that cytochrome *c* release may not always commit a cell to die.

This appears surprising at first glance since cytochrome *c* and the integrity of the mitochondrial intermembrane space would appear to be essential for respiration and cell survival. Some cells do survive after cytochrome *c* release and therefore may be without respiration (cell survival in the absence of respiration has been observed in rho<sup>o</sup> cells<sup>29,30</sup>), however in order for mitochondria to exert their metabolic functions such as amino acid, heme and steroid metabolism, the membrane potential, protein import function and integrity of the inner membrane have to be maintained after cytochrome *c* release. It is therefore obvious that for cells to survive the release of cytochrome *c*, the mechanism of this release must be delicate enough to allow for the recovery of mitochondria provided that downstream events such as caspase activation are blocked.

In this review we will summarize the accumulating evidence that mitochondrial structure and function are well preserved after cytochrome *c* release in many experimental systems and discuss the mechanistic implications of this finding. We will also discuss the kinetics of cytochrome *c* release, the possible order of events in apoptotic cells and possible reasons for the loss of mitochondrial function being associated with cytochrome *c* release in some experimental settings.

## Cytochrome *c* release from isolated mitochondria

Isolated mitochondria provide a model system that has the natural protein and lipid composition of the outer membrane, and therefore reflect the target of cytochrome *c* releasing factors in the cell. Thus using isolated mitochondria and recombinant proapoptotic Bcl-2 proteins it should be possible to describe events that are associated with cytochrome *c* release that is mediated by proapoptotic Bcl-2 family proteins, and to distinguish between events that are sufficient and/or necessary for the release. An advantage of isolated mitochondria is the elimination of secondary events caused by cytosolic factors such as activated caspases.

Mitochondrial membrane potential ( $\Delta\psi_m$ ) is produced by the respiratory chain that uses the free energy of substrate oxidation to pump protons out of the matrix. This electrochemical potential across the inner membrane is used by ATP-synthase to produce ATP from ADP and  $P_i$ . Since mitochondria form the main ATP producing system of the cell,  $\Delta\psi_m$  is essential for efficient ATP production under aerobic conditions.  $\Delta\psi_m$  is also required for mitochondrial protein import and regulated metabolite transport. Because mitochondria are involved in pathways metabolising steroids, heme, pyrimidines and amino acids,  $\Delta\psi_m$  is often used as an indication of cellular viability.

In one of the first papers to describe cytochrome *c* release, Newmeyer and colleagues<sup>11</sup> used a *Xenopus* cell free system to show that  $\Delta\psi_m$  was not lost when cytochrome *c* was released, indicating that mitochondria remain relatively intact. Further work showed that cytochrome *c* was released from *Xenopus* mitochondria in a process involving only a limited permeabilization of the outer membrane.<sup>31</sup> The limited degree of outer membrane permeabilization in this system was further demonstrated by the observation that a cytosolic activity (named PEF for permeability enhancing factor) can enhance the limited permeability of the outer membrane caused by recombinant Bid or Bax.<sup>31</sup> The outer membranes of Bid or Bax treated mitochondria are permeable to cytochrome *c*, but the rate of cytochrome *c* traffic across the outer membrane (measured by oxidation of added cytochrome *c* catalyzed by cytochrome *c* oxidase) can be further enhanced by PEF. However, even the PEF-enhanced permeabilization of the outer membrane does not lead to damage of the inner membrane.<sup>31</sup> Recently von Ahsen *et al.* demonstrated that recombinant Bid and Bax can release cytochrome *c* from mitochondria without causing mitochondrial depolarization or inducing changes in mitochondrial ultrastructure.<sup>32</sup> Mitochondria that have lost their cytochrome *c* were shown to be in the same condensed conformation as control mitochondria. Electron microscopy as well as tomography<sup>33</sup> could not reveal any significant change in the ultrastructure. No holes, tears or ruptures in the outer membrane could be detected. This suggests a very specific release mechanism possibly involving a very small pore, too small to be visible by electron microscopy. Using mitochondrial protein import as an assay for mitochondrial membrane potential they could also rule out artifacts from other assays such as those observed when some fluorescent dyes were used to measure membrane potential. One problem of the dye

uptake assay is the influence of the plasma membrane potential.<sup>34</sup> The uptake of positively charged lipophilic dyes like rhodamine-123 has also been questioned because artifacts may arise due to self-quenching.<sup>35</sup> Furthermore, rhodamine 123 inhibits complex V,<sup>36</sup> while Mito Tracker orange and DioC6(3) inhibit complex I of the respiratory chain.<sup>37,38</sup> The posttranslational import of nuclear-encoded proteins into mitochondria however is a vital, natural function of mitochondria. The work of von Ahsen *et al.*<sup>32</sup> demonstrated that protein import was preserved after complete release of cytochrome *c*. This could be shown for isolated mitochondria from *Xenopus* oocytes and human HL-60 cells. Mitochondrial membrane potential was also maintained in apoptotic HeLa cells that were induced to undergo apoptosis by treatment with UV or staurosporine, provided that caspase activation was prevented by zVAD-fmk. Furthermore, after permeabilization of the plasma membrane by digitonin, the protein import function of mitochondria was shown to be maintained in apoptotic cells after cytochrome *c* release.

Others have published data that confirm a subtle mechanism of cytochrome *c* release in other systems: For example Jurgensmeier *et al.*<sup>9</sup> showed that recombinant Bax could induce cytochrome *c* release from isolated rat liver mitochondria without swelling, suggesting that a mechanical disruption of the outer membrane is not the mechanism. Similarly, Finucane *et al.*<sup>10</sup> have shown that Bax can induce cytochrome *c* release from isolated mouse liver mitochondria without swelling. Furthermore it was shown that mitochondrial depolarization in apoptotic 293T cells after Bax expression was a secondary event due to caspase activation since it could be blocked by coexpression of XIAP or addition of zVAD-fmk. This result provides an explanation for mitochondrial depolarization often observed in apoptotic cells.

## Cellular studies confirm a non-disruptive mechanism of cytochrome *c* release

The dynamic interaction of mitochondria with cells has made it difficult to objectively compare cytochrome *c* release with other events that occur during apoptosis. This has been compounded by the nature of the techniques used to assess cytochrome *c* release in cells (cellular fractionation followed by Western blotting or immunocytochemistry). Cellular fractionation generally involves mechanical disruption of cells and often results in incomplete rupturing of a large proportion of cells or excessive disruption of many of the mitochondria. Subsequent analysis of the fractions by Western blotting can therefore only be regarded as qualitative. Immunocytochemistry is quantitative; however the fixing process used in this technique limits the mitochondrial parameters other than cytochrome *c* release that can be assessed. For instance using immunocytochemistry, it is not possible to determine whether mitochondria that have released cytochrome *c* in cells can maintain protein import and many of the dyes used to measure  $\Delta\psi_m$  are released from the mitochondria upon fixation. In addition, dyes that measure  $\Delta\psi_m$  and can withstand fixation have been shown to induce mitochondrial permeability transition<sup>37</sup> and

many of these dyes inhibit various aspects of mitochondria metabolism (e.g. Rhodamine 123 inhibits ATP synthase,<sup>36</sup> and MitoTracker orange and DioC6(3) inhibit complex I,<sup>37,38</sup> as discussed).

More recently, green fluorescent protein (GFP)-tagged cytochrome *c* expressed in HeLa cells was shown to localize to the mitochondria and to be released concomitantly with endogenous cytochrome *c* during apoptosis.<sup>39</sup> The release of cytochrome *c*-GFP was complete in that it was evenly distributed throughout the cell. This excludes the involvement of an active export mechanism since cytochrome *c*-GFP appeared at the same concentration both inside and outside the mitochondria. Further, the addition of the caspase inhibitor zVAD-fmk did not prevent the release of cytochrome *c*-GFP during cytotoxic drug induced apoptosis, confirming previous data showing that cytochrome *c* release is caspase independent. These studies clearly demonstrate that mitochondria do not maintain a large proportion of their cytochrome *c* during apoptosis and exclude the possibility that some mitochondria release cytochrome *c* via a destructive mechanism (such as membrane rupture) while a subpopulation of mitochondria evade the cytochrome *c* releasing signal to aid in cell recovery (if required) after cytochrome *c* release. At least in these cells, all of the mitochondria release most of their cytochrome *c*.

Using these cells treated with a variety of apoptosis-inducing stimuli, it was shown that mitochondria which have released cytochrome *c*-GFP do not take up TMRE (a fluorescent dye used to measure  $\Delta\psi_m$ ). However, there were no circumstances where cells dying by apoptosis had a low membrane potential prior to the release of cytochrome *c*-GFP in the mitochondria. Further, in the presence of the caspase inhibitor zVAD-fmk, mitochondria of cells treated with a variety of stimuli maintained  $\Delta\psi_m$  (took up TMRE) for an extended period of time after they had released cytochrome *c*-GFP into the cytoplasm. In the cells that had released cytochrome *c* in the presence of zVAD-fmk, the  $\Delta\psi_m$  was maintained in the presence of oligomycin (an inhibitor of ATP synthase) but was not maintained in the presence of sodium azide (an inhibitor of complex IV of the electron transport chain), suggesting that even after cytochrome *c* release  $\Delta\psi_m$  is maintained by the electron transport chain. Since the contribution of complex IV to the  $\Delta\psi_m$  is dependent on acceptance of electrons from cytochrome *c*, this suggests that even when cytochrome *c* is diffuse through the cell, the concentration is sufficient to maintain respiration. These data demonstrate that the loss of  $\Delta\psi_m$  is not the cause of cytochrome *c* release as is predicted in some hypotheses, but is rather a consequence of cytochrome *c* release that is probably mediated by caspase-proteases. These data also showed clearly that mitochondria can survive the release of cytochrome *c*-GFP without compromising the integrity of the inner mitochondrial membrane. In support of this, cytochrome *c* release has been reported in HL-60 cells and neuronal cells treated with various death-inducing stimuli without a detectable loss of  $\Delta\psi_m$ <sup>40,41</sup> and addition of the caspase inhibitor zVAD-fmk prevented  $\Delta\psi_m$  loss in HeLa cells treated with UV but did not inhibit cytochrome *c*

release.<sup>42</sup> These studies in cells also confirmed the observations made using isolated mitochondria treated with pro-apoptotic Bcl-2 family members (as discussed previously).

Time lapse confocal microscopy using HeLa cells expressing cytochrome *c*-GFP has greatly extended our current knowledge of the mechanism of cytochrome *c* release in cells. These studies showed irrefutably that cytochrome *c* release is an early event during apoptosis occurring hours before phosphatidylserine exposure and loss of plasma membrane integrity. These studies also showed that although cytochrome *c*-GFP release is an early event, mitochondria retain cytochrome *c*-GFP for many hours after the initial apoptosis-inducing stimulus, however once initiated all mitochondria release cytochrome *c*-GFP in approximately 5 min regardless of the nature of the initial stimulus. The relatively invariable duration of cytochrome *c*-GFP release (~5 min) indicates that a universal pathway of cytochrome *c* release may be used during cytotoxic drug-induced and TNF-induced apoptosis or that the mechanisms of release are remarkably similar.

During TNF induced apoptosis, caspase-8 is activated early and mediates the release of cytochrome *c* via the activation of the pro-apoptotic Bcl-2 family member Bid.<sup>7,8</sup> This has led to the speculation that during apoptosis induced by cytotoxic drugs (that rely on the release of cytochrome *c* for caspase activation), caspases may signal for increased cytochrome *c*-release. The addition of zVAD-fmk inhibited the release of cytochrome *c*-GFP during TNF-induced apoptosis but did not extend the duration of cytochrome *c* release during apoptosis induced by other stimuli, thus confirming the participation of caspases in the release of cytochrome *c* during TNF induced apoptosis.<sup>39</sup> However the involvement of caspases in an amplification loop during apoptosis induced by other stimuli is unlikely in these cells.

Goldstein *et al.*<sup>39</sup> also showed that duration of cytochrome *c*-GFP release from mitochondria was similar over a temperature range of 13°C. Since the activity of most enzymes changes dramatically over 10°C, if enzymes were rate limiting during cytochrome *c* release, lower temperatures would be expected to slow the period over which cytochrome *c* is released. Individual mitochondria or groups of mitochondria were shown to release cytochrome *c* within 2 min and therefore the duration of cytochrome *c* release in cells was reported to reflect the time between cytochrome *c* release from the first and last mitochondria in the cell. At lower temperatures a slowed enzymatic process would result in cytochrome *c* release from the last mitochondria with a delayed kinetics after the first (if enzyme kinetics is slowed and plays a critical role in the kinetics), resulting in an apparent extension of the duration of cytochrome *c* release. Since an increase in duration was not observed, this reasoning suggested that the direct mechanism of cytochrome *c* release was temperature independent and did not rely on enzymes.

From the experiments performed on isolated mitochondria as well as in living cells, we conclude that proapoptotic Bcl-2 family members are sufficient to release cytochrome *c* and other proteins from the intermembrane space without

causing severe damage seen as depolarization and swelling of mitochondria. This is clearly inconsistent with all models based on matrix swelling and outer membrane rupture (caused by either permeability transition or hyperpolarization) as the primary mechanism of cytochrome *c* release in apoptotic cells.

The question arises of why mitochondria can keep their membrane potential high after cytochrome *c* release. If cytochrome *c* is not actively transported, it will only equilibrate within the total cell volume. Under conditions that require no or only low ATP synthase activity, the residual cytochrome *c* concentration might be sufficient to sustain respiration. So using isolated mitochondria in buffer, the dilution may be limiting for respiration. In cells, this dilution would be determined by mitochondrial content and cytoplasmic volume of the cell. Another factor may be the rate of ATP consumption in the cell. If ATP is consumed quickly, the remaining membrane potential may be used up, since the respiratory chain will not work efficiently under conditions of cytochrome *c* depletion. Some cells however may even compensate for low or absent respiration by enhanced glycolysis and reverse action of complex V to


maintain mitochondrial  $\Delta\psi_m$ . Thus if glucose is sufficient, cells can maintain  $\Delta\psi_m$  and thus may be able to survive cytochrome *c* release.

### Possible mechanisms of cytochrome *c* release

Data from our lab and from others clearly show that cytochrome *c* release can happen in a very subtle way leaving much of mitochondrial structure and function intact. What kind of a mechanism may account for such a gentle process? Some mechanisms for cytochrome *c* release have been proposed, among them very disruptive ways but also some ways of limited outer membrane permeabilization (Figure 1)

A delicate mechanism of cytochrome *c* release involving only the permeabilization of the outer membrane is suggested by findings that Bcl-2 family proteins can form pores in liposomes or artificial membranes. Structural studies of Bcl-x<sub>L</sub><sup>43</sup> revealed that the protein had a similar arrangement to the translocation domain of diphtheria toxin and of the colicins. Upon exposure to acid pH, the transmembrane domain of diphtheria toxin is inserted into

## Possible escape routes for Cytochrome *c*



**Figure 1** Models of cytochrome *c* release. Two of these models rely on matrix swelling causing mechanical disruption of the outer mitochondrial membrane. One model suggests that the opening of the permeability transition pore (PTP) is the initial change that leads to cytochrome *c* release. PTP is thought to be a complex composed of many proteins including hexokinase, voltage dependent anion channel (VDAC), adenine nucleotide translocator (ANT) and cyclophilin D (CypD). Opening of this pore leads to equilibration of solutes up to 1.5 kDa across mitochondrial membranes. This results in depolarization and subsequent matrix swelling due to colloid osmotic pressure. The second model suggests an initial hyperpolarization due to inhibition of ATP/ADP exchange via VDAC as reason for matrix swelling. Other models propose formation of pores for cytochrome *c* release. Bax (and maybe also other proapoptotic members of the bcl-2 family) could form channels in the outer membrane or alternatively cause membrane instability that leads to cytochrome *c* release. It has also been proposed that Bax regulates the pore size of VDAC to allow cytochrome *c* translocation

the lipid bilayer and forms an ion channel, followed by translocation of the catalytic domain through the lipid bilayer.<sup>44,45</sup> Similarly, recombinant Bcl-x<sub>L</sub> lacking the C-terminal hydrophobic region (Bcl-x<sub>L</sub>ΔC) was shown to form an ion channel in lipid bilayers at pH 4–5.<sup>46</sup> Subsequently it was shown that Bcl-2ΔC also exhibited an ion channel activity and released Cl<sup>-</sup> from liposomes at pH below 5.5.<sup>47</sup> Whether ion channel activity could account for the physiological role of Bcl-2 proteins remains an open question as (a) it is unlikely that mitochondria would be exposed to such extreme acidic pH and (b) Bcl-2 proteins regulate cytochrome *c* release from isolated mitochondria at neutral pH (and presumably in cells).

Subsequently the structure of Bid, a pro-apoptotic member of the Bcl-2 family, was determined. Its striking similarity to that of Bcl-x<sub>L</sub>ΔC suggested that Bcl-2 family proteins all interact with lipid bilayers in a similar manner.<sup>48,49</sup> As expected, Bid induced Cl<sup>-</sup> efflux from liposomes and exhibited ion channel activity in planar bilayers, with both activities facilitated by acidic pH.<sup>50</sup> More recently, there have been reports showing that recombinant Bid releases encapsulated proteins (cytochrome *c* and trypsin)<sup>51</sup> as well as carboxyfluorescein (CF)<sup>52</sup> from liposomes at neutral pH, without involving other proteins. In addition, the pro-apoptotic Bcl-2 protein, Bax, released CF from liposomes at neutral pH although this activity was increased at lower pH.<sup>53</sup> Significantly, the release of CF by BaxΔC was inhibited by addition of Bcl-2ΔC at neutral pH,<sup>53</sup> raising the possibility that Bcl-2ΔC exerts its effect without forming an ion channel.

Kudla *et al.*<sup>52</sup> found that Bid did not form ion channels, but ruptured the planar bilayer. In addition, although BaxΔC could form ion channels at low concentrations, at higher levels it also ruptured the lipid bilayer. *In vitro* translated full length Bax has also been found to rupture the membrane.<sup>54</sup> In summary, Bcl-2 family proteins can alter the permeability of lipid bilayers composed of various lipids, though negatively charged lipids were preferred,<sup>46,47,50</sup> suggesting that they can interact with the mitochondrial outer membrane in the same manner. Future studies utilizing the lipid composition reflecting that of the outer membrane of mitochondria at physiological pH will enhance our understanding of this process.

Although evidence has been published that Bax and Bid can permeabilize membranes on their own, Tsujimoto and colleagues<sup>55</sup> found that VDAC (voltage dependent anion channel), the most abundant protein located in the outer membrane of mitochondria was necessary for Bax to exert its effect. Bax also co-immunoprecipitated with VDAC from isolated mitochondria and in cells undergoing apoptosis.<sup>55</sup> Liposomes became permeable to sucrose upon incorporation of VDAC and this permeability was enhanced by recombinant full length Bax. The Bax effect was inhibitable by Bcl-x<sub>L</sub>.<sup>55</sup> Further, electrophysiological studies showed that Bax protein, together with VDAC in the planar bilayer, formed a novel ion channel larger than VDAC or Bax alone.<sup>56</sup> Permeability to cytochrome *c* was also demonstrated in the VDAC-loaded liposomes and in the planar bilayer systems in response to recombinant Bak or Bax.<sup>55,56</sup>

Thus, Bcl-2 family proteins, as their structures suggest, can interact with lipid bilayers, making them permeable to ions and small molecules and possibly to proteins as well. Alternatively, Bax, Bak and Bcl-x<sub>L</sub> may require proteins such as VDAC to release cytochrome *c* from mitochondria. Future studies including more comprehensive reconstitution will reveal whether Bcl-2 proteins regulate mitochondrial outer membrane proteins during apoptosis or act via direct pore formation.

Another suggested mechanism of apoptotic cytochrome *c* release is the mitochondrial permeability transition followed by large amplitude swelling that ultimately results in mechanical disruption of the outer membrane. Since a loss of mitochondrial membrane potential can be detected in apoptotic cells,<sup>57,58</sup> permeability transition (PT) was proposed as a possible mechanism for cytochrome *c* release. Permeability transition is an event described in the early days of mitochondrial research since some time after isolation, the mitochondria lose their ability for oxidative phosphorylation and eventually swell. It was then discovered that to promote mitochondrial survival, calcium must be excluded from the buffers. Now we know that the permeability transition can be induced by calcium, phosphate, low membrane potential or atractyloside (ATR) and inhibited by high membrane potential, adenine nucleotides, magnesium, bongkrekic acid and cyclosporin A. Until recently no physiological consequence was attributed to this event in cells, although regulation of intracellular calcium concentration, solute exchange between mitochondria and cytosol or even a role in heat generation, by analogy to brown adipose tissue were proposed.<sup>59</sup>

To investigate whether permeability transition could lead to release of cytochrome *c* and so be involved in apoptosis regulation, Kroemer and colleagues<sup>60,61</sup> treated isolated mitochondria with PT inducers and have shown that PT inducers (e.g. atractyloside, tBHP, CCCP) were sufficient to trigger apoptotic changes in added nuclei and that this was inhibitable by Bcl-2. Later however Marzo and colleagues<sup>62,63</sup> showed that active caspases can induce PT pore opening and release of proteins from the mitochondrial intermembrane space suggesting the possibility that PT is a secondary event caused by active caspases that might work as an amplification loop. Very recently, a mass spectroscopic approach to identify proteins released from mitochondria upon ATR induced PT revealed that this treatment led to a general disruption of membranes and to the release of mitochondrial matrix proteins and even proteins from contaminating peroxisomes and lysosomes.<sup>64</sup> These studies on artificial induction of PT proved that PT is sufficient to induce apoptosis, however since any mechanism of cytochrome *c* release will initiate an apoptotic response in cells, it has to be determined which events are necessary under physiological conditions to release cytochrome *c*. Since cytochrome *c* release in apoptotic cells is mediated by proapoptotic Bcl-2 proteins, the mechanism of cytochrome *c* release should be investigated using the effector proteins that mediate this event *in vivo*.

Taking this approach, Marzo *et al.*<sup>65</sup> showed release of cytochrome *c* and loss of  $\Delta\psi_m$  in isolated mitochondria treated with Bax and inhibition of Bax-dependent events by CsA and bongkrekic acid. However, it was not investigated whether  $\Delta\psi_m$  loss preceded cytochrome *c* release or was a late consequence of this event. Bax and the adenine nucleotide transporter (ANT) were proposed to physically interact, and a yeast strain lacking all three ANT genes did not die from Bax transfection, so ANT was proposed to be necessary for Bax to kill yeast. It was not shown however, whether Bax could mediate cytochrome *c* release from yeast mitochondria independently of ANT, and we don't know yet how Bax mediated death of yeast relates to cytochrome *c* release and apoptosis in multicellular organisms.

Most results obtained using yeast as a model system seem to show that an intact mitochondrial energy metabolism is necessary for Bax toxicity. Inner membrane proteins like ANT, respiratory chain components and a working ATP synthase were reported to be needed<sup>65–68</sup> whereas the involvement of VDAC and even the occurrence of cytochrome *c* release gave controversial results.<sup>55,67,69,70</sup> Since cytochrome *c* release is not convincingly associated with Bax toxicity in yeast this organism might not be a suitable model for apoptotic cytochrome *c* release.

Pastorino *et al.*<sup>71</sup> showed that moderate concentrations of Bax can release intermembrane space proteins without loss of membrane potential whereas high Bax concentrations induced a permeability transition as indicated by loss of membrane potential and large amplitude swelling.<sup>71</sup> They proposed that transient permeability transition pore openings may lead to release of proteins from the mitochondrial intermembrane space but that a permanent opening is needed for a measurable depolarization and large amplitude swelling.<sup>71</sup> However the authors failed to explain or even discuss how intermembrane space proteins could be released without large amplitude swelling, an event that seems to be required for the rupture of the outer membrane due to PT. A possible explanation may be that low concentrations of Bax are already sufficient for cytochrome *c* release to occur via a PT independent mechanism but that high Bax may lead to inner membrane damage by its pore forming or membrane destabilizing capabilities (see above).

Another hypothesis for cytochrome *c* release involves an initial hyperpolarization of mitochondria. Altered mitochondrial metabolism is proposed to result in an increased negative charge across the mitochondrial inner membrane. This increased stress would then lead to an influx of water to the matrix followed by rupturing of the outer membrane and subsequent cytochrome *c* release. However it is not clear whether this would result in an immediate loss of  $\Delta\psi_m$  in cells, since mitoplasts can maintain membrane potential even though the outer membrane is fragmented.

Using the fluorescent dye rhodamine 123, an increase in  $\Delta\psi_m$  (hyperpolarization) followed by a dramatic loss of  $\Delta\psi_m$  (depolarization) was detected early during apoptosis of FL5.12 cells induced to die by growth factor withdrawal.<sup>13</sup> The re-addition of growth factor rescued cells with hyperpolarized mitochondria but did not rescue cells that

had lost  $\Delta\psi_m$ . These results suggested that hyperpolarization occurred before commitment to apoptosis, and therefore probably before cytochrome *c* release. Both hyperpolarization and apoptosis were inhibited by Bcl-x<sub>L</sub>.<sup>13</sup> In recent reports, Vander Heiden *et al.*<sup>72,73</sup> demonstrated that during growth factor withdrawal induced apoptosis the outer membrane became impermeable to complex anions preventing the transport of ATP/ADP while the transfer of ATP/ADP across the inner membrane via the ANT was unaffected. This change in permeability was similar to that observed in planar lipids containing VDAC and indicated that VDAC may take on a closed state during apoptosis. After the ADP stores in the matrix have been used up, it is possible that continuing action of the electron transport chain may lead to an increase in mitochondrial  $\Delta\psi_m$ . The data presented also supported an anti-apoptotic role for Bcl-2 and Bcl-x<sub>L</sub> by maintaining ATP/ADP exchange across the outer membrane thereby preventing the increase in  $\Delta\psi_m$ .

Matsuyama *et al.*<sup>68</sup> recently demonstrated that oligomycin, an inhibitor of ATP synthase could inhibit apoptosis. ATP synthase utilizes the mitochondrial  $\Delta\psi_m$  to form ATP from ADP and Phosphate. Although ATP synthase can operate in reverse, under normal physiological conditions oligomycin would be expected to reduce the demand on the mitochondrial  $\Delta\psi_m$  resulting in hyperpolarization.<sup>68</sup> If hyperpolarization were the mechanism for cytochrome *c* release oligomycin would be thought to facilitate the release of cytochrome *c* and induce apoptosis. Indeed others have shown that oligomycin can induce apoptosis, however this appears to take some time, rather than by direct hyperpolarization of the mitochondria.<sup>74</sup> More recently, Matsuyama *et al.*<sup>75</sup> demonstrated that although it induced a mild increase in  $\Delta\psi_m$ , oligomycin appears to prevent the extent of hyperpolarization observed during staurosporine induced death.<sup>75</sup> This study also reported that an increase in matrix pH also preceded cytochrome *c* release. It is therefore possible that hyperpolarization is a consequence of altered metabolic events and that these events rather than hyperpolarization *per se* are responsible for cytochrome *c* release. Neither hyperpolarization nor an increase in pH of the mitochondrial matrix have been reported in isolated mitochondria treated with pro-apoptotic Bcl-2 family members and there is no hypothesis as to why an increased pH in the mitochondria would lead to cytochrome *c* release. Therefore we conclude that these changes may reflect metabolic alterations in dying cells but are not strictly required for cytochrome *c* release.

## Conclusions

Since considerable evidence shows that mitochondria can release cytochrome *c* without severe damage of structure and function, we conclude that cytochrome *c* release can be mediated by a 'delicate' mechanism. Mitochondrial structure and membrane potential dependent functions like protein import are retained after cytochrome *c* release.

Disruption of mitochondrial structure and function that are sometimes proposed to be linked with cytochrome *c* release were shown to be secondary in time and inhibitable

with caspase inhibitors when studied in detail. Thus the physiological release of cytochrome *c* in apoptosis appears to be mediated in a nondestructive way and mitochondrial damage to be caused later in time by activated caspases. The delicate pathway of cytochrome *c* release could allow cell recovery provided that other signals interfere with the downstream signaling pathway leading to caspase activation. From the data discussed in this review, we also conclude that signalling pathways converge at the mitochondria to release cytochrome *c* and the released cytochrome *c* leads to caspase activation in the cytoplasm if no downstream regulations prevent this. Once the caspase cascade is initiated by Apaf-1, all cellular caspases quickly become active and mediate the apoptotic degradation of cellular structures including mitochondria and the nucleus.

## References

1. Newmeyer DD, Farschon DM and Reed JC (1994) Cell-free apoptosis in *Xenopus* egg extracts: inhibition by Bcl-2 and requirement for an organelle fraction enriched in mitochondria. *Cell*. 79: 353–364
2. Nakai M, Takeda A, Cleary ML and Endo T (1993) The bcl-2 protein is inserted into the outer membrane but not into the inner membrane of rat liver mitochondria in vitro. *Biochem. Biophys. Res. Commun.* 196: 233–239
3. Nguyen M, Millar DG, Yong VW, Korsmeyer SJ and Shore GC (1993) Targeting of Bcl-2 to the mitochondrial outer membrane by a COOH-terminal signal anchor sequence. *J. Biol. Chem.* 268: 25265–25268
4. Zamzami N, Marchetti P, Castedo M, Decaudin D, Macho A, Hirsch T, Susin SA, Petit PX, Mignotte B and Kroemer G (1995) Sequential reduction of mitochondrial transmembrane potential and generation of reactive oxygen species in early programmed cell death. *J. Exp. Med.* 182: 367–377
5. Liu X, Kim CN, Yang J, Jemmerson R and Wang X (1996) Induction of apoptotic program in cell-free extracts: requirement for dATP and cytochrome *c*. *Cell*. 86: 147–157
6. Zhivotovskiy B, Orrenius S, Brustugun OT and Døskeland SO (1998) Injected cytochrome *c* induces apoptosis [letter] [see comments]. *Nature*. 391: 449–450
7. Luo X, Budihardjo I, Zou H, Slaughter C and Wang X (1998) Bid, a Bcl2 interacting protein, mediates cytochrome *c* release from mitochondria in response to activation of cell surface death receptors. *Cell*. 94: 481–490
8. Li H, Zhu H, Xu CJ and Yuan J (1998) Cleavage of BID by caspase 8 mediates the mitochondrial damage in the Fas pathway of apoptosis. *Cell*. 94: 491–501
9. Jurgensmeier JM, Xie Z, Deveraux Q, Ellerby L, Bredesen D and Reed JC (1998) Bax directly induces release of cytochrome *c* from isolated mitochondria. *Proc. Natl. Acad. Sci. USA* 95: 4997–5002
10. Finucane DM, Bossy-Wetzler E, Waterhouse NJ, Cotter TG and Green DR (1999) Bax-induced caspase activation and apoptosis via cytochrome *c* release from mitochondria is inhibitable by Bcl-xL. *J. Biol. Chem.* 274: 2225–2233
11. Kluck RM, Bossy-Wetzler E, Green DR and Newmeyer DD (1997) The release of cytochrome *c* from mitochondria: a primary site for Bcl-2 regulation of apoptosis. *Science*. 275: 1132–1136
12. Yang J, Liu X, Bhalla K, Kim CN, Ibrado AM, Cai J, Peng T-I, Jones DP and Wang X (1997) Prevention of apoptosis by Bcl-2: release of cytochrome *c* from mitochondria blocked. *Science*. 275: 1129–1132
13. Vander Heiden MG, Chandel NS, Williamson EK, Schumacker PT and Thompson CB (1997) Bcl-xL regulates the membrane potential and volume homeostasis of mitochondria [see comments]. *Cell*. 91: 627–637
14. Zou H, Henzel WJ, Liu X, Lutschg A and Wang X (1997) Apaf-1, a human protein homologous to *C. elegans* CED-4, participates in cytochrome *c*-dependent activation of caspase-3 [see comments]. *Cell*. 90: 405–413
15. Zou H, Li Y, Liu X and Wang X (1999) An APAF-1/cytochrome *c* multimeric complex is a functional apoptosome that activates procaspase-9. *J. Biol. Chem.* 274: 11549–11556
16. Thornberry NA and Lazebnik Y (1998) Caspases: enemies within. *Science*. 281: 1312–1316
17. Rodriguez A, Oliver H, Zou H, Chen P, Wang X and Abrams JM (1999) Dark is a *Drosophila* homologue of Apaf-1/CED-4 and functions in an evolutionarily conserved death pathway [see comments]. *Nat. Cell. Biol.* 1: 272–279
18. Colussi PA, Quinn LM, Huang DC, Coombe M, Read SH, Richardson H and Kumar S (2000) Debcl, a proapoptotic Bcl-2 homologue, is a component of the *Drosophila melanogaster* cell death machinery [see comments]. *J. Cell. Biol.* 148: 703–714.
19. Kanuka H, Sawamoto K, Inohara N, Matsuno K, Okano H and Miura M (1999) Control of the cell death pathway by Dapaf-1, a *Drosophila* Apaf-1/CED-4-related caspase activator. *Mol. Cell*. 4: 757–769.
20. Scaffidi C, Fulda S, Srinivasan A, Friesen C, Li F, Tomaselli KJ, Debatin KM, Kramer PH and Peter ME (1998) Two CD95 (APO-1/Fas) signaling pathways. *EMBO J.* 17: 1675–1687.
21. Deveraux QL, Roy N, Stennicke HR, Van Arsdale T, Zhou Q, Srinivasula SM, Alnemri ES, Salvesen GS and Reed JC (1998) IAPs block apoptotic events induced by caspase-8 and cytochrome *c* by direct inhibition of distinct caspases. *EMBO J.* 17: 2215–2223.
22. Deveraux QL, Leo E, Stennicke HR, Welsh K, Salvesen GS and Reed JC (1999) Cleavage of human inhibitor of apoptosis protein XIAP results in fragments with distinct specificities for caspases. *EMBO J.* 18: 5242–5251.
23. Jaattela M, Wissing D, Kokholm K, Kallunki T and Egeblad M (1998) Hsp 70 exerts its anti-apoptotic function downstream of caspase-3-like proteases. *EMBO J.* 17: 6124–6134
24. Samali A and Orrenius S (1998) Heat shock proteins: regulators of stress response and apoptosis. *Cell. Stress Chaperones* 3: 228–236
25. Beere HM, Wolf BB, Cain K, Mosser DD, Mahboubi A, Kuwana T, Taylor P, Morimoto RI, Cohen GM and Green DR (2000) Heat-shock protein 70 inhibits apoptosis by preventing recruitment of procaspase-9 to the Apaf-1 apoptosome. *Nat. Cell. Biol.* 2: 469–475.
26. Yoshida H, Kong YY, Yoshida R, Elia AJ, Hakem A, Hakem R, Penninger JM and Mak TW (1998) Apaf1 is required for mitochondrial pathways of apoptosis and brain development. *Cell*. 94: 739–750.
27. Hakem R, Hakem A, Duncan GS, Henderson JT, Woo M, Soengas MS, Elia A, de la Pompa JL, Kagi D, Khoo W, Potter J, Yoshida R, Kaufman SA, Lowe SW, Penninger JM and Mak TW (1998) Differential requirement for caspase 9 in apoptotic pathways in vivo. *Cell*. 94: 339–352
28. Martinou I, Desagher S, Eskes R, Antonsson B, Andre E, Fakan S and Martinou JC (1999) The release of cytochrome *c* from mitochondria during apoptosis of NGF-deprived sympathetic neurons is a reversible event. *J. Cell. Biol.* 144: 883–889
29. Donnelly M and Scheffler IE (1976) Energy metabolism in respiration-deficient and wild type Chinese hamster fibroblasts in culture. *J. Cell. Physiol.* 89: 39–51
30. Martinou RD, Linnane AW and Nagley P (1993) Growth of rho 0 human Namalwa cells lacking oxidative phosphorylation can be sustained by redox compounds potassium ferricyanide or coenzyme Q10 putatively acting through the plasma membrane oxidase. *Biochem. Mol. Biol. Int.* 31: 997–1005
31. Kluck RM, Esposti MD, Perkins G, Renken C, Kuwana T, Bossy-Wetzler E, Goldberg M, Allen T, Barber MJ, Green DR and Newmeyer DD (1999) The proapoptotic proteins, Bid and Bax, cause a limited permeabilization of the mitochondrial outer membrane that is enhanced by cytosol. *J. Cell. Biol.* 147: 809–822
32. von Ahlsen O, Renken C, Perkins G, Kluck RM, Bossy-Wetzler E and Newmeyer DD. (2000) Preservation of mitochondrial structure and function after Bid- or Bax-mediated cytochrome *c* release. *J. Cell. Biol.* 150: 1027–1036
33. Perkins GA, Renken CW, Song JY, Frey TG, Young SJ, Lamont S, Martone ME, Lindsey S and Ellisman MH (1997) Electron tomography of large, multi-component biological structures. *J. Struct. Biol.* 120: 219–227
34. Salvioli S, Ardizzone A, Franceschi C and Cossarizza A (1997) JC-1, but not DiOC6(3) or rhodamine 123, is a reliable fluorescent probe to assess delta psi changes in intact cells: implications for studies on mitochondrial functionality during apoptosis. *FEBS Lett.* 411: 77–82
35. Emaus RK, Grunwald R and Lemasters JJ (1986) Rhodamine 123 as a probe of transmembrane potential in isolated rat liver mitochondria: spectral and metabolic properties. *Biochim. Biophys. Acta.* 850: 436–448
36. Modica-Napolitano JS, Weiss MJ, Chen LB and Aprille JR (1984) Rhodamine 123 inhibits bioenergetic function in isolated rat liver mitochondria. *Biochem. Biophys. Res. Commun.* 118: 717–723

37. Scorrano L, Petronilli V, Colonna R, Di Lisa F and Bernardi P (1999) Chloromethyltetramethylrosamine (Mitotracker Orange) induces the mitochondrial permeability transition and inhibits respiratory complex I. Implications for the mechanism of cytochrome c release. *J. Biol. Chem.* 274: 24657–24663
38. Anderson WM, Wood JM and Anderson AC (1993) Inhibition of mitochondrial and Paracoccus denitrificans NADH-ubiquinone reductase by oxocarboxyanine dyes. A structure-activity study. *Biochem. Pharmacol.* 45: 2115–2122.
39. Goldstein JC, Waterhouse NJ, Juin P, Evan GI and Green DR (2000) The coordinate release of cytochrome c during apoptosis is rapid, complete and kinetically invariant [see comments]. *Nat. Cell. Biol.* 2: 156–162.
40. Finucane DM, Waterhouse NJ, Amarante-Mendes GP, Cotter TG and Green DR (1999) Collapse of the inner mitochondrial transmembrane potential is not required for apoptosis of HL60 cells. *Exp. Cell. Res.* 251: 166–174
41. Krohn AJ, Wahlbrink T and Prehn JH (1999) Mitochondrial depolarization is not required for neuronal apoptosis. *J. Neurosci.* 19: 7394–7404
42. Bossy-Wetzell E, Newmeyer DD and Green DR (1998) Mitochondrial cytochrome c release in apoptosis occurs upstream of DEVD-specific caspase activation and independently of mitochondrial transmembrane depolarization. *EMBO. J.* 17: 37–49
43. Muchmore SW, Sattler M, Liang H, Meadows RP, Harlan JE, Yoon HS, Nettlesheim D, Chang BS, Thompson CB, Wong S-L, Ng S-C and Fesik SW (1996) X-ray and NMR structure of human Bcl-xL, an inhibitor of programmed cell death. *Nature* 381: 335–341
44. Moskaug JO, Stenmark H and Olsnes S (1991) Insertion of diphtheria toxin B-fragment into the plasma membrane at low pH. Characterization and topology of inserted regions. *J. Biol. Chem.* 266: 2652–2659
45. Choe S, Bennett MJ, Fujii G, Curmi PM, Kantardjiev KA, Collien RJ and Eisenberg D (1992) The crystal structure of diphtheria toxin. *Nature* 357: 216–222
46. Minn AJ, Velez P, Schendel SL, Liang H, Muchmore SW, Fesik SW, Fill M and Thompson CB (1997) Bcl-x(L) forms an ion channel in synthetic lipid membranes. *Nature* 385: 353–357
47. Schendel SL, Xie Z, Montal MO, Matsuyama S, Montal M and Reed JC (1997) Channel formation by antiapoptotic protein Bcl-2. *Proc. Natl. Acad. Sci. USA* 94: 5113–5118
48. Chou JJ, Li H, Salvesen GS, Yuan J and Wagner G (1999) Solution structure of BID, an intracellular amplifier of apoptotic signaling. *Cell* 96: 615–624
49. McDonnell JM, Fushman D, Milliman CL, Korsmeyer SJ and Cowburn D (1999) Solution structure of the proapoptotic molecule BID: a structural basis for apoptotic agonists and antagonists. *Cell* 96: 625–634
50. Schendel SL, Azimov R, Pawlowski K, Godzik A, Kagan BL and Reed JC (1999) Ion channel activity of the BH3 only Bcl-2 family member, BID. *J. Biol. Chem.* 274: 21932–21936
51. Zhai D, Huang X, Han X and Yang F (2000) Characterization of tBid-induced cytochrome c release from mitochondria and liposomes [In Process Citation]. *FEBS Lett.* 472: 293–296
52. Kudla G, Montessuit S, Eskes R, Berrier C, Martinou JC, Ghazi A and Antonsson B (2000) The destabilization of lipid membranes induced by the C-terminal fragment of Caspase 8-cleaved Bid is inhibited by the N-terminal fragment. *J. Biol. Chem.* 275: 22713–22718
53. Antonsson B, Conti F, Ciavatta A, Montessuit S, Lewis S, Martinou I, Bernasconi L, Bernard A, Mermod JJ, Mazzei G, Maundrell K, Gambale F, Sadoul R and Martinou JC (1997) Inhibition of Bax channel-forming activity by Bcl-2. *Science* 277: 370–372
54. Basanez G, Nechushtan A, Drozhinin O, Chanturiya A, Choe E, Tutt S, Wood KA, Hsu Y, Zimmerberg J and Youle RJ (1999) Bax, but not Bcl-xL, decreases the lifetime of planar phospholipid bilayer membranes at subnanomolar concentrations. *Proc. Natl. Acad. Sci. USA* 96: 5492–5497
55. Shimizu S, Narita M and Tsujimoto Y (1999) Bcl-2 family proteins regulate the release of apoptogenic cytochrome c by the mitochondrial channel VDAC [see comments]. *Nature* 399: 483–487
56. Shimizu S, Ide T, Yanagida T and Tsujimoto Y (2000) Electrophysiological study of a novel large pore formed by Bax and the voltage-dependent anion channel that is permeable to cytochrome c. *J. Biol. Chem.* 275: 12321–12325
57. Vayssiere JL, Petit PX, Rislis Y and Mignotte B (1994) Commitment to apoptosis is associated with changes in mitochondrial biogenesis and activity in cell lines conditionally immortalized with simian virus 40. *Proc. Natl. Acad. Sci. USA* 91: 11752–11756
58. Zamzami N, Marchetti P, Castedo M, Zanin C, Vayssiere JL, Petit PX and Kroemer G (1995) Reduction in mitochondrial potential constitutes an early irreversible step of programmed lymphocyte death in vivo. *J. Exp. Med.* 181: 1661–1672
59. Zoratti M and Szabo I (1995) The mitochondrial permeability transition. *Biochim. Biophys. Acta.* 1241: 139–176
60. Susin SA, Zamzami N, Castedo M, Hirsch T, Marchetti P, Macho A, Daugas E, Geuskens M and Kroemer G (1996) Bcl-2 inhibits the mitochondrial release of an apoptogenic protease. *J. Exp. Med.* 184: 1331–1341
61. Zamzami N, Susin SA, Marchetti P, Hirsch T, Gomez-Monterrey I, Castedo M and Kroemer G (1996) Mitochondrial control of nuclear apoptosis [see comments]. *J. Exp. Med.* 183: 1533–1544
62. Marzo I, Brenner C, Zamzami N, Susin SA, Beutner G, Brdiczka D, Remy R, Xie ZH, Reed JC and Kroemer G (1998) The permeability transition pore complex: a target for apoptosis regulation by caspases and bcl-2-related proteins. *J. Exp. Med.* 187: 1261–1271
63. Marzo I, Susin SA, Petit PX, Ravagnan L, Brenner C, Larochette N, Zamzami N and Kroemer G (1998) Caspases disrupt mitochondrial membrane barrier function. *FEBS Lett.* 427: 198–202
64. Patterson SD, Spahr CS, Daugas E, Susin SA, Irinopoulou T, Koehler C and Kroemer G (2000) Mass spectrometric identification of proteins released from mitochondria undergoing permeability transition. *Cell Death Differ.* 7: 137–144
65. Marzo I, Brenner C, Zamzami N, Jurgensmeier JM, Susin SA, Vieira HL, Prevost MC, Xie Z, Matsuyama S, Reed JC and Kroemer G (1998) Bax and adenine nucleotide translocator cooperate in the mitochondrial control of apoptosis. *Science* 281: 2027–2031
66. Harris MH, Vander Heiden MG, Kron SJ and Thompson CB (2000) Role of oxidative phosphorylation in Bax toxicity. *Mol. Cell. Biol.* 20: 3590–3596
67. Priault B, Chaudhuri B, Clow A, Camougrand N and Manon S (1999) Investigation of bax-induced release of cytochrome c from yeast mitochondria permeability of mitochondrial membranes, role of VDAC and ATP requirement. *Eur. J. Biochem.* 260: 684–691
68. Matsuyama S, Xu Q, Velours J and Reed JC (1998) The Mitochondrial F0F1-ATPase proton pump is required for function of the proapoptotic protein Bax in yeast and mammalian cells. *Mol. Cell.* 1: 327–336
69. Roucou X, Prescott M, Devenish RJ and Nagley P (2000) A cytochrome c-GFP fusion is not released from mitochondria into the cytoplasm upon expression of Bax in yeast cells. *FEBS Lett.* 471: 235–239
70. Gross A, Pilcher K, Blachly-Dyson E, Basso E, Jockel J, Bassik MC, Korsmeyer SJ and Forte M (2000) Biochemical and genetic analysis of the mitochondrial response of yeast to BAX and BCL-X(L). *Mol. Cell. Biol.* 20: 3125–3136
71. Pastorino JG, Tafani M, Rothman RJ, Marcinekivicute A, Hoek JB, Farben JL and Marcinekivicute A (1999) Functional consequences of the sustained or transient activation by Bax of the mitochondrial permeability transition pore [published erratum appears in *J. Biol. Chem.* 2000 Mar 17; 275(11): 8262]. *J. Biol. Chem.* 274: 31734–31739
72. Vander Heiden MG, Chandel NS, Schumacker PT and Thompson CB (1999) Bcl-xL prevents cell death following growth factor withdrawal by facilitating mitochondrial ATP/ADP exchange. *Mol. Cell.* 3: 159–167
73. Vander Heiden MG, Chandel NS, Li XX, Schumacker PT, Colombini M and Thompson CB (2000) Outer mitochondrial membrane permeability can regulate coupled respiration and cell survival. *Proc. Natl. Acad. Sci. USA* 97: 4666–4671
74. Wolvetang EJ, Johnson KL, Krauer K, Ralph SJ and Linnane AW (1994) Mitochondrial respiratory chain inhibitors induce apoptosis. *FEBS Lett.* 339: 40–44
75. Matsuyama S, Llopis J, Deveraux QL, Tsien RY and Reed JC (2000) Changes in intramitochondrial and cytosolic pH: early events that modulate caspase activation during apoptosis. *Nat. Cell. Biol.* 2: 318–325