

EGF activates an inducible survival response via the RAS-> Erk-1/2 pathway to counteract interferon- α -mediated apoptosis in epidermoid cancer cells

M Caraglia¹, P Tagliaferri², M Marra¹, G Giuberti¹, A Budillon³, E Di Gennaro³, S Pepe⁴, G Vitale⁴, S Improta¹, P Tassone², S Venuta², AR Bianco⁴ and A Abbruzzese^{*,1}

¹ Dipartimento di Biochimica e Biofisica, Seconda Università di Napoli, Via Costantinopoli, 16 80138, Naples, Italy

² Dipartimento di Medicina Sperimentale, Università 'Magna Grecia' di Catanzaro, Italy

³ Istituto Nazionale dei Tumori 'G Pascale', Naples, Italy

⁴ Dipartimento di Endocrinologia ed Oncologia Molecolare e Clinica, Università 'Federico II' di Napoli, Naples, Italy

* Corresponding author: A Abbruzzese, Dipartimento di Biochimica e Biofisica, Seconda Università di Napoli, Via Costantinopoli, 16 80138, Naples, Italy

Received 23.5.02; revised 24.7.02; accepted 28.8.02

Edited by G. Ciliberto

Abstract

The mechanisms of tumor cell resistance to interferon- α (IFN α) are at present mostly unsolved. We have previously demonstrated that IFN α induces apoptosis on epidermoid cancer cells and EGF antagonizes this effect. We have also found that IFN α -induced apoptosis depends upon activation of the NH₂-terminal Jun kinase-1 (Jnk-1) and p³⁸ mitogen-activated protein kinase, and that these effects are also antagonized by EGF. At the same time, IFN α increases the expression and function of the epidermal growth factor receptor (EGF-R). Here we report that the apoptosis induced by IFN α occurs together with activation of caspases 3, 6 and 8 and that EGF also antagonizes this effect. On the basis of these results, we have hypothesized that the increased EGF-R expression and function could represent an inducible survival response that might protect tumor cells from apoptosis caused by IFN α via extracellular signal regulated kinase 1 and 2 (Erk-1/2) cascades. We have found an increased activity of Ras and Raf-1 in IFN α -treated cells. Moreover, IFN α induces a 50% increase of the phosphorylated isoforms and enzymatic activity of Erk-1/2. We have also demonstrated that the inhibition of Ras activity induced by the transfection of the dominant negative Ras plasmid RASN17 and the inhibition of Mek-1 with PD098059 strongly potentiates the apoptosis induced by IFN α . Moreover, the selective inhibition of this pathway abrogates the counteracting effect of EGF on the IFN α -induced apoptosis. All these findings suggest that epidermoid tumor cells counteract the IFN α -induced apoptosis through a survival pathway that involves the hyperactivation of the EGF-dependent Ras->Erk signalling. The selective targeting of this pathway appears to be a promising approach in order to enhance the antitumor activity of IFN α .

Cell Death and Differentiation (2003) 10, 218–229. doi:10.1038/sj.cdd.4401131

Keywords: Epidermal growth factor receptor; Dominant negative Ras; Mek-1 inhibitor; Caspases; New anticancer strategies

Abbreviations: IFN α , Interferon- α ; JNK-, Jun kinase-1; EGF-R, Epidermal growth factor receptor; MAPK, Mitogen-activated protein kinase

Introduction

Interferon- α (IFN α) is a biological agent widely used in the therapy of several neoplasms such as myeloma, renal cell carcinoma, epidermoid cervical and head and neck tumors and melanoma.^{1–5} IFN α , the first cytokine to be produced by recombinant DNA technology, has emerged as an important regulator of cancer cell growth and differentiation, affecting cellular communication and signal transduction pathways, and is also produced during the immunological control of tumor growth.⁵ However, inconsistent data have been obtained regarding the clinical effectiveness of IFN α in the therapy of solid tumors. In fact, the benefit of IFN α treatment is limited to some neoplasms while others are completely or partially resistant. The mechanisms of tumor resistance to IFN α have been studied *in vitro*. The alteration of JAK-STAT components of the IFN α -induced signalling can be indeed a mechanism of resistance to IFN. In fact, it has been shown that melanoma cell lines refractory to the antiproliferative effects of IFNs are deficient in STATs and that the expression of STATs can be restored by *in vitro* gene therapy.⁶ More germane to clinical practice is the possibility that IFN treatment could be improved by the concomitant administration of agents known to enhance JAK-STAT responses; the use of retinoids in combination with IFN α in cancer therapy is a salient example.^{7,8}

The way in which tumor cell growth is suppressed by IFN α is not well known. A possibility is that cancer cells undergo apoptosis after exposure to the cytokine. In fact, it is reported that IFN α induces apoptosis in human squamous cancer,^{9,10} glioma¹¹ and virus-infected cells.¹² Therefore, it is likely that this cytokine acts, at least in part, through the triggering of programmed cell death.

On the other hand, EGF is able to protect eukaryotic cells from the onset of apoptosis.^{13–16} EGF acts through the binding to its specific receptor, EGF-R a transmembrane protein with a cytoplasmic tyrosine kinase domain.^{17,18} The phosphorylation of the intracytoplasmic tail allows the interaction of EGF-R with cytoplasmic factors that can induce Ras activation and the subsequent stimulation of the mitogen-activated protein kinase (MAPK) cascade composed of three

intracellular protein kinases (MKKK, MKK and MAPK), which are activated successively by phosphorylation events.^{17,18} Raf-1, stimulated after steric interaction with Ras, phosphorylates and activates an MKK whose main component is Mek-1. Mek-1 phosphorylates the MAPKs extracellular signal regulated kinase 1 and 2 (Erk-1/2) that translocate to the nucleus and phosphorylate gene transactivators, such as the serum response factor-1, that are involved in the regulation of cell proliferation.^{17,18} Moreover, it has been recently reported that Erks mediate a strong antiapoptotic effect.^{19,20} For instance, Erk plays an important role in protecting cardiac myocytes from apoptotic death following oxidative stress.²¹ Protection of PC12 cell death by N-acetylcysteine requires Erk activation²⁰ that is, moreover, involved in the phorbol myristate acetate-mediated inhibition of drug-induced apoptosis in tumor cells.²²

We have reported that IFN α increases the functional expression of the epidermal growth factor receptor (EGF-R) at the surface of human epidermoid carcinoma cells.^{23,24} Moreover, we have shown that IFN α induces apoptosis in these cells through the triggering of a stress response that leads to the activation of NH₂-terminal Jun kinase-1 (Jnk-1).¹⁰ EGF also antagonizes these effects. On the basis of these findings, we have hypothesized that increased EGF-R expression and function could be part of an inducible survival pathway, which is activated in the tumor cells by exposure to IFN α . In this paper, we have investigated the molecular basis of the counteracting effects of IFN α and EGF on the apoptosis of human epidermoid cancer cells. Specifically, we have evaluated the involvement of the caspase cascade and of EGF-dependent signalling in the apoptotic and antiapoptotic events recorded in our experimental model. We have then studied the effects of IFN α and EGF on the activity of Ras, Raf-1 and Erk-1/2. We have finally evaluated if the selective inhibition of Ras and Mek-1 could enhance apoptosis induced by IFN α in order to provide direct evidence on the antiapoptotic activity of this pathway in our experimental model and to identify novel molecular targets to enhance the antitumor effects of this cytokine.

Results

Effects of IFN α and EGF on the apoptosis of human epidermoid cancer cells

We have found that IFN α (48 h 1000 IU/ml) induced apoptotic DNA fragmentation in the KB cell line and that the exposure of IFN α -treated cells to 10 nM EGF for 12 h antagonized this effect (Figure 1a). Before performing apoptosis detection assays, both attached and suspended cells were collected. In these experimental conditions, 30% of cells exposed to IFN α for 48 h were apoptotic while only about 8% of cells treated for 48 h with IFN α and exposed for 12 h to 10 nM EGF underwent to apoptosis (Figure 1b). Interestingly, also the exposure to 10 nM EGF alone for 12 h is able to induce apoptosis in 15% of the cell population (Figure 1a and b). Such an effect is not surprising since apoptosis and cell proliferation are coupled processes, and EGF can induce apoptosis in squamous cancer cells. In order to identify the molecular effectors of the apoptosis induced by IFN α , we have evaluated the expression

of p53 and c-myc and the activity of caspases in KB cells. We have found that apoptosis induced by either IFN α or EGF was paralleled by a significant increase of p53 and c-myc expression in KB cells, as evaluated with Western blot assay. The antagonism of EGF on apoptosis occurred together with the restoration of p53 and c-myc expression to basal levels (Figure 1C). We have thereafter studied the role of caspases in the modulation of apoptosis by IFN α and EGF in our experimental model. KB cells were treated with IFN α for 12, 24, 48 and 72 h and then the activity of caspases was evaluated with a fluorimetric protease assay. Caspases 6, 8 and 9 showed an activation peak after 24 h exposure to IFN α (Figure 2a). The activation peak of caspase 3 was instead found after 48 h exposure to the cytokine. Moreover, the exposure of KB cells to 10 nM EGF for 12 h was always able to antagonize the caspase activation induced by IFN α both at 24 h and 48 h (Figure 2b-d). In fact, EGF completely abrogated caspases 6, 8 and 9 and caspase 3 activities. EGF alone also induced a slight increase of caspase stimulation, but the extent of activation was far lesser than that recorded in IFN α -treated cells. Therefore, in our experimental model the IFN α -induced apoptosis is correlated with the activation of the caspase cascade, and EGF was able to antagonize both apoptotic death and caspase activation. Moreover, the apoptosis recorded in KB cells exposed to EGF alone did not appear to be induced by a caspase pathway similar to that recorded in IFN α -treated cells.

Effects of EGF and IFN α on Ras activity

We have previously shown that IFN α induces growth inhibition paralleled by increased EGF-R expression and sensitivity to the growth-promoting effects of EGF in epidermoid cancer cells.^{23,24} Therefore, we have investigated if the activity of the Ras-dependent MAPK pathway was increased in IFN α -treated cells. This and all the following experiments were performed without serum starvation in order to set up more physiological experimental conditions. Moreover, EGF was added for short-term exposures in order to evaluate the responsiveness of each signalling component to its physiological ligand in untreated and IFN α -treated KB cells. We have firstly studied the activity of Ras through the affinity precipitation of Ras with the minimal binding domain of Raf-1. We have found that the levels of activated Ras (linked to the minimal binding domain of Raf-1) were increased in IFN α -treated cells. In fact, the activated Ras levels were about two-fold higher in IFN α -treated cells than in control cells, and the stimulation of Ras activity by exposure to EGF for 10 min was further enhanced in IFN α -treated cells reaching levels 3.5-fold higher than untreated controls (Figure 3). At longer times of exposure to EGF, Ras activity decreased either in EGF- or EGF/IFN α -treated cells (Figure 3). Therefore, in IFN α -treated KB cells, the increased activity of EGF-R signalling was indeed coupled to enhanced Ras activity and was further stimulated by the addition of exogenous EGF.

Effects of IFN α and EGF on Raf-1 expression and activity

We have evaluated the effect of IFN α and EGF on Raf-1 activity by a kinase assay using histone H1 as substrate of




Figure 1 EGF antagonizes apoptosis induced by IFN α in human epidermoid cancer cells. KB cells were seeded and treated with IFN α and/or EGF as described in the Materials and methods section. (a) The internucleosomal DNA fragmentation was assessed as described in the Materials and methods section. (b) The percent of apoptotic cells, evaluated by FACS analysis, is shown as bars. **CTR**: untreated cells; **IFN α** : 48 h 1000 IU/ml IFN α ; **EGF**: 12 h 10 nM EGF; **IFN α +EGF**: 48 h 1000 IU/ml IFN α +12 h 10 nM EGF. The experiment was performed three times and S.E.s were always less than 5%. (c) Evaluation of the expression of p53, c-myc and β -actin. Proteins were extracted and run on SDS-PAGE as described in 'Materials and methods section'. The experiment was performed at least three times and the results were always similar. **CTR**: untreated cells; **IFN α** : 48 h 1000 IU/ml IFN α ; **EGF**: 12 h 10 nM EGF; **IFN α +EGF**: 48 h 1000 IU/ml IFN α +12 h 10 nM EGF; **VP-16**: 18 h 50 μ M etoposide

Raf-1, and we have found that Raf-1 activity is increased either by EGF or IFN α and is potentiated after EGF addition for 10 min to KB cells exposed to IFN α (Figure 4). Moreover, we

have determined Raf-1 levels by Western blotting, and found that they were almost unmodified by the different treatments (Figure 4). These experiments demonstrate that an increase of basal and EGF-induced enzymatic activity of Raf-1 can be recorded downstream of the EGF-Ras signalling in IFN α -treated cells.

Effects of IFN α and EGF on Erk-1/2 activity

On the basis of the finding that IFN α induced a hyperactivation of Mek-1, we have evaluated the effects of this cytokine on the activity of Erk-1/2, the downstream enzymatic targets of Raf-1. We found that the exposure of KB cells to IFN α for short times (5 min–12 h) had no effects on MAPK activity, while they were maximally activated after 48 h of treatment with the cytokine (Figure 5a, lower panel). The latter effect paralleled EGF-R upregulation, which reached a peak after 48 h of exposure to IFN α (Figure 5a, upper panel and see also ref. 24). Therefore, the timing of MAPK²⁵ activation by IFN α overlapped that of EGF-R upregulation, suggesting that the stimulation of MAPK activity could be a consequence of the increased surface expression of EGF-R with subsequent amplification of downstream signalling.

We have subsequently evaluated the kinetics of Erk-1/2 activation by EGF in untreated and IFN α -treated KB cells. We again found an almost two-fold increase of the two enzyme activities in 48 h 1000 IU/ml IFN α -treated cells. In both untreated and 48 h 1000 IU/ml IFN α -treated KB cells, the maximal activation of Erk-1/2 occurred after 10 min of exposure to 10 nM EGF, as evaluated with an immunoprecipitated kinase assay (Figure 5b and c). However, the exposure to IFN α caused an almost two-fold increase of the maximal activation of Erk-1/2 induced by EGF (Figure 5b and c). Therefore, 48 h exposure to IFN α increased the activity of Erk-1/2 without affecting the sensitivity to further stimulation by exogenous EGF.

IFN α enhanced the apoptosis induced by dominant negative Ras plasmid RASN17 in KB cells

We have shown that the EGF-dependent Ras-> Erk pathway was hyperactivated in cells exposed to IFN α . In order to demonstrate that such signalling could indeed retain an antiapoptotic activity in the cells exposed to IFN α , we have selectively inhibited Ras activity by a dominant negative mutant. KB cells have been cotransfected with a plasmid encoding for a dominant negative *Ras* RASN17 and with the *E. coli* β -galactosidase reporter gene. The cells were analyzed by FACS after PI labelling as described above. In all cases, the cotransfection resulted in specific green fluorescence in a fraction of the cell population owing to the expression of β -galactosidase (see Figure 6 β Gal NEG versus β Gal POS). The latter effect allowed us the gating of nontransfected (G2) or transfected cells (G1) in order to perform cell cycle analysis (red fluorescence determined by PI labelling). We have found that β -galactosidase expression alone did not cause apoptosis (2 and 6% of apoptosis either in nontransfected or β -galactosidase-transfected cells, respectively) (Figure 6a, G2




Figure 2 Modulation of caspases 3, 6, 8 and 9 by IFN α and EGF. (b) KB cells were seeded and treated with 1000 IU/ml IFN α for the indicated times. The specific caspase activity was determined with a fluorimetric assay as described in the 'Materials and methods section'. Caspase 3 (●); caspase 6 (■); caspase 8 (▽); caspase 9 (○). The experiment was performed three times and S.D.s were always less than 5%. Bars, s.d. (b–d) KB cells were seeded and treated with 1000 IU/ml IFN α for the indicated times. EGF (10 nM) was added to the cells for the last 12 h of incubation alone or in combination with IFN α . The specific caspase activity was determined with a fluorimetric assay as described in the 'Materials and Methods section'. Caspase 3 (b); caspase 6 (c); caspase 8 (d); caspase 9 (e). 1000 IU/ml IFN α (▨); 10 nM EGF (■); IFN α +12 h 10 nM EGF (□). The experiment was performed three times and S.D.s were always less than 5%. Bars, S.D.

and G1). In the cells cotransfected with RASN17 and β -galactosidase almost 29% of apoptotic cells were detected (Figure 6b, G1), while only 2% of apoptosis was found in nontransfected cells (Figure 6b, G2). The exposure of β -galactosidase-transfected cells to 1000 IU/ml IFN α for 24 h induced apoptosis in 16% of the cell population (Figure 6c, G1) against 14% of apoptotic cells in nontransfected population (Figure 6c, G2). The treatment of β -galactosidase and RASN17 cotransfected cells with 1000 IU/ml IFN α induced an almost 76% of apoptosis (Figure 6D, G1), while programmed cell death was found in only 13% of the nontransfected cell

population (Figure 6D, G2). We have performed these experiments after 24 h of exposure to IFN α because at this time maximal β -galactosidase expression occurred. Moreover, almost 80% of cells were apoptotic after 48 h from the transfection with RASN17 (data not shown). In this experiment about 30% of the cell population was transfected with β -galactosidase plasmid. It was therefore demonstrated that the abrogation of Ras activity by a transfected dominant negative mutant sensitized tumor cells to IFN α -induced apoptosis, indicating an antiapoptotic function of Ras in this experimental model.




Figure 3 Effects of EGF and IFN α on Ras activity. (Upper gel) Affinity precipitation of Ras performed with the minimal binding domain of Raf-1 conjugated with agarose as described in the 'Materials and methods section'. **CTR**, untreated; **IFN α** , 48 h 1000 IU/ml IFN α ; **E10'**, 10 min 10 nM EGF; **I/E 10'**, 48 h 1000 IU/ml IFN α and 10 min 10 nM EGF; **E1 h**, 1 h 10 nM EGF; **I/E 1 h**, 48 h 1000 IU/ml IFN α and 1 h 10 nM EGF; **Asp.** minimal binding domain of Raf-1 incubated only with Mab raised against Ras without cell extracts. (Lower gel) Western blot assay for the expression of the total Ras protein. **CTR**, untreated; **IFN α** , 48 h 1000 IU/ml IFN α ; **E10'**, 10 min 10 nM EGF; **I/E 10'**, 48 h 1000 IU/ml IFN α and 10 min 10 nM EGF; **E1 h**, 1 h 10 nM EGF; **I/E 1 h**, 48 h 1000 IU/ml IFN α and 1 h 10 nM EGF. The experiments were performed at least three different times and the results were always similar. (Bars) Quantitization of the intensity of the different bands with an appropriate apparatus (Bio-Rad) and expression as arbitrary units. **CTR**, untreated; **IFN α** , 48 h 1000 IU/ml IFN α ; **E10'**, 10 min 10 nM EGF; **I/E 10'**, 48 h 1,000 IU/ml IFN α and 10 min 10 nM EGF; **E1 h**, 1 h 10 nM EGF; **I/E 1 h**, 48 h 1000 IU/ml IFN α and 1 h 10 nM EGF; **Asp.** minimal binding domain of Raf-1 incubated only with Mab raised against Ras without cell extracts.

MEK-1 inhibitor PD098059 specifically abrogated recovery from IFN α -induced apoptosis by EGF

Since Ras is implicated in the triggering of several different signal transduction pathways, we have verified that the integrity of the downstream ERK-targeted pathway was




Figure 4 Effects of IFN α and EGF on Raf-1 activity and expression. KB cells have been cultured for 48 h in the absence or presence of 1000 IU/ml IFN α with or without the addition of 10 nM EGF for the last 5 min as described in the figure by + and -. Raf-1 was immunoprecipitated and analyzed for kinase activity by immunocomplex kinase assay using histone H1 as substrate. Lane C1 indicated a control where the immunocomplex kinase assay was performed on untreated EGF-stimulated cells in the absence of the substrate. Lane C2 indicated a control assay performed using nonimmune serum to immunoprecipitate untreated EGF-stimulated cells. At the same time, the cells were also processed for the determination of Raf-1 expression as described in the 'Materials and Methods section'.

indeed required for the antiapoptotic effects of EGF.²⁶ Therefore, we have evaluated if the specific inhibition of Mek-1 could also potentiate the growth inhibition and apoptosis induced by IFN α and if the Ras \rightarrow Erk-1/2 pathway plays a crucial role in the survival of epidermoid cancer cells exposed to IFN α . Only 5% untreated KB cells were apoptotic as assessed by FACS and *in situ* TUNEL analysis (Figure 7a). Again 48 h 1000 IU/ml IFN α induced 30% apoptosis and 12 h 10 nM EGF antagonized this effect (only 8% apoptotic cells were recorded under these experimental conditions) (Figure 7b and e and relative insets, respectively). In total, 50 μ M PD098059 and 10 nM EGF alone induced 35 and 15% apoptosis, respectively (Figure 7c and d, respectively). PD098059 was again able to enhance the effect induced by IFN α causing programmed cell death in about 80% of cells, thus suggesting a supra-additive effect induced by the addition of MEK-1 inhibitor to IFN α (Figure 7f). Moreover, 50 μ M PD098059 abrogated the anti-apoptotic effects of EGF in IFN α -treated cells since an almost 30% of apoptosis was recorded in cells exposed to IFN α , EGF and PD098059 (Figure 7h). On the other hand, PD098059 caused almost no effects on the apoptosis induced by EGF (Figure 7g). Interestingly, PD098059 reduced MAPK phosphorylation almost completely already after 6 h in KB cells (Figure 7i). All together, these data suggest that the antiapoptotic effects induced by EGF occurred through the activation of Erk, which is therefore the final target of an antiapoptotic kinase cascade.

Discussion

IFN α is a cytokine that has shown a well-defined but still limited activity against human tumors.¹⁻⁵ Resistance to IFN α has been demonstrated to be in part based on receptor or post-receptor signalling defects in the tumor cells.⁶⁻⁸ However, the cancer cell escape from the antiproliferative activity of IFN α and the mechanisms by which this cytokine causes





Figure 5 IFN α increases Erk-1 and Erk-2 activity in human epidermoid cancer cells. (A) upper panel) ^{125}I -EGF binding on human epidermoid KB cells. KB cells were treated for different times with 1000 IU/ml IFN α . At the indicated times, cells were processed for the determination of the ^{125}I -EGF specific binding, as described in the 'Materials and methods section'. The values were expressed as % of control. The experiments were performed three times and SDs were always less than 5%. (a, lower panel) KB cells were cultured for the indicated times in the absence or presence of 1000 IU/ml IFN α . Then the cells were processed for the determination of the expression and phosphorylation of Erk-1 and -2 evaluated after blotting with an anti-MAPK and an anti-pMAPK specific Mab, respectively, as described in the 'Materials and methods section'. The experiments were performed at least three different times and the results were always similar. (b and c) KB cells were cultured for 48 h in the absence or presence of 1000 IU/ml IFN α with or without the addition of 10 nM EGF for different times. Then an immunocoupled kinase assay of (b) Erk-1 and (c) Erk-2 was performed as described in 'Materials and Methods section'. Untreated KB cells (■); IFN α -treated KB cells (●). The experiment was performed three times and s.d.s were always less than 5%. Bars, s.d.

direct antitumor effect remain still unsolved. Apoptosis induction appears to be a way by which IFN α exerts its antiproliferative effects.⁹⁻¹² It is well established that tumor cells overcome the natural propensity to undergo apoptosis with the overexpression and/or overactivation of growth factor-induced signal transduction pathways.¹³⁻¹⁵ EGF-R overexpression is a common event in epidermoid cancers and is considered an unfavorable prognostic factor.²⁷ It has also been demonstrated that *Erk-1/2* antagonize apoptosis in several experimental models.^{22,28-30} We have previously reported that IFN α induces growth inhibition and increases the expression and function of EGF-R in human epidermoid cancer cells.^{23,24} Moreover, we have demonstrated that the growth inhibition induced by IFN α is, at least in part, because of the apoptosis triggered by a stress response leading to the activation of JNK-1, and that EGF antagonizes growth inhibition, apoptosis and the biochemical events induced by the cytokine.¹⁰ On the basis of these findings, we have

hypothesized that the increased expression and function of the EGF-R could be part of a survival response that could protect tumor cells from the apoptosis triggered by IFN α . In this paper, we have investigated the molecular bases of the counteracting effects of IFN α and EGF on apoptosis. We have firstly studied the involvement of caspases in the execution of the apoptotic process. We have found that IFN α induced an activation peak of caspases 6 and 8 after 24 h, while the maximal activation of caspase 3 was recorded 24 h later. The differential timing of caspase activation is in agreement with the sequence of activation of these proteases during the apoptotic process and suggests an involvement of caspases in the execution of apoptosis induced by IFN α .³¹ Our findings are, moreover, in agreement with the recent observations by Thyrell *et al.*,³² who found induction of caspase activity by IFN α in apoptotic tumor cells. In our experimental conditions, we have found that the addition of EGF to IFN α -treated KB cells completely antagonized apoptosis induction and the

triggering of the caspase cascade, suggesting that the EGF-R signalling suppresses apoptosis, thereby antagonizing the activation of its executioners. These results appear also in line with the recent findings demonstrating the involvement of

growth-factor-dependent pathways in the protection from caspase activation induced by Bad overexpression. Moreover, it has been demonstrated that the EGF-R-dependent pathway controls keratinocyte survival and the expression of



the proapoptotic bcl-x_L expression through a Mek-dependent pathway.³⁰

We have then investigated the involvement of EGF-mediated intracellular signalling in the protection of epidermoid cancer cells from IFN α -induced apoptosis. We have found that the EGF- and Ras-dependent MAPK cascade is hyperactivated in IFN α -treated cells and could be further stimulated by the addition of EGF. In fact, we have demonstrated an increased activity and responsiveness to EGF stimulation of Ras, Raf-1 and *Erk-1/2* in KB cells exposed to IFN α . These findings suggest that the EGF-R function is preserved in IFN α -treated cells, taking also into account the activation threshold of the EGF-dependent signalling. Interestingly, maximal MAPK hyperactivation occurred after 48 h from the beginning of the treatment with IFN α . It must be considered that the maximal upregulation of EGF-R on the cell surface occurs with the same timing, suggesting that overstimulation of the MAPK pathway could be produced by the amplification of the EGF signalling owing to the increased expression of the receptor.^{23,24} We have previously described that other antiproliferative agents, such as cytosine arabinoside, 5aza-2'-deoxycytidine and 8-chloro-cAMP (8ClcAMP), also increase EGF-R expression on KB cells.³³⁻³⁵ On this basis, we have hypothesized that the up-regulation of growth factor receptors is a common event in growth-inhibited tumor cells and could represent a protective response towards the antiproliferative stimuli.³⁶ Also in the case of 8ClcAMP, the EGF-induced signalling MAPK is amplified likely as a consequence of the increased expression of EGF-R.³⁵ However, MAPK activity is reduced in 8Cl-cAMP-treated KB cells, suggesting a selective inhibition of Erks or of a still unknown upstream activator induced by the drug.³⁵ It has to be considered that chronic activation of Erk has been reported to induce apoptosis in eukaryotic cells.³⁷⁻⁴⁰ However, we can exclude that IFN α -induced apoptosis occurs through the hyperactivation of Erks on the basis of both the timing of Erk activation and the effects of the combined treatment of KB cells with IFN α and EGF. In fact, we have found that the exposure of IFN α -treated KB cells to EGF did not simply induce a recovery of cell growth, but it also caused a proliferative effect much greater than that recorded in KB cells exposed to EGF alone.²⁴

The involvement of the Ras->MAPK pathway in the protection of KB cells from the apoptosis induced by IFN α is further demonstrated by both Ras inactivation by RASN17 transfection and Mek-1 inhibition by exposure to PD098059. In fact, the transfection of RASN17 in KB cells caused apoptosis as demonstrated by the FACS analysis performed on transfected cells and also potentiated the apoptosis

induced by IFN α . These data suggest that the integrity of Ras function was necessary to produce an anti-apoptotic signal that mediates a survival response in cells exposed to IFN α via Erk-1/2 activation. In fact, we have demonstrated that Ras-dependent survival signalling targets Erk-1/2 since the reduction of MAPK activity by PD098059 enhanced apoptosis caused by IFN α . An additional important finding is that PD098059 specifically abrogated the recovery from apoptosis induced by EGF in IFN α -treated cells. A second important antiapoptotic pathway involves signalling via Akt/PKB.^{41,42} In fact, it has been demonstrated that Akt can be activated concomitantly or independently from Ras->ERK-1/2 signalling by growth factors.⁴³⁻⁴⁵ Additionally, it has been reported that PC12 cells display a protective antiapoptotic pathway in response to hypoxic stimuli.⁴⁶ The protection from apoptosis by Akt could be because of the regulation of mitochondrial physiology since Akt is involved in the regulation of bcl-related proteins such as Mcl-1.⁴³ However, the requirement of Akt for the protection from apoptotic events is highly variable depending upon the experimental model used.^{44,45,47} In our experimental conditions, we have demonstrated that PD098059 completely antagonized the antiapoptotic effects of EGF. Moreover, we have preliminarily found that the specific EGF-R-associated kinase inhibitor ZD1839 (IRESSA) synergizes with IFN α in inducing the growth inhibition and apoptosis of several human epidermoid cancer cell lines, which is coupled to complete inhibition of ERK activity (A Budillon *et al.*, manuscript in preparation). Therefore, our results suggest that the activation of Ras->Raf-1->Mek1->Erk-1/2 signalling has a prominent role in the antiapoptotic effects exerted by EGF in epidermoid cancer cells exposed to IFN α , providing evidence of the potential benefits of the molecular interference with this pathway. However, the occurrence of other survival pathways will warrant further investigations and we cannot presently completely exclude a role of the Akt pathway in the modulation of apoptosis of KB cells.

Consistent with our results, additional findings have been recently reported on the interaction between the Ras->Erk pathway and the myc oncogene on apoptosis onset.⁴⁸ In fact, it has been recently found that the activation of the Ras/Mek signalling suppresses myc-dependent apoptosis in cells with MAPK cascade amplification because of Ras activation.⁴⁸ In our experimental model, we have also found increased expression of p53 and c-myc in cells exposed to IFN α while the EGF anti-apoptotic effects were paralleled by down-regulation of myc and p53 expression.

In conclusion, we have provided evidence that IFN α makes the survival of human epidermoid cancer cells dependent on

Figure 6 IFN α enhances the apoptosis induced by the dominant negative Ras plasmid RasN17 in KB cells. KB cells were seeded, electroporated with dominant negative Ras plasmid RASN17 and/or β -galactosidase plasmid, and treated with IFN α for 24 h as described above. Then the cells were fixed in methanol, incubated with anti- β -galactosidase MA b and anti-mouse FITC rabbit antiserum and analyzed after DNA labelling with propidium iodide. Cell cycle analysis was performed after appropriate gating on cell population expressing β -galactosidase as described in the 'Materials and methods section'. (a) Parental untreated cells (CTR); (b) RASN17 transfected cells (RASN17); (c) 24 h 1000 IU/ml IFN α -treated parental cells (IFN α 24 h); (D) 24 h 1000 IU/ml IFN α -treated RasN17 transfected cells (IFN α 24 h+RASN17). β Gal NEG, cells exposed to an irrelevant mouse IgG and subsequently labelled with an anti-mouse FITC rabbit antiserum. β Gal POS, cells exposed to an anti- β -galactosidase mouse IgG and subsequently labelled with an anti-mouse FITC rabbit antiserum. G1, gating of cells expressing β -galactosidase; G2, gating of cells not expressing β -galactosidase. The lower histograms show the analysis of cell cycle performed on the G2 and G1 population, respectively. The apoptotic cells were shown as a percentage of total cell population indicated by a bar. The experiments were performed at least three different times and the results always gave less than 5% s.d. In each experimental point 10,000 events were analyzed.




Figure 7 The antiapoptotic effects of EGF are mediated by the selective activation of ERK. KB cells were seeded and treated with IFN α and/or 50 μ M PD098059 for 48 h and/or 10 nM EGF for 12 h. At the time of the experiment, cells were fixed and processed with TUNEL techniques as described in the 'Materials and methods section'. The experiments were performed at least three different times and the results were always similar. (a) Control cells; (b) 48 h 1000 IU/ml IFN α -treated cells; (c) 48 h 50 μ M PD098059-treated cells; (d) 12 h 10 nM EGF-treated cells; (e) 48 h 1000 IU/ml IFN α - and 12 h 10 nM EGF-treated cells; (f) 48 h 1000 IU/ml IFN α - and 48 h 50 μ M PD098059 treated cells; (g) 12 h 10 nM EGF- and 48 h 50 μ M PD098059-treated cells; (h) 48 h 1000 IU/ml IFN α - and 48 h 50 μ M PD098059-treated cells and 12 h 10 nM EGF. *Insets*: The cells were seeded and treated with the different substances as described above. At the time of the experiment KB cells were FACS analyzed after detachment and DNA labelling with propidium iodide as described in the 'Materials and methods section'. The experiments were performed at least three different times and the results always gave less than 5% SD. In each experimental point, 10,000 events were analyzed. (i) The expression and phosphorylation of Erk-1 and -2 was evaluated after blotting with an anti-p-MAPK (upper gel) and an anti-MAPK-specific MAb (lower gel) in KB cells treated for different times with 50 μ M PD098059. The experiments were performed at least three different times and the results were always similar

Ras/Erk signalling, which acts as a survival pathway. The specific disruption of the latter could be a useful approach to potentiate the antitumor activity of IFN α against human epidermoid tumors as indicated by the results achieved by selective interference with Ras and Mek-1 function in cells exposed to IFN α .

Materials and Methods

Materials

DMEM, BSA and FBS were purchased from Flow Laboratories (Milan, Italy). Tissue culture plasticware was from Becton Dickinson (Lincoln Park, NJ, USA). IFN α was a gift of Schering (Schering-Plough, NJ, USA).

Receptor grade EGF and protein Sepharose were purchased from Sigma (St. Louis, MO, USA). Rabbit antisera raised against Raf-1 C-12, β actin, Erk-1 K-23 and Erk-2 MAb C-14 were purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Anti-pan-Ras MAb clone 10 was purchased from Calbiochem.

Cell culture

The human oropharyngeal epidermoid carcinoma KB cell line, obtained from the American Type Tissue Culture Collection, Rockville, MD, USA, was grown in DMEM supplemented with heat-inactivated 10% FBS, 20 mM HEPES, 100 U/ml penicillin, 100 μ g/ml streptomycin, 1% L-glutamine and 1% sodium pyruvate. The cells were grown in a humidified atmosphere of 95% air/5% CO $_2$ at 37°C.

Western blot analysis

KB cells were grown for 48 h with or without IFN α and were thereafter exposed for the indicated times to 10 nM EGF at 37°C. For cell extract preparation, the cells were washed twice with ice-cold PBS/BSA, scraped, and centrifuged for 30 min at 4°C in 1 ml of lysis buffer (1% Triton, 0.5% sodium deoxycholate, 0.1 NaCl, 1mM EDTA, pH 7.5, 10 mM Na₂HPO₄, pH 7.4, 10 mM PMSF, 25 mM benzamidin, 1 mM leupeptin, 0.025 U/ml aprotinin). Equal amounts of cell proteins were separated by SDS-PAGE. The proteins on the gels were electrotransferred to nitrocellulose and reacted with the different MABs.

Affinity precipitation of Ras

KB cells were treated with IFN α and EGF as described above. The cells were lysed in Mg²⁺ buffer containing 20 mM HEPES, pH 7.5, 150 mM NaCl, 1% Igepal CA-630, 10 mM MgCl₂, 1 mM EDTA and 2% glycerol. Then, 10 μ l Ras binding domain (RBD) conjugated to agarose was added to 1 mg of cell lysate and the mixture was incubated at 4°C for 1 h. The agarose beads were collected by microcentrifugation at 14,000 \times g for 5 s. and washed three times with Mg²⁺ buffer. The agarose beads were boiled for 5 min in 2 \times Laemmli sample buffer and collected by a microcentrifuge pulse. The supernatants were run on 12% SDS-PAGE, then the proteins were electrotransferred on a nitrocellulose film. The nitrocellulose was incubated overnight with 1 μ g/ml of anti-Ras Mab, clone RAS10 and with a secondary Mab, a goat α -mouse HRP conjugated IgG, for 1.5 h. The film was washed with PBS/0.05% Tween 20 and detected by ECL, chemiluminescence's technique (Amersham).

Internucleosomal DNA fragmentation (ladder)

For all apoptosis evaluation experiments (gel ladder and FACS analysis), both attached and suspended cells were collected prior to the processing. DNA fragmentation was measured after extraction of low molecular weight DNA. Briefly, 10 \times 10⁶ cells were resuspended in 900 μ l 1 \times Tris-EDTA buffer and lysed with 25 μ l 20% SDS. DNA was precipitated in ethanol for 6 h in the presence of 5 M NaCl. The high molecular weight fraction was sedimented by high-speed centrifugation, and the fragmented DNA was extracted from the aqueous phase with phenol and chloroform and then precipitated with ethanol. After resuspension in water, DNA was electrophoresed using 1.5% agarose gel and visualized by ultraviolet light following ethidium bromide staining.

Evaluation of apoptosis by DNA-flow cytometry

Cells were centrifuged and directly stained in a propidium iodide (PI) solution (50 μ g PI in 0.1% sodium citrate, 0.1% NP40, pH 7.4) overnight at 4°C in the dark. Flow cytometric analysis was performed using a FACScan flow cytometer (Becton Dickinson, San Jose, CA, USA) interfaced with a Hewlett-Packard computer (mod. 310) for data analysis. To evaluate cell apoptosis, PI fluorescence was collected as FL2 (log scale) by the CellFIT software (Becton Dickinson). For the evaluation of apoptosis after transfection with RASN17 and/or β -galactosidase plasmids, cells were fixed in methanol and then incubated with an anti- β -galactosidase MAB for 1 h at 4°C and labelled with an FITC anti-mouse rabbit antibody for 1 h. Thereafter, the cells were stained with PI as described above. In the latter case, apoptosis and cell cycle analysis were performed on the whole cell population and on β -galactosidase expressing cells after appropriate gating. For the evaluation of intracellular DNA content, at least 10,000

events for each point were analyzed in at least three different experiments giving a s.d. less than 5%.

MAPK assay

KB cells were cultured and treated as described above. Then the cells were washed twice with ice-cold PBS, scraped and lysed for 1 h at 4°C in the following buffer: 10 mM TRIS, 150 mM NaCl, 2 mM EGTA, 2 mM DTT, 1 mM sodium orthovanadate, 1 mM PMSF, 10 μ g/ml aprotinin, 10 μ g/ml leupeptin and 1% glycerol. Equalized amounts of proteins from lysates of KB cells (50 μ l) were incubated for 90 min at 4°C with 50 μ l of protein A-Sepharose conjugated to 25 μ g anti-Erk-1 or anti-Erk-2 MABs. The suspension was centrifuged at 14,000 \times g, the supernatant discarded, and the immunoprecipitated washed twice with 250 μ l of the following buffer: 20 mM Tris-acetate, pH 7.0 (20°C), 0.27 M sucrose, 1 mM EDTA, 1 mM EGTA, 1 mM sodium orthovanadate, 10 mM sodium glycerophosphate, 50 mM sodium fluoride, 5 mM sodium pyrophosphate, 1% Triton X-100, 0.1% 2-mercaptoethanol, 1 mM benzamidin, 0.2 mM phenylmethylsulfonyl fluoride and 0.5 M NaCl. The immunoprecipitates were washed once with 250 μ l of the following buffer: 50 mM Tris-HCl, pH 7.5 (20°C), 0.03% Brij-35, 0.1 mM EGTA and 0.1% 2-mercaptoethanol. Then the immunoprecipitates were incubated for 30 min at 30°C with 5 μ l of a solution containing 1.2 mM Mg⁺⁺ ATP with 200 μ Ci/ml of [γ -³²P]ATP and 15 μ l of a solution containing a peptide that is specifically phosphorylated by MAPK (Amersham, Biotrak, Milan, Italy).²⁵ The reaction was stopped with the addition of 2.94% (w/v) orthophosphoric acid and red carmosin solution and then microfuged for 15 s. The reaction mixtures were spotted onto phosphocellulose filters (Whatman P81) and washed three times in 1% acetic acid. Filters were air-dried and then counted by liquid scintillation using Omnifluor/toluene (DuPont-New England Nuclear, Boston, MA, USA).

Raf-1 kinase assay

Assay for Raf-1 kinase activity was performed by immunocomplex kinase assay as already described for MAPK using anti-Raf-1 antiserum. H1 histone (10 μ g) (Upstate Biotechnology Inc.) was used as substrate in 30 μ l of buffer A containing 10 μ Ci of [γ -³²P]ATP, the reaction was allowed to proceed for 30 min at 30°C, and the proteins were separated by SDS-12.5% PAGE.

Electroporation

Cells were detached from confluent 100 mm- dishes. Cells (100 \times 10⁶) were incubated in appropriate electroporation vials with 800 μ l of electroporation buffer (20 mM HEPES, 137 mM NaCl, 5 mM KCl, 0.7 mM Na₂HPO₄ and 6 mM glucose) and 15 μ g of the RASN17 DNA and/or 5 μ g/ml β -galactosidase DNA in 20 mM HEPES. Then cells were electroporated at 250 V and at 975 μ F for 6 s. The cells were incubated at 37°C with or without 1000 IU/ml IFN α for 24 h. After the incubation, the cells were processed for FACS analysis as described above.

TUNEL technique

For TUNEL assay, after washing in PBS supplemented in 0.1% BSA, cells were treated with an *in situ* detection kit, according to the manufacturers instructions (Boehringer Mannheim Biochemicals). Nuclei with fragmented DNA were visualized by a fluorescence microscope.

Determination of caspase activity

Cells were seeded and treated with 1000 IU/ml IFN α for different times and/or 10 nM EGF for 12 h. At the time of caspase determination, cells were lysed. For each reaction, 50 μ l of cell lysate was added to 1 ml reaction mixture composed of reaction buffer, DTT 10 mM and the specific substrate peptide (DEVD for caspase-3, VEID for caspase-6, IETD for caspase-8 and LEHD for caspase-9) conjugated to 7-amino-4-trifluoromethylcoumarin (AFC). The mixtures were incubated for 1 h at 37°C. The levels of free AFC were measured using a spectrofluorometer with an excitation wavelength of 400 nm and an emission wavelength range of 480–520 nm (peak at 505 nm).

Acknowledgments

We thank Prof. N Perrotti and G Cuda of Catanzaro University for the careful reading of the paper and Ms. G Granata and Dr. P Sands for their help in the manuscript preparation. This work was partially supported by the Italian Association for Cancer Research (AIRC) and by grants from the Italian Ministry for Research (PRIN2001) and the Italian Ministry of Health (FSN99 and FSN2000).

References

- Agarwala SS and Kirkwood JM (1998) Adjuvant therapy of melanoma. *Semin. Surg. Oncol.* 14: 302–310
- Vlock DR Andersen J Kalish LA Johnson JT Kirkwood JM Whiteside T Herberman RB Adams GS Oken MM and Haselow RE (1996) Phase II trial of interferon-alpha in locally recurrent or metastatic squamous cell carcinoma of the head and neck: immunological and clinical correlates. *J. Immunother. Emphasis Tumor Immunol.* 19: 433–442
- Kirkwood JM, Resnick GD and Cole BF (1997) Efficacy, safety, and risk-benefit analysis of adjuvant interferon alfa-2b in melanoma. *Semin. Oncol.* 24: 16–23
- Kirkwood JM Strawderman MH Ernstoff MS Smith TJ Borden EC and Blum RH (1996) Interferon alfa-2b adjuvant therapy of high-risk resected cutaneous melanoma: the Eastern Cooperative Oncology Group Trial EST 1684. *J. Clin. Oncol.* 14: 7–17
- Gutterman JU (1994) Cytokine therapeutics: lessons from interferon alpha. *Proc. Natl. Acad. Sci. USA* 91: 1198–1205
- Wong LH Kraver KG Hatzinisiriou I Estcourt M J Hersey P Tam ND Edmondson S Devenish RJ and Ralph SJ (1997) Interferon-resistant human melanoma cells are deficient in ISGF3 components, STAT1, STAT2, and p48-ISGF3gamma. *J. Biol. Chem.* 272: 28779–28785
- Ransohoff RM. (1998) Cellular responses to interferons and other cytokines: the JAK-STAT paradigm. *N. Engl. J. Med.* 338: 616–618
- Harvat BL Jetten AM and Seth P (1997) The role of p27Kip1 in gamma interferon-mediated growth arrest of mammary epithelial cells and related defects in mammary carcinoma cells. *Oncogene*, 14: 2111–2122
- Rodriguez-Villanueva J and McDonnell TJ (1995) Induction of apoptotic cell death in non-melanoma skin cancer by interferon-alpha. *Int. J. Cancer* 61: 110–114
- Caraglia M Abbruzzese A Leardi A Pepe S Budillon A Baldassarre G Selleri C De Lorenzo S Fabbrocini A Giuberti G Vitale G Lupoli G Bianco AR and Tagliaferri P (1999) Interferon-alpha induces apoptosis in human KB cells through a stress-dependent mitogen activated protein kinase pathway that is antagonized by epidermal growth factor. *Cell Death Differ.* 6: 773–780
- Roth W Wagenknecht B Dichgans J and Weller MJ (1998) Interferon-alpha enhances CD95L-induced apoptosis of human malignant glioma cells. *J. Neuroimmunol.* 87: 121–129
- Tanaka N Sato M Lamphier MS Nozawa H Oda E Noguchi S Schreiber RD Tsujimoto Y and Taniguchi T (1998) Type I interferons are essential mediators of apoptotic death in virally infected cells. *Genes Cells* 3: 29–37
- Tilly JL Billig H Kowalski KI and Hsueh AJ (1992) Epidermal growth factor and basic fibroblast growth factor suppress the spontaneous onset of apoptosis in cultured rat ovarian granulosa cells and follicles by a tyrosine kinase-dependent mechanism. *Mol. Endocrinol.* 6: 1942–1950
- Peng H Wen TC Tanaka J Maeda N Matsuda S Desaki J Sudo S Zhang B and Sakanaka M (1998) Epidermal growth factor protects neuronal cells in vivo and *in vitro* against transient forebrain ischemia- and free radical-induced injuries. *J. Cereb. Blood Flow Metab.* 18: 349–360
- Stoll SW Benedict M Mitra R Hiniker A Elder JT and Nunez G (1998) EGF receptor signalling inhibits keratinocyte apoptosis: evidence for mediation by Bcl-XL. *Oncogene* 16: 1493–1499
- Sibilia M Fleischmann A Behrens A Stingl L Carroll J Watt FM Schlessinger J and Wagner EF (2000) The EGF receptor provides an essential survival signal for SOS-dependent skin tumor development. *Cell*, 102: 211–220
- Widmann C Gibson S Jarpe MB and Johnson GL (1999) Mitogen-activated protein kinase: conservation of a three-kinase module from yeast to human. *Physiol. Rev.* 79: 143–180
- Garrington TP and Johnson GL (1999) Organization and regulation of mitogen-activated protein kinase signaling pathways. *Curr. Opin. Cell Biol.* 11: 211–218
- Xia Z Dickens M Raingeaud J Davis RJ and Greenberg ME (1995) Opposing effects of ERK and JNK-p38 MAP kinases on apoptosis. *Science* 270: 1326–1331
- Yan CYI and Greene LA (1998) Prevention of PC12 cell death by *N*-acetylcysteine requires activation of the Ras pathway. *J. Neurosci.* 18: 4042–4049
- Aikawa R Komuro I Yamazaki T Zou Y Kudoh S Tanaka M Shiojima I Hiroi Y and Yazaki Y (1997) Oxidative stress activates extracellular signal-regulated kinases through Src and Ras in cultured cardiac myocytes of neonatal rats. *J. Clin. Invest.* 100: 1813–1821
- Stadheim TA and Kucera GL (1998) Extracellular signal-regulated kinase (ERK) activity is required for TPA-mediated inhibition of drug-induced apoptosis. *Biochem. Biophys. Res. Commun.* 245: 266–271
- Budillon A Tagliaferri P Caraglia M Torrisi MR Normanno N Iacobelli S Palmieri G Stoppelli MP Frati L and Bianco AR (1991) Upregulation of epidermal growth factor receptor induced by alpha-interferon in human epidermoid cancer cells. *Cancer Res.*, 51: 1294–1299
- Caraglia M Leardi A Corradino S Ciardiello F Budillon A Guarasi R Bianco AR and Tagliaferri P (1995) alpha-Interferon potentiates epidermal growth factor receptor-mediated effects on human epidermoid carcinoma KB cells. *Int. J. Cancer* 61: 342–347
- Clark-Lewis I Sanghera JS and Pelech SL (1991) Definition of a consensus sequence for peptide substrate recognition by p44mpk, the meiosis-activated myelin basic protein kinase. *J. Biol. Chem.* 266: 15180–15184
- Shields JM Pruitt K Mc Fall A Shaub A and Der CJ (2000) Understanding Ras: it ain't over til it's over. *Trends Cell. Biol.* 10: 147–154
- Gullick WJ Marsden JJ and Whittle N (1986) Expression of epidermal growth factor receptors on human cervical, ovarian, and vulval carcinomas. *Cancer Res.* 46: 285–292
- Foncea R Galvez A Perez V Morales MP Calixto A Melendez J Gonzalez-Jara F Diaz-Araya G Sapag-Hagar M Sugden PH LeRoith D and Lavandero S (2000) Extracellular regulated kinase, but not protein kinase C, is an antiapoptotic signal of insulin-like growth factor-1 on cultured cardiac myocytes. *Biochem. Biophys. Res. Commun.* 273: 736–744
- Kang C Yoo S Hwang B Kim K Kim D Kim C Kim S and Chung B (2000) The inhibition of ERK/MAPK not the activation of JNK/SAPK is primarily required to induce apoptosis in chronic myelogenous leukemic K562 cells. *Leuk. Res.* 24: 527–534
- Jan MS Liu HS and Lin YS (1999) Bad overexpression sensitizes NIH/3T3 cells to undergo apoptosis which involves caspase activation and ERK inactivation. *Biochem. Biophys. Res. Commun.* 264: 724–729
- Nunez G Benedict MA Hu Y and Inohara N (1998) Caspases: the proteases of the apoptotic pathway. *Oncogene* 17: 3237–3245
- Thyrell L Erickson S Zhivotovsky B Pokrovskaja K Sangfelt O Castro J Einhorn S and Grander D (2002) Mechanisms of interferon-alpha induced apoptosis in malignant cells. *Oncogene* 21: 1251–1262
- Caraglia M Tagliaferri P Correale P Genua G Pepe S Pinto A Del Vecchio S Esposito G and Bianco AR (1993) Cytosine arabinoside increases the binding of 125I-labelled epidermal growth factor and 125I-transferrin and enhances the *in vitro* targeting of human tumor cells with anti-(growth factor receptor) mAb. *Cancer Immunol. Immunother.* 37: 150–156

34. Caraglia M Pinto A Correale P Zagonel V Genua G Leardi A Pepe S Bianco AR and Tagliaferri P (1994) 5-Aza-2'-deoxycytidine induces growth inhibition and upregulation of epidermal growth factor receptor on human epithelial cancer cells. *Ann. Oncol.* 5: 269–276
35. Budillon A Di Gennaro E Caraglia M Barbarulo D Abbruzzese A and Tagliaferri P (1999) 8-Cl-cAMP antagonizes mitogen-activated protein kinase activation and cell growth stimulation induced by epidermal growth factor. *Br. J. Cancer* 81: 1134–1141
36. Tagliaferri P Caraglia M Muraro R Budillon A Pinto A and Bianco AR (1994) Pharmacological modulation of peptide growth factor receptor expression on tumor cells as a basis for cancer therapy. *Anti-Cancer Drugs* 5: 379–393
37. Zimmermann S Moelling K (1999) Phosphorylation and regulation of Raf by Akt (protein kinase B). *Science* 286: 1741–1744
38. Kauffmann Zeh A Rodriguez-Viciana P Ulrich E Gilbert C Coffey P Downward J and Evan G (1997) Suppression of c-Myc-induced apoptosis by Ras signalling through PI(3)K and PKB. *Nature* 385: 544–548
39. Serrano M Lin AW Mc Currach ME Beach D and Lowe SW (1997) Oncogenic Ras provokes premature cell senescence associated with accumulation of p53 and p16INK4a. *Cell* 88: 593–602
40. Arase Y Hiwasa T Hasegawa R Nomura J Ito H and Suzuki N (2000) Prevention of v-Ha-Ras-dependent apoptosis by PDGF coordinates in phosphorylation of ERK and Akt. *Biochem. Biophys. Res. Commun.* 267: 33–39
41. von Gise A Lorenz P Wellbrock C Hemmings B Berberich-Siebelt F Rapp UR and Troppmair J (2001) Apoptosis suppression by Raf-1 and MEK1 requires MEK- and phosphatidylinositol 3-kinase-dependent signals. *Mol. Cell. Biol.* 21: 2324–2336
42. Zhou H Li X-M Meinkoth J and Pittman RN (2000) Akt regulates cell survival and apoptosis at a postmitochondrial level. *J. Cell Biol.* 151: 483–494
43. Kuo ML Chuang SE Lin MT and Yang SY (2001) The involvement of PI 3-K/Akt-dependent up-regulation of Mcl-1 in the prevention of apoptosis of Hep3B cells by interleukin-6. *Oncogene* 20: 677–685
44. Liu B Fang M Lu Y Mendelsohn J and Fan Z (2001) Fibroblast growth factor and insulin-like growth factor differentially modulate the apoptosis and G1 arrest induced by anti-epidermal growth factor receptor monoclonal antibody. *Oncogene* 20: 1913–1922
45. Mitsui H Takuwa N Maruyama T Maekawa H Hirayama M Sawatari T Hashimoto N Takuwa Y and Kimura S (2001) The MEK1-ERK map kinase pathway and the PI 3-kinase-Akt pathway independently mediate anti-apoptotic signals in HepG2 liver cancer cells. *Int. J. Cancer* 92: 55–62
46. Alvarez-Tejado M Naranjo-Suarez S Jimenez C Carrera AC Landazuri MO and del Peso L (2001) Hypoxia induces the activation of the phosphatidylinositol 3-kinase/Akt cell survival pathway in PC12 cells: protective role in apoptosis. *J. Biol. Chem.* 276: 22368–22374
47. Chaudhary LR and Hruska KA (2001) The cell survival signal Akt is differentially activated by PDGF-BB, EGF, and FGF-2 in osteoblastic cells. *J. Cell Biochem.* 81: 304–311
48. Tsuneoka M and Mekada E (2000) Ras/MEK signalling suppresses Myc-dependent apoptosis in cells transformed by c-myc and activated Ras. *Oncogene*, 19: 115–123