

**Cover illustration**

Simulated outcome of a proton-proton collision at the LHC, showing the tracks left by particles. (Courtesy of CERN)

Editor, Nature

Philip Campbell

Insights Publisher

Sarah Greaves

Publishing Assistant

Claudia Banks

Insights Editor

Karl Ziemelis

Production Editors

Davina Dudley-Moore

Sarah Archibald

Anna York

Senior Art Editor

Martin Harrison

Art Editor

Nik Spencer

Sponsorship

Emma Green

Production

Susan Gray

Marketing

Katy Dunningham

Elena Woodstock

Editorial Assistant

Alison McGill

THE LARGE HADRON COLLIDER

We are on the threshold of a new era in particle-physics research. In 2008, the Large Hadron Collider (LHC) — the highest-energy accelerator ever built — will come into operation at CERN, the European laboratory that straddles the French-Swiss border near Geneva.

In the debris of the collisions between protons in the LHC, physicists are hoping for a sign. Hypotheses such as the Higgs mechanism (and associated particle) and supersymmetry remain to be proved. The high energy of the LHC collisions will give the best ever 'reach': that is, the best chance yet of finding the decisive signature of a Higgs or supersymmetric particle.

In looking forward, we should not forget what lies behind us — the phenomenal success of the standard model of particle physics over the past three decades. The standard model has been tested to a greater degree of precision than any other model in science and has withstood every challenge. But it is incomplete, and the search for the missing pieces of the puzzle is the prime motivation for building the LHC.

The LHC programme is, however, much wider than a search for the Higgs. Alongside the 'general-purpose' detectors, known as ATLAS and CMS, the LHCb experiment will analyse the production of bottom quarks in LHC collisions. This rich system is the key to a better understanding of the phenomenon of CP violation and its connection to the dominance of matter over antimatter in the Universe. In addition, during dedicated runs in which lead ions will collide in the LHC instead of protons, the ALICE experiment will study a phase of matter called quark-gluon plasma, which might have existed shortly after the Big Bang.

This is the story told in this Insight: how the standard model was developed and tested; how it was agreed to build the LHC; how the programme has been realized through decades of effort by thousands of scientists — and how marvellous the rewards might be.

Alison Wright, Chief Editor, *Nature Physics*

Richard Webb, Senior Editor, *Nature News & Views*

270 Glossary**PERSPECTIVE****271 The making of the standard model**

G. 't Hooft

REVIEW**274 High-energy colliders and the rise of the standard model**

T. Wyatt

PERSPECTIVE**281 How the LHC came to be**

C. Llewellyn Smith

REVIEWS**285 Building a behemoth**

O. Brüning & P. Collier

290 Detector challenges at the LHC

S. Stapnes

297 Beyond the standard model with the LHC

J. Ellis

302 The quest for the quark-gluon plasma

P. Braun-Munzinger & J. Stachel

PERSPECTIVE**310 The God particle *et al.***

L. Lederman

nature
insight