SHORT REPORT

Expanding the phenotype of craniofrontonasal syndrome: two unrelated boys with *EFNB1* mutations and congenital diaphragmatic hernia

Pradeep C Vasudevan^{*,1,4}, Stephen RF Twigg^{2,4}, John B Mulliken³, Jackie A Cook¹, Oliver WJ Quarrell¹ and Andrew OM Wilkie²

¹Department of Clinical Genetics, Sheffield Children's Hospital, Sheffield, South Yorkshire, UK; ²Weatherall Institute of Molecular Medicine, John Radcliffe Hospital, Oxford, UK; ³Craniofacial Centre, Children's Hospital, Boston, MA, USA

Craniofrontonasal syndrome (CFNS, MIM 304110) is an X-linked craniofacial disorder that shows paradoxically greater severity in heterozygous females than in hemizygous males. Mutations have been identified in the *EFNB1* gene that encodes a member of the ephrin-B family of transmembrane ligands for Eph receptor tyrosine kinases. Here, we describe two unrelated families, in both of which a mother and her son have proven mutations in *EFNB1*. The mothers have classical features of CFNS; although the sons have no major craniofacial features other than telecanthus, both had a congenital diaphragmatic hernia (CDH). Our cases represent the first in which CDH has been confirmed in males with mutations in *EFNB1*, highlighting an important role for signalling by ephrin-B1 in the development of the diaphragm.

European Journal of Human Genetics (2006) 14, 884–887. doi:10.1038/sj.ejhg.5201633; published online 26 April 2006

Keywords: craniofrontonasal syndrome; congenital diaphragmatic hernia; *EFNB1*

Introduction

In 1979, craniofrontonasal syndrome (CFNS, MIM 304110) was identified as a subgroup of frontonasal dysplasia by Cohen,¹ who coined the term craniofrontonasal dysplasia. The condition is characterised in females by hypertelorism, coronal craniosynostosis, craniofacial asymmetry, frontal bossing, downslanting palpebral fissures, broad bifid nose, low posterior hairline with an anterior widow's peak, frizzy hair, occasional cleft lip and palate, longitudinally grooved fingernails and other digital anomalies.^{2–7} The fewer males described have mostly been more mildly affected with hypertelorism and, occasionally, cleft lip or palate.^{2–5,7} This greater severity in females than males has always

appeared surprising, given evidence from pedigree analysis that CFNS is an X-linked disorder.^{3,7}

The X-linkage of CFNS was recently confirmed by the identification of mutations in the *EFNB1* gene, which maps to Xq13.1, in both familial and sporadic cases.^{8–11} *EFNB1* encodes ephrin-B1, a transmembrane ligand for Eph receptor tyrosine kinases that itself exhibits receptor-like effects. Significantly, mice with null mutations of ephrin-B1 have malformations of the axial and appendicular skeleton that also show greater severity in heterozygous females than in either homozygous females or hemizygous males.^{12,13} Because of X-inactivation, heterozygous females are uniquely mosaic for cells that either do, or do not, make functional protein; this is proposed to lead to abnormal sorting of cell populations and disturbance of normal tissue boundary formation. The term cellular interference has been suggested for this process.⁹ This ectopic signalling mechanism of pathogenesis makes it difficult to draw firm conclusions about the normal functions of ephrin-B1 from the phenotype of CFNS females. By contrast, the usually mild male phenotype

*Correspondence: Dr PC Vasudevan, Department of Clinical Genetics, Sheffield Children's Hospital, Western Bank, Sheffield, South Yorkshire, S10 2 TH, UK.

Tel: +44 114 2717025 or +44 114 2260570; Fax: +44 114 2737467; E-mail: pradeepcv@yahoo.com

⁴These authors contributed equally to this work

Received 5 January 2006; revised 1 March 2006; accepted 1 March 2006; published online 26 April 2006

associated with ephrin-B1 mutations in both humans and mice has been attributed to functional redundancy with other ephrins. Here we report a more severe phenotype, congenital diaphragmatic hernia (CDH), in two unrelated males hemizygous for *EFNB1* mutations. This suggests a nonredundant role for ephrin-B1 in the development of the diaphragm.

Case reports and mutation analysis

Family 1

A 4-year-old boy (case 1) born to nonconsanguineous Caucasian parents was found antenatally to have a left-sided CDH and sacroccygeal teratoma. He was born by spontaneous normal vaginal delivery at 34 weeks' gestation. He had a primary repair of his diaphragmatic defect (Figure 1a) on day 2 with good result. His sacroccygeal teratoma was excised on day 21. On examination at 6 months of age, he had a prominent metopic suture, telecanthus (inner canthal distance of 28 mm; +2 s.d. above the mean), brachydactyly and pectus excavatum. Later he grew curly blonde hair (Figure 1b). There was no significant developmental delay. His head circumference and weight were between the 0.4th and 2nd centile and height between the 2nd and 9th centile. Investigations revealed a normal male karyotype (46, XY). His cranial ultrasound scan and skull X-ray were normal. A clinical diagnosis of male CFNS was made on the basis of his mild dysmorphic features and characteristic clinical features in the mother (Figure 1c and d). Mutation analysis of *EFNB1* using previously described methods⁸ demonstrated a frameshifting mutation (432delG, predicting L145WfsX14) in exon 3, which was heterozygous in the mother and hemizygous in her son (Figure 1e and f). While sacroccygeal teratoma is not a recognised association of CFNS, we note that a similarly affected female CFNS patient with a confirmed *EFNB1* mutation was described previously.⁸

Family 2

A male infant (case 2) of nonconsanguineous Caucasian parents was born at 37 weeks' gestation with left-sided CDH, talipes equinovarus and swallowing difficulties. The CDH was repaired at age 3 days with a Gore-Tex patch, fundoplication and gastrostomy tube placement, with a good result. On examination at 2 years of age, he had telecanthus (inner canthal distance of 33 mm; more than +2 s.d. above the mean) but no other dysmorphic features (Figure 2a). His mother (Figure 2b) and her two half sisters (Figure 2c and d), each of whom had different fathers, all had characteristic clinical features of CFNS. A clinical diagnosis of male CFNS was made in the proband on the basis of the hypertelorism and family history. Analysis of *EFNB1* demonstrated a 3-nucleotide deletion in exon 2 (151_153delGTG, predict-

Figure 1 Clinical features and mutation analysis of case 1 and mother. (a) Plain radiograph of case 1 aged 3 h. Note large left diaphragmatic hernia with stomach bubble in chest and gross mediastinal shift to right. (b) Case 1 aged 3 years. Note hypertelorism, prominent metopic ridge and curly hair. (c and d) Mother showing craniofacial asymmetry, telecanthus, broad bifid nose and digital anomalies with longitudinal ridging in nails. (e) Forward sequence of *EFNB1* exon 3 from case 1 and control DNA. The nucleotide deleted in case 1 (432delG) is shaded in the control sequence. (f) Confirmation of 432delG mutation. *EFNB1* exon 3 was amplified⁸ from case 1, his mother and control DNA and incubated with *Bgl*I, which specifically digests the 432delG allele (normal fragment 201 bp; mutant, 96 and 104 bp). Case 1 is hemizygous for the mutation, his mother is heterozygous and the control is negative. (–) undigested, (+) digested with *Bgl*I.

ing V51del), present in heterozygous state in the mother and one of her sisters (a sample from the other sister was unavailable), and in hemizygous state in the son (Figure 2e and f).

Figure 2 Clinical features and mutation analysis of case 2 and family. (a) Case 2 showing hypertelorism. (b) Mother of case 2 exhibiting craniofacial asymmetry and sloping shoulders. (c and d) Maternal aunts of case 2. (e) Reverse sequence of *EFNB1* exon 2 from case 2 and control DNA. Three nucleotides (shown shaded in control) are deleted in case 2 (151_153delGTG). (f) Confirmation of 151_153delGTG mutation. *EFNB1* exon 2 was amplified⁸ and digested with *HphI*, a site for which is abolished by the mutation (normal allele 75, 125 and 167 bp; mutant allele, 197 and 167 bp). The expected normal allele products are present in the control; the mother and aunt illustrated in panel c show the heterozygous pattern; case 2 is hemizygous for the mutation. PCR product not digested with *HphI* (uncut) is 367 bp.

Discussion

The identification of two unrelated males hemizygous for *EFNB1* mutations, both of whom have CDH, provides new insight into the pathogenesis of this disorder. CDH occurs about once in 3000 newborns and is associated with a 30–60% mortality rate with significant morbidity among survivors.¹⁴ CDH has been described in association with many chromosome abnormalities and in a number of syndromes;^{15–17} however, the only known genetic defect associated with nonsyndromic congenital diaphragmatic defect in humans is mutation in the *FOG2* gene.¹⁸ Studies in rodents have shown three general causes of developmental error leading to CDH. Failure of the normal closure of the pleuroperitoneal folds from embryonic day 12.5 in rats (equivalent to week 5 in humans) causes the common posterolateral defect seen in the cases described here. Alternatively, defects in the formation of the central tendon, or in migration of the muscle and nerve precursor cells from the cervical somites to the diaphragm, can also result in CDH.¹⁹

CDH was previously described in four females with CFNS,^{8,20–22} but the developmental interpretation of these cases is complicated by the presence of X-inactivation (see Introduction). The only previous report of CDH in affected males was by Morris *et al*,²³ who described a family of six individuals with apparent CFNS including two affected brothers with CDH. However, several authors have disputed this diagnosis as only one family member had craniosynostosis and the affected males had additional manifestations not described in CFNS, including short stature, delayed bone age, shawl scrotum and hypospadias; an alternative diagnosis of brachycephalofrontal dysplasia (Teebi syndrome; MIM 145420) has been proposed.^{5,7,22} To our knowledge, no molecular analysis has been undertaken in this family, so the issue remains unresolved. The two male CDH cases presented here represent the first in which the diagnosis has been confirmed by the finding of pathogenic mutations in *EFNB1*. Both mutations are likely to lead to the loss of function. In family 1, the frameshift mutation truncates the C-terminal portion of the extracellular domain and, in addition, may lead to nonsense-mediated decay of the mutant transcript; in family 2, the deleted valine is within a β -sheet²⁴ and therefore normal folding of the protein is likely to be disrupted. The spectrum of *EFNB1* mutations observed in males without CDH^{9–11} is qualitatively similar, arguing against a distinct pathogenic action of the mutations observed here: presumably, the occurrence of CDH in our cases was caused by additional, unknown modifier effects.

In conclusion, our data indicate that the male CFNS phenotype is not always mild and that loss of ephrin-B1 function may interfere with normal diaphragmatic development. These observations need to be incorporated into genetic counselling for families with CFNS. Previous studies of ephrin-B1 expression in mouse²⁵ and chick²⁶

do not indicate which aspects of diaphragmatic development might be affected, and diaphragmatic defects were not reported in either of the published murine *Efnb1* mutants.^{14,15} Our observations provide impetus to further experimental studies of ephrin-B1 signalling in the development of the diaphragm.

Acknowledgements

We are very grateful to the families for their participation in this work. The mutation analysis described in this work was funded by the Wellcome Trust (AOMW).

References

- Cohen Jr MM: Craniofrontonasal dysplasia. *Birth Defects Orig Art Ser* 1979; **15**: 85–89.
- Slover R, Sujansky E: Frontonasal dysplasia with coronal craniosynostosis in three sibs. *Birth Defects Orig Art Ser* 1979; **15**: 75–83.
- Grutzner E, Gorlin RJ: Craniofrontonasal dysplasia: phenotypic expression in females and males and genetic considerations. *Oral Surg Oral Med Oral Pathol* 1988; **65**: 436–444.
- Kere J, Ritvanen A, Marttinen E, Kaitila I: Craniofrontonasal dysostosis: variable expression in a three-generation family. *Clin Genet* 1990; **38**: 441–446.
- Saavedra D, Richieri-Costa A, Guion-Almeida ML, Cohen Jr MM: Craniofrontonasal syndrome: study of 41 patients. *Am J Med Genet* 1996; **61**: 147–151.
- Orr DJA, Slaney S, Ashworth GJ, Poole MD: Craniofrontonasal dysplasia. *Br J Plast Surg* 1997; **50**: 153–161.
- Cohen Jr MM: Syndromes with craniosynostosis—craniofrontonasal syndrome; in Gorlin RJ, Cohen Jr MM, Hennekam RCM (ed): *Syndromes of the Head and Neck*. Oxford: Oxford University Press, 2001, pp 981–983.
- Twigg SRF, Kan R, Babbs C *et al*: Mutations of ephrin-B1 (*EFNB1*), a marker of tissue boundary formation, cause craniofrontonasal syndrome. *Proc Natl Acad Sci USA* 2004; **101**: 8652–8657.
- Wieland I, Jakubiczka S, Muschke P *et al*: Mutations of the ephrin-B1 gene cause craniofrontonasal syndrome. *Am J Hum Genet* 2004; **74**: 1209–1215.
- Wieland I, Reardon W, Jakubiczka S *et al*: Twenty-six novel *EFNB1* mutations in familial and sporadic craniofrontonasal syndrome (CFNS). *Hum Mutat* 2005; **26**: 113–118.
- Twigg SRF, Matsumoto K, Kidd AMJ *et al*: The origin of *EFNB1* mutations in craniofrontonasal syndrome: frequent somatic mosaicism and explanation of the paucity of carrier males. *Am J Hum Genet* 2006 (in press).
- Compagni A, Logan M, Klein R, Adams R: Control of skeletal patterning by ephrinB1-EphB interactions. *Dev Cell* 2003; **5**: 217–230.
- Davy A, Aubin J, Soriano P: Ephrin-B1 forward and reverse signalling are required during mouse development. *Genes Dev* 2004; **18**: 572–583.
- Harrison MR, Adzick NS, Estes JM, Howell LJ: A prospective study of the outcome for fetuses with diaphragmatic hernia. *JAMA* 1994; **271**: 382–384.
- Witters I, Legius E, Moerman P *et al*: Associated malformations and chromosomal anomalies in 42 cases of prenatally diagnosed diaphragmatic hernia. *Am J Med Genet* 2001; **103**: 278–282.
- Lurie IW: Where to look for the genes related to diaphragmatic hernia? *Genet Couns* 2003; **14**: 75–93.
- Slavotinek AM: The genetics of congenital diaphragmatic hernia. *Semin Perinatol* 2005; **29**: 77–85.
- Ackerman KG, Herron BJ, Vargas SO *et al*: *Fog2* is required for normal diaphragm and lung development in mice and humans. *PLoS Genet* 2005; **1**: 58–65.
- Babiuk RP, Zhang W, Clugston R *et al*: Embryological origins and development of the rat diaphragm. *J Comp Neurol* 2003; **455**: 477–487.
- Hurst J, Baraitser M: Craniofrontonasal dysplasia. *J Med Genet* 1988; **25**: 133–138.
- Brooks AS, van Dooren M, Hoozeboom J *et al*: Congenital diaphragmatic hernia in a female patient with craniofrontonasal syndrome. *Clin Dysmorphol* 2002; **11**: 151–153.
- McGaughan J, Rees M, Battin M: Craniofrontonasal syndrome and diaphragmatic hernia. *Am J Med Genet* 2002; **110**: 391–392.
- Morris CA, Palumbos JC, Carey JC: Delineation of the male phenotype in craniofrontonasal syndrome. *Am J Med Genet* 1987; **27**: 623–631.
- Nikolov DB, Li C, Barton WA, Himanen JP: Crystal structure of the ephrin-B1 ectodomain: implications for receptor recognition and signaling. *Biochemistry* 2005; **44**: 10947–10953.
- Flenniken AM, Gale NW, Yancopoulos GD, Wilkinson DG: Distinct and overlapping expression patterns of ligands for Eph-related receptor tyrosine kinases during mouse embryogenesis. *Dev Biol* 1996; **179**: 382–401.
- Baker RK, Antin PB: Ephs and ephrins during early stages of chick embryogenesis. *Dev Dyn* 2003; **228**: 128–142.