

SHORT REPORT

A locus for sacral/anorectal malformations maps to 6q25.3 in a 0.3 Mb interval region

Luigi Titomanlio^{1,2}, Irina Giurgea³, Clarisse Baumann², Monique Elmaleh⁴, Philippe Sachs¹, François Chalard⁴, Azzedine Aboura⁵ and Alain Verloes^{*,2,6}

¹Department of Child Neurology, AP-HP Robert Debré Hospital, Paris, France; ²Clinical Genetics Unit, AP-HP Robert Debré Hospital, Paris, France; ³Department of Genetics and Biochemistry, INSERM U654, AP-HP Henri Mondor Hospital, Créteil, France; ⁴Medical Imaging Department, AP-HP Robert Debré Hospital, Paris, France; ⁵Cytogenetics Unit, AP-HP Robert Debré Hospital, Paris, France; ⁶INSERM U676, AP-HP Robert Debré Hospital, Paris, France

Partial absence of the sacrum is a rare congenital defect that also occurs as an autosomal-dominant trait, whereas imperforate/ectopic anus is a relatively common malformation, usually observed in multiple congenital anomalies syndromes. We report on a girl born to healthy consanguineous parents (first cousins once removed) with anal imperforation and associated rectovaginal fistula and partial sacral agenesis. Facial dysmorphism included a high forehead, epicanthic folds, downslanting palpebral fissures, hypertelorism and a depressed nasal root. Brain MRI showed a bilateral opercular dysplasia with a unilateral (right) pachygyria; MRI and X-ray imaging of the spine disclosed a tethered cord associated with partial sacral agenesis. She showed a moderate developmental delay. Ophthalmologic examination evidenced bilateral microphthalmos and relative microcornea. Cytogenetic studies in our patient disclosed a pure *de novo* 6q25.3 → qter deletion. By genotype analysis, we detected in our patient a maternal allele loss encompassing D6S363 and D6S446. Pure distal 6q deletion is a rare anomaly, reported in association with sacral/anorectal malformations (sacral agenesis, anal imperforation/ectopia) and never with cortical dysplasia. Pooling deletion mapping information in patients with pure terminal and interstitial 6q deletion allowed us to define a critical region spanning 0.3 Mb between the markers D6S959 and D6S437 for sacral/anal malformations. We hypothesize that haploinsufficiency for a gene within the deleted region may impair normal development of caudal structures, possibly acting on the notochordal development.

European Journal of Human Genetics (2006) 14, 971–974. doi:10.1038/sj.ejhg.5201635; published online 17 May 2006

Keywords: 6q deletion; sacrum agenesis; pachygyria; anal atresia; genotype/phenotype correlation; opercular dysplasia

Introduction

Terminal deletions of 6q (6q25 → 6qter) have been rarely reported in the literature and have been associated to a specific phenotype. Findings in a group of 26 patients¹ included mental retardation (100%), ear anomalies (88%),

hypotonia (86%), microcephaly (82%), limb anomalies (71%), brain anomalies (67%), eye anomalies (50%), cardiac defects (48%), genital anomalies (48%) and seizures (38%). However, only in the patient by McLeod *et al.*,² a pure deletion was found. To date, subtelomeric deletion of 6q have been reported in three patients,^{3–5} and patients carrying a pure terminal or interstitial deletion of 6q have been rarely observed.^{6–12}

Partial absence of the sacrum is a rare congenital defect, which also occurs as an autosomal-dominant trait; association with anterior meningocele, presacral teratoma and

*Correspondence: Professor A Verloes, Clinical Genetics Unit, AP-HP Robert Debré Hospital, 48 bd Sérurier, Paris 75019, France.

Tel: +33 1 4003 5342; Fax: +33 1 4003 5344;

E-mail: alain.verloes@rdb.aphp.fr

Received 26 September 2005; revised 8 February 2006; accepted 23 February 2006; published online 17 May 2006




Figure 1 Clinical and neuroradiological aspects. (a) Facial appearance at age 3. (b) MRI: coronal section (T2), note normal gyral pattern. (c) MRI: frontal section (T1), bilateral opercular dysplasia and unilateral (right) pachygyria. (d) Sagittal MRI section of dorso-lumbar spinal canal disclosing tethered spinal cord.

anorectal abnormalities constitutes the Currarino triad (MIM 176450).

Imperforate/ectopic anus is a relatively common malformation. It has been rarely reported in familial cases (MIM 207500, MIM 301800), but no mapping data are available.

Patient report

The girl was born at 35 weeks of gestation to healthy consanguineous Malagasi parents (first cousins once removed). Family history and gestation were unremarkable. Birth weight was 2500 g (10th–25th centile), length was 48 cm (25th–50th centile) and cranial circumference was 32 cm (5th–10th centile). She was immediately admitted to a tertiary care center for surgical correction of an anal imperforation with an associated rectovaginal fistula. Imaging studies showed partial sacral agenesis and a tethered cord. Echocardiographic evaluation revealed atrial and ventricular septal defects that were corrected at the age of 12 months. Clinical evaluation at 24 months showed a severe developmental delay, microcephaly, short stature and minor anomalies. Weight was 10 kg (–1.3 SD), height was 80 cm (–1.4 SD) and head circumference was 44 cm (–2.8 SD). Facial dysmorphism included a high forehead, epicanthal folds, downslanting palpebral fissures, hypertelorism and a depressed nasal root (Figure 1a). Brain MRI showed a bilateral opercular dysplasia with a unilateral (right) pachygyria (Figure 1b and c); MRI and X-ray

imaging of the spine disclosed a tethered cord associated with a partial sacral agenesis (Figure 1d).

At 3 years, 2/12 of age ophthalmologic examination evidenced bilateral microphthalmos (diameter of the globe <20 mm) and relative microcornea. Visual-evoked potentials and electroretinogram were normal.

Methods

Cytogenetic analysis

Subtelomeric FISH probing was performed with Cytocell Chromoprobe Multiprobe kit.

Genotype analysis

DNA from leukocytes of patient and their parents were used for genotyping. Simple fluorescent PCR assays were performed using polymorphic markers: D6S292, D6S308, D6S441, D6S1577, D6S415, D6S959, D6S363, D6S437, D6S1614, D6S1581, D6S264 and D6S446 (<http://www.gdb.org/>). PCR reactions were performed following standard procedures. After denaturation, each sample was loaded for electrophoresis on an Applied Biosystems model 3100 automated sequencer (PE Applied Biosystems, Perkin-Elmer). Data were analyzed using the Gene Scanner Model 3.7 Fluorescent Fragment Analyzer (PE Applied Biosystems, Perkin-Elmer) and electropherograms were generated for each sample. The samples from the patient and from his parents were processed for each marker. Data were analyzed using the Gene Scanner Model 3.7 Fluorescent Fragment Analyzer (PE Applied Biosystems, Perkin-Elmer).

Results

Metaphase cells analyzed from cultures of peripheral blood on the patient revealed a normal female chromosome complement at the 650-band level. A terminal 6q deletion was found by subtelomeric FISH. The patient's karyotype was designated as 46, XX, del (6)(q25.3qter). In both parents, FISH using identical probe for the subtelomeric region of 6p and 6q yielded normal result. There was no evidence for a balanced rearrangement in the parents. By genotype analysis, we detected in our patient a maternal allele loss for the D6S363, D6S1581, D6S264 and D6S446 markers (Figure 2).

Discussion

Present patient shows a multiple congenital anomalies syndrome owing to a pure *de novo* 6q23qter deletion. A peculiar finding is the presence of a bilateral opercular dysplasia associated to a unilateral pachygyria at the brain MRI. These brain malformations are probably consequent to the chromosomal deletion, although they have never been reported to date in other patients who carry an isolate interstitial or terminal 6q25 deletion. Moreover, no locus for brain malformation has been mapped in this region, with the exception of corpus callosum agenesis (6q25). Because of the presence of parental consanguinity, a recessive phenotype compounded by a homozygote mutation is also possible. A long-range effect of the deletion on gene expression outwith the deletion is another possibility to be considered.

On the other hand, distal 6q deletions have been frequently reported in association with sacral/anorectal malformations (sacral agenesis, anal imperforation/ectopia).

Our patient showed anal imperforation with associated rectovaginal fistula and partial sacral agenesis. Probably, a common pathogenetic mechanism is involved in producing both malformations. At an early stage of development, the notochord is known to organize normal development of central axial structures, such as the spinal cord, vertebral column and anorectum. However, its role has not been completely elucidated.¹³ Recently, Qi *et al*¹⁴ suggested that an alteration in sonic hedgehog signaling may be pivotal in producing abnormal notochord development and consequently sacral/anorectal malformations.

By comparing the reported cases of affected patients carrying a pure deletion, we delimited a critical region of 0.3 Mb for sacral/anorectal malformations, lying between the markers D6S959 and D6S437 (see Table 1). The patient by Pirola *et al*¹⁰ shows an ectopic anus, which can be considered as a mild form of anorectal malformation. Our patient's contribution to the definition of a critical region should be cautiously considered, because of the simultaneous presence of another midline defect (agenesis of the corpus callosum).

We hypothesize that there is a gene in the deleted region whose haploinsufficiency impairs the normal development of these structures, possibly acting on the notochordal development or interfering with SHH signaling.




Figure 2 (a) Microsatellite analysis of the deletion. Dotted arrow is used for noninformative markers. (b) Comparative deletion mapping in patients with 6q deletion and anorectal malformations. The overlapping segment between Pirola's case and our patient is depicted in the bottom line, with the four known genes.

Table 1 Clinical findings in selected cases with pure interstitial or terminal 6q deletions

| | <i>Present case</i> | <i>Evers et al</i> ⁸ | <i>McLeod et al</i> , ² case 1 | <i>Meng et al</i> , ⁷ case 1 | <i>Pirola et al</i> ¹⁰ | <i>Lukusa et al</i> ¹² |
|-------------------------------|--|---------------------------------|---|---|-----------------------------------|--|
| Deletion | q25.3qter | q25.1qter | q23q25 | q24.3qter | q25.1q25.3 | q25.3qter |
| Sex | F | M | M | M | F | M |
| Age at report | 3y 2m | 2y | 1m | 4m | 9m | 10y |
| Microcephaly | + | + | – | + | – | – |
| Hypotonia | + | – | – | + | + | + |
| Seizures | – | – | – | – | – | – |
| Developmental delay | + | + | NM | NM | + | + |
| Brain neuroimaging | Bilateral opercular dysplasia pachygyria | Hydrocephalus | NM | Agenesis of corpus callosum | Agenesis of corpus callosum | Hydrocephalus |
| Retinal abnormalities | – | NM | – | – | – | + |
| Cardiac malformation | + | + | – | + | – | – |
| Sacral/anorectal malformation | Imperforate anus partial sacral agenesis | Bony appendix of the coccyx | Imperforate anus sacral agenesis | Imperforate anus | Ectopic anus | Spina bifida bony appendix of the coccyx |

+, characteristic present; –, characteristic absent; NM, not mentioned.

To date, four genes are positioned in the deleted region: *SYNJ2* (synaptojanin 2), *SERAC1* (serine-active site containing 1), *GTF2H5* (general transcription factor iih, polypeptide 5) and *TULP4* (Tubby-like protein 4) (UCSC Genome Browser, <http://genome.ucsc.edu/> and Ensembl Genome Browser, <http://www.ensembl.org>). *GTF2H5* mutations are responsible for trichothiodystrophy group A, a DNA repair syndrome, and for a form of ichthyosiform erythroderma with hair abnormality, and mental and growth retardation.¹⁵ Sacral/anorectal malformations have not been observed in the reported patients. The Synj2b protein isoforms are localized in nerve terminals in rat brain and at spermatid manchette in rat testis. In glioblastoma cell lines, Synj2b seems implicated in the regulation of the formation of invadopodia and lamellipodia.¹⁶ Mutations in *SERAC1* or *SYNJ2* cause male mouse sterility.¹⁷ *TULP4* is a putative transcription factor of unknown function. No role has been attributed for the latter three genes in human developmental anomalies and/or diseases.

Other genes mapping outside the reported critical region could be involved, because of possible modifications on gene expression.

Report of further patients is needed to evaluate these genes as candidates in sacral/anorectal malformations, and their hypothetical role in notochordal development.

References

- Hopkin RJ, Schorry E, Bofinger M *et al*: New insights into the phenotypes of 6q deletions. *Am J Med Genet* 1997; **70**: 377–386.
- McLeod DR, Fowlow SB, Robertson A *et al*: Chromosome 6q deletions: a report of two additional cases and a review of the literature. *Am J Med Genet* 1990; **35**: 79–84.
- Lorda-Sanchez I, Lopez-Pajares I, Roche MC *et al*: Cryptic 6q subtelomeric deletion associated with a paracentric inversion in a mildly retarded child. *Am J Med Genet* 2000; **95**: 336–338.
- Anderlid BM, Schoumans J, Anneren G *et al*: FISH-mapping of a 100-kb terminal 22q13 deletion. *Hum Genet* 2002; **110**: 439–443.
- Stevenson DA, Brothman AR, Carey JC *et al*: 6q subtelomeric deletion: is there a recognizable syndrome? *Clin Dysmorphol* 2004; **13**: 103–106.
- Narahara K, Tsuji K, Yokoyama Y *et al*: Specification of small distal 6q deletions in two patients by gene dosage and *in situ* hybridization study of plasminogen and alpha-L-fucosidase 2. *Am J Med Genet* 1991; **40**: 348–353.
- Meng J, Fujita H, Nagahara N *et al*: Two patients with chromosome 6q terminal deletions with breakpoints at q24.3 and q25.3. *Am J Med Genet* 1992; **43**: 747–750.
- Evers LJ, Schrandt-Stumpel CT, Engelen JJ *et al*: Deletion of the long arm of chromosome 6: two new patients and literature review. *Clin Genet* 1996; **50**: 138–144.
- Rubtsov N, Senger G, Kuzcera H *et al*: Interstitial deletion of chromosome 6q: precise definition of the breakpoints by microdissection, DNA amplification, and reverse painting. *Hum Genet* 1996; **97**: 705–709.
- Pirola B, Bortotto L, Giglio S *et al*: Agenesis of the corpus callosum with Probst bundles owing to haploinsufficiency for a gene in an 8 cM region of 6q25. *J Med Genet* 1998; **35**: 1031–1033.
- Sukumar S, Wang S, Hoang K *et al*: Subtle overlapping deletions in the terminal region of chromosome 6q24.2–q26: three cases studied using FISH. *Am J Med Genet* 1999; **87**: 17–22.
- Lukusa T, Willekens D, Lukusa N *et al*: Terminal 6q25.3 deletion and abnormal behaviour. *Genet Counsel* 2001; **12**: 213–221.
- Stemple DL: Structure and function of the notochord: an essential organ for chordate development. *Development* 2005; **132**: 2503–2512.
- Qi BQ, Beasley SW, Frizelle FA: Evidence that the notochord may be pivotal in the development of sacral and anorectal malformations. *J Pediatr Surg* 2003; **38**: 1310–1316.
- Giglia-Mari G, Coin F, Ranish JA *et al*: A new, tenth subunit of TFIIF is responsible for the DNA repair syndrome trichothiodystrophy group A. *Nat Genet* 2004; **36**: 714–719.
- Chuang YY, Tran NL, Rusk N *et al*: Role of synaptojanin 2 in glioma cell migration and invasion. *Cancer Res* 2004; **64**: 8271–8275.
- Schimenti JC, Reynolds JL, Planchart A: Mutations in Serac1 or Synj2 cause proximal t haplotype-mediated male mouse sterility but not transmission ratio distortion. *Proc Natl Acad Sci USA* 2005; **102**: 3342–3347.