Association of early life factors and acute lymphoblastic leukaemia in childhood: historical cohort study

L Murray^{*,1}, P McCarron^{1,2}, K Bailie³, R Middleton¹, G Davey Smith⁴, S Dempsey⁵, A McCarthy⁵ and A Gavin¹

¹Northern Ireland Cancer Registry, Department of Epidemiology and Public Health, The Queens University, Belfast, Riddel Hall, Stranmillis Road, Belfast BT9 5EE, UK; ²Division of Cancer Control and Population Sciences, National Cancer Institute, Rockville, Maryland, USA; ³Ulster Community & Hospitals Trust, Belfast, UK; ⁴Department of Social Medicine, University of Bristol, Bristol, UK; ⁵Royal Group of Hospitals Trust, Belfast, UK

In a historical cohort study of all singleton live births in Northern Ireland from 1971–86 ($n=434\,933$) associations between early life factors and childhood acute lymphoblastic leukaemia were investigated. Multivariable analyses showed a positive association between high paternal age (≥ 35 years) and acute lymphoblastic leukaemia (relative risk=1.49; 95% confidence interval (CI)=0.96–2.31) but no association with maternal age. High birth weight (≥ 3500 g) was positively associated with acute lymphoblastic leukaemia (relative risk=1.66; 95% CI=1.18–2.33). Children of mothers with a previous miscarriage or increased gestation (≥ 40 weeks) had reduced risks of ALL (respective relative risks=0.49; 95% CI=0.29–0.80, and 0.67; 95% CI=0.48–0.94). Children born into more crowded households (≥ 1 person per room) had substantially lower risks than children born into less crowded homes with also some evidence of a lower risk for children born into homes with three adults (relative risks=0.56; 95% CI=0.35–0.91 and 0.58; 95% CI=0.21–1.61 respectively). These findings indicate that several early life factors, including living conditions in childhood and maternal miscarriage history, influence risk of acute lymphoblastic leukaemia in childhood.

British Journal of Cancer (2002) **86**, 356–361. DOI: 10.1038/sj/bjc/6600012 www.bjcancer.com
© 2002 The Cancer Research Campaign

Keywords: ALL; childhood; early life factors; historical cohort

The aetiology of childhood leukaemia, the most common childhood malignancy, is poorly understood and there are few established risk factors. Recently there has been growing interest in the relationship between prenatal and early life influences on risk of childhood leukaemia (Van Steensel-Moll *et al*, 1985; Shu *et al*, 1988; Kaye *et al*, 1991; Westergaard *et al*, 1997; McKinney *et al*, 1999). It is clear that mutations occurring in-utero in lymphoid stem cells contribute to the development of childhood leukaemia (Greaves, 1999; Wiemels *et al*, 1999) but it is not known which prenatal exposures result in these mutations or how they may be prevented. The apparent association between high birth weight and increased risk of acute lymphoblastic leukaemia (ALL) has stimulated research into the role of insulin growth factor-1 (IGF-1) and relevant receptors in abnormal haematopoiesis. Other possible prenatal or early life influences, as indexed, for example, by birth order or social class at birth have less well understood potential mechanisms. Evidence from studies on population mixing (Kinlen, 1995) indicate a role for infectious agents and delayed exposure to infection has been suggested as influencing risk of childhood leukaemia (Greaves and Alexander, 1993; Greaves, 1997).

There is a lack of consistency among reports examining prenatal and early life associations with childhood leukaemia, suggesting that some of the observed associations are spurious. Study designs utilized to date may have contributed to this lack of consistency and very few population-based cohort studies of childhood leukaemia

have been performed (Westergaard *et al*, 1997; Mogren *et al*, 1999). We describe a population-based cohort study, undertaken in Northern Ireland, which includes unique information recorded at birth on socio-economic position and living conditions in the household.

MATERIALS AND METHODS

Since 1971, the Northern Ireland Child Health System has collected information on all births to mothers normally resident in Northern Ireland. The Department of Health, Social Services and Public Safety maintains the data relating to births between 1971 and 1986. Subject to an appropriate confidentiality agreement, access was granted to these data which include: date and place of birth; birth weight (in grams); gestational age (to the nearest complete week); number of previous pregnancies, live births and miscarriages; mother's and father's year of birth; social class (based on father's occupation); method of infant feeding on discharge from place of confinement; number of persons aged above and below 15 years of age per household and; number of living rooms and bedrooms in the household at the time of the birth.

Details of incident cases of leukaemia which occurred before the age of 16 years in Northern Ireland between 1975 and 1997 were obtained from the Northern Ireland Cancer Registry's Register of Childhood Cancers. Data on cases are obtained from multiple sources, including: the Northern Ireland Leukaemia and Lymphoma Register; the Oxford Childhood Cancer Register; clinicians' records and; databases of specific research projects into childhood cancer. Using sex, surname, and date of birth, cases of ALL born in Northern Ireland between 1971 and 1986 were

*Correspondence: Dr L Murray; E-mail: l.murray@qub.ac.uk

Received 14 August 2001; revised 2 October 2001; accepted 15 October 2001

identified within the Child Health System database. Exact matches, and matches where one character was incorrect were accepted after manual checking. Perinatal information relating to the children who subsequently developed ALL was compared to that of children who did not develop ALL. Multiple births were excluded from the analyses. There were too few cases of acute myeloid leukaemia ($n=27$) to permit investigation in the same manner.

Univariable analyses

Approximate maternal and paternal age, at time of the child's birth, were calculated by subtracting mother's and father's year of birth from year of birth of the child. Household density was calculated as the number of residents of all ages (including the newborn) divided by the number of rooms in the accommodation. These variables and number of previous miscarriages, birth order, diagnosis of Down's syndrome at birth, method of feeding, and number of children and adults in the household were categorized as in Table 1. Social class, based on father's occupation, was coded in the Child Health System according to the Office of Population Censuses and Surveys classification of occupations into six groups as categorized in Table 2. In analyses this was dichotomized into non-manual (groups 1–3) and manual/other (groups 4–6) occupations. Differences in proportions in these predictor variables were compared between those children who did and those who did not develop ALL using the χ^2 test and the χ^2 test for trend. Where appropriate, variables were dichotomized and χ^2 tests repeated. Parental age was dichotomized into <35 years and ≥ 35 years, and gestational age into <40 weeks and ≥ 40 weeks.

Multivariable analyses

As we did not actively follow up the cohort for migration or death we did not feel justified in calculating person years and modelling the risk of disease using Poisson or Cox regression. Instead we approximated relative risks by the odds ratios obtained from logistic regression models, an approximation which is reasonable for a rare disease such as childhood ALL. Variables with a $P < 0.2$ level in univariable analyses were included in multivariable models. As data relating to living conditions were unavailable before 1976 or after 1984, two multivariable models were constructed: model I did not include number of adults in the household or household density; model II included these variables. These models were also adjusted for sex and Down's syndrome and models were also constructed to include only those children with a diagnosis of ALL before the age of 6 years. All statistical analyses were performed using SPSS for Windows, Version 10.0 (SPSS Inc., Chicago, USA). Ethical approval was obtained from the Research Ethics Committee of The Queen's University of Belfast.

RESULTS

The Child Health System contained data on 444 168 live births (including 9235 multiple births) which occurred between January 1971 and December 1986 to mothers normally resident in Northern Ireland. The Registrar General's Office, which registers all births within Northern Ireland, registered 444 111 live births for the same period. The Child Health System may include a small number of duplicate records. The Northern Ireland Cancer Registry contains 342 cases of ALL diagnosed in children aged less than 16

Table 1 Association between selected early life factors and birth characteristics and risk of childhood ALL, univariable analyses

Variable	Categories	Cases N (%)	Non-cases N (%)	Relative risk (95% CI)	P value (trend)
Maternal age (years)	<25	59 (31.4)	146 527 (33.8)	0.95 (0.69, 1.30)	0.33 (0.21)
	25–34	99 (52.7)	232 722 (53.7)	1.00 ^b	
	≥ 35	30 (16.0)	54 069 (12.5)	1.30 (0.87, 1.96)	
Paternal age (years)	<25	27 (16.2)	77 793 (20.0)	0.86 (0.60, 1.30)	0.06 (0.03)
	25–34	92 (55.1)	227 073 (58.5)	1.00 ^b	
	≥ 35	48 (28.7)	83 552 (21.5)	1.42 (1.00, 2.00)	
Miscarriage history	None	146 (86.9)	318 909 (80.5)	1.00 ^b	0.04
	1 or more	22 (13.1)	77 012 (19.5)	0.62 (0.40, 0.98)	
Gestational age	<38 weeks	19 (10.2)	36 297 (8.4)	0.96 (0.58, 1.59)	0.17 (0.45)
	38–39 weeks	98 (52.4)	255 229 (59.2)	1.00 ^b	
	≥ 40 weeks	70 (37.4)	139 679 (32.4)	0.73 (0.45, 1.20)	
Birth order	First-born	52 (30.2)	119 828 (29.8)	1.00 ^b	0.87
	Not first born	120 (69.8)	282 605 (70.2)	0.98 (0.71, 1.36)	
Down's syndrome	Yes	6 (3.2)	434 (0.1)	33.0 (14.5, 74.7)	<0.001
	No	182 (96.8)	434 311 (99.9)	1.00 ^b	
Breast fed	Yes	35 (19.7)	79 548 (19.4)	1.00 ^b	0.93
	No	143 (80.3)	330 421 (80.6)	0.98 (0.68, 1.42)	
Birth weight (g)	<3500 g	96 (51.1)	252 328 (58.4)	1.00 ^b	0.04
	≥ 3500 g	92 (48.9)	179 796 (41.6)	1.34 (1.01, 1.79)	
Social class	Non-manual	130 (72.6)	269 625 (64.7)	1.00 ^b	0.03
	Manual/other	49 (27.4)	147 357 (35.3)	0.69 (0.50, 0.96)	
Number of children in the household ^a	<3	51 (64.6)	113 260 (63.2)	1.00 ^b	0.81
	≥ 3	28 (35.4)	65 831 (36.8)	0.94 (0.60, 1.50)	
Number of adults in the household	<3	98 (92.5)	207 858 (87.4)	1.00 ^b	0.12
	≥ 3	8 (7.5)	29 929 (12.6)	0.57 (0.28, 1.17)	
Household density	<1 person/room	39 (49.4)	63 114 (35.4)	1.00 ^b	0.01
	≥ 1 person/room	40 (50.6)	115 001 (64.6)	0.55 (0.36, 0.88)	

^aIncludes the newborn; ^bReference category.

Table 2 Characteristics of cases of ALL and non-cases

	Cases number (%)	Non-cases number (%)
Sex		
Male	101 (53.7)	223 945 (51.5)
Female	87 (46.3)	210 785 (48.5)
Age (years)		
< 1	4 (2.1)	
1	9 (4.8)	
2	34 (18.0)	
3	27 (14.4)	
4	25 (13.3)	
5	22 (11.7)	
6–9	40 (21.3)	
10–14	26 (13.8)	
15+	1 (0.5)	
Year of birth		
1971–72	21 (11.2)	60 594 (13.9)
1973–74	23 (12.2)	53 953 (12.4)
1975–76	19 (10.1)	51 733 (11.9)
1977–78	24 (12.8)	51 490 (11.8)
1979–80	25 (13.3)	55 045 (12.7)
1981–82	24 (12.8)	53 411 (12.3)
1983–84	28 (14.9)	53 855 (12.4)
1985–86	24 (12.8)	54 645 (12.6)
Social class		
1. Academic/professional	8 (4.3)	18 045 (4.2)
2. Management/executive	35 (18.6)	74 160 (17.1)
3. Clerical/craftsmen	88 (46.3)	177 420 (40.8)
4. Semiskilled	20 (10.6)	60 805 (14.0)
5. Unskilled	12 (6.4)	33 737 (7.8)
6. Armed forces, students, unmarried unemployed, other	17 (9.0)	52 816 (12.1)
Missing data	9 (4.8)	17 762 (4.1)

years in Northern Ireland between January 1975 and December 1997. Of these, 212 were born between 1971 and 1986, and 189 (89.2%) were identified within the Child Health System. One was a multiple birth and was excluded from the analyses. This study therefore involves 188 cases of childhood ALL and 434 745 non cases. Characteristics of the identified cases are shown in Table 2. The male to female ratios of identified cases and unidentified cases were 1:1.16 and 1:1.09 respectively. Mean age at diagnosis in identified and unidentified cases was 5.19 and 5.26 years respectively, *t*-test value 0.1, *P*=0.92.

Univariable analyses

Table 1 shows the association between variables of interest and the risk of childhood ALL. There were 446 children who had a diagnosis of Down's syndrome made at birth, six of whom subsequently developed ALL (relative risk=33.0; 95% CI=14.5–74.7). High paternal and maternal age (35 years) were both associated with a modest increase in risk of ALL while the risk associated with young parental age (<25 years), especially in mothers, did not differ substantially from that of parental age between 25 and 34 years. When dichotomized into <35 and ≥35 years the relative risk associated with high paternal age was 1.47 (95% CI=1.05–2.06; *P*=0.02), while that associated with high maternal age was 1.33 (95% CI=0.90–1.97; *P*=0.15). Women who had at least one miscarriage were substantially less likely to have a child who developed ALL than women who had no history of miscarriage (relative risk=0.62; 95% CI=0.40–0.98). Gestation of more than 40 weeks appeared to be protective against future risk of ALL while the risk associated with gestation of less than 38 weeks was similar to that for 38–39 weeks. When gestational age was dichotomized into <40 weeks or ≥40 weeks, longer gestation had a relative risk of

0.76 (95% CI=0.57–1.01; *P*=0.06). Higher birth weight (≥3500 g) was associated with a modest increase in risk of ALL (relative risk=1.34; 95% CI=1.01–1.79). There was no association between risk of ALL and birth order or breast-feeding on discharge from place of confinement. The strongest associations seen were for social class at birth and living conditions in the household into which the child was born. Children whose fathers were in manual occupations at the time of birth had a risk of ALL more than 30% lower than children whose fathers were employed in non-manual occupations. Although number of children in the household at the time of birth was unrelated to risk of ALL, children born into households in which three or more adults were living appeared to have substantially reduced risk of ALL (relative risk=0.57; 95% CI=0.28, 1.17). Furthermore, being born into a more crowded home was associated with a 45% reduction in risk of ALL.

Multivariable analyses

Tables 3 and 4 show the mutually adjusted relative risks for the associations between predictor variables and risk of ALL. Not all data items were available for every subject. Model Ia (Table 3) is based on 146 cases (77.7% of all cases) and 345 473 children who did not develop ALL (79.5% of total). Model IIa (Table 4) is based on 72 children who developed ALL and 150 772 who did not; 66.7 and 62.9%, respectively, of those born in the years during which data on living conditions were collected. In Model Ia high maternal age was no longer associated with risk of ALL, although high paternal age continued to be associated with a 50% increase in risk. However, maternal and paternal age were closely correlated; correlation co-efficient 0.81, *P*<0.001. The remaining adjusted risks in Model Ia were either unaltered or stronger than the unadjusted risks.

Further adjustment for number of adults in the household and household density at birth (Model IIa) again strengthened the associations between paternal age, history of miscarriage, and gestational age, and risk of ALL. In this model, paternal age of 35 years or over was associated with an 80% increase in the risk of ALL, while history of miscarriage and longer gestational age were associated with 60 and 50% lower risks, respectively. The negative association between social class and ALL was attenuated while that for birth weight was unaltered. The strong association between household density and risk of childhood ALL seen in the univariable analysis remained in this multivariable model, with children born into less crowded households having almost twice the risk of ALL compared to children from more crowded households.

Restricting the multivariate models to only those cases diagnosed before 6 years (Models Ib and IIb, Tables 3 and 4 respectively) either strengthened or did not alter the relative risks observed in the models including all cases. Maternal age was the exception – the direction of the relative risk observed changed from weakly positive to weakly negative between Models Ia and Ib, and between Models IIa and IIb.

DISCUSSION

A major strength of this study is that it involved follow-up of a whole population for virtually the full period of risk of childhood ALL. As far as we are aware it is only the third such population-based cohort study of childhood ALL and it is unique in that the data, collected at the time of birth, included information on childhood socioeconomic circumstances and household living conditions. The use of several different sources of data to identify incident cases of ALL in a relatively small geographic region, where few centres treat this condition will have resulted in a very high level of case ascertainment. Selection bias through inclusion of only those cases treated at specific centers has been avoided. Non-inclusion of cases was uncommon and was due only to failure to

Table 3 Association between early life factors and birth characteristics and risk of childhood ALL, multivariate analysis – model I

Variable	Model Ia		Model Ib	
	Relative risk	P-value	Relative risk	P-value
Female sex	0.89 (0.64, 1.24)	0.49	0.94 (0.643, 1.41)	0.73
Down's syndrome	35.0 (14.0, 88.0)	<0.001	44.3 (15.7, 124.8)	<0.001
Maternal age ≥ 35 years	1.03 (0.61, 1.75)	0.91	0.79 (0.40, 1.50)	0.46
Paternal age ≥ 35 years	1.49 (0.96, 2.31)	0.07	1.68 (1.00, 2.82)	0.05
History of miscarriage (1 or more vs none)	0.49 (0.29, 0.80)	<0.01	0.52 (0.28, 0.96)	0.04
Gestational age ≥ 40 weeks	0.67 (0.48, 0.94)	0.02	0.50 (0.33, 0.76)	<0.01
Birth weight ≥ 3500 g	1.66 (1.18, 2.33)	<0.01	1.84 (1.22, 2.80)	<0.01
Social class (non-manual vs manual/other)	0.68 (0.47, 1.00)	0.05	0.61 (0.38, 1.00)	0.05

Table 4 Association between early life factors and birth characteristics and risk of childhood ALL, multivariate analyses – model II

Variable	Model IIa		Model IIb	
	Relative risk	P-value	Relative risk	P-value
Female sex	1.02 (0.64, 1.63)	0.94	1.11 (0.63, 1.95)	0.72
Down's syndrome	24.1 (5.7, 102.50)	<0.001	38.8 (8.9, 168.6)	<0.001
Maternal age ≥ 35 years	1.08 (0.51, 2.29)	0.84	0.72 (0.27, 1.88)	0.49
Paternal age ≥ 35 years	1.78 (0.96, 3.29)	0.07	1.92 (0.93, 3.94)	0.08
History of miscarriage (1 or more vs none)	0.42 (0.19, 0.92)	0.03	0.46 (0.18, 1.17)	0.10
Gestational age ≥ 40 weeks	0.52 (0.34, 0.85)	<0.01	0.40 (0.22, 0.72)	<0.01
Birth weight ≥ 3500 g	1.58 (0.98, 2.56)	0.06	1.71 (0.96, 3.05)	0.07
Social class (non-manual vs manual/other)	0.76 (0.43, 1.33)	0.34	0.69 (0.34, 1.41)	0.31
Three or more adults in the household	0.58 (0.21, 1.61)	0.29	0.70 (0.21, 2.30)	0.56
Household density (≥ 1 person per room)	0.56 (0.35, 0.91)	0.02	0.50 (0.28, 0.89)	0.02

identify their birth records, which with the exception of children moving into Northern Ireland was likely to be entirely random. The data used in the analyses were documented before the onset of illness, thus avoiding recall bias and observer bias. Inclusion of all live births also avoids the biases associated with control selection and participation in case-control studies. Furthermore, as the Northern Irish population is ethnically homogenous, racial and genetic factors which may obscure important associations in more mixed populations may be discounted.

The study has some potential limitations. The relatively small number of cases precludes important subgroup analyses. Birth weight was the only available indicator of prenatal development, and data on living conditions within the household were not collected for the whole study period. Data were collected by many observers and, although attempts were made to standardize procedures, variation in recording is likely to have occurred. However, such misclassification bias will result in the underestimation of true associations. The matching process will not have identified ALL cases born outside marriage, who were initially given their mother's surname and had a subsequent name change such as when their mother married. As unmarried mothers are more likely to be younger and live with relatives (higher household densities) than their married counterparts, failure to identify these cases will have reduced the power of the study to detect associations between household density, or young maternal age, and ALL. However, as birth records were identified for almost 90% of possible cases, this is a minor deficiency.

Birth weight

We found a reasonably strong positive association between birth weight and risk of childhood ALL. To date, the findings on birth weight and risk of childhood leukaemia have been inconsistent.

Several studies have reported positive relationships with ALL (Robison *et al*, 1987; Kaye *et al*, 1991; Westergaard *et al*, 1997) while others (McKinney *et al*, 1987, 1999; Roman *et al*, 1997; Dockerty *et al*, 1999), found no association or a relationship with both high and low birth weight (Schuz *et al*, 1999). It is possible that selection bias, particularly among controls, may have resulted in the failure of some investigators to observe a positive association as controls are likely to be of higher socio-economic position and thus have infants of higher average birth weight, than the general population (Kogan, 1995). Our study did not have this problem and multivariable analyses suggest that differences in birth weight across social classes do not explain our observation. Choice of threshold for high birth weight may also contribute to the inconsistency between reports. We found that the relationship between high birth weight and risk of ALL was much less obvious if a threshold of 4000 g rather than 3500 g was chosen and at least one other group has found the threshold employed to be crucial (Kaye *et al*, 1991). This sensitivity to the choice of threshold may suggest that the association is spurious. On the other hand, we displayed the same relationship, and a similar magnitude of effect, as one other population-based cohort (Westergaard *et al*, 1997). The relationship is also biologically plausible and evidence is accumulating to suggest that it may be mediated through IGF-1 (Ross *et al*, 1996; Yeazel *et al*, 1997; Petridou *et al*, 1999; Blatt, 2000; Eshet *et al*, 2000).

Parental age

Although higher maternal age has generally been associated with higher risk of ALL (Kaye *et al*, 1991; Dockerty *et al*, 1999; Hemminki *et al*, 1999; Mogren *et al*, 1999), not all studies have observed this relationship (Shaw *et al*, 1984; Van Steensel-Moll *et al*, 1985; Shu *et al*, 1988; Westergaard *et al*, 1997; Rosenbaum *et*

al, 2000) and one has demonstrated higher risk at younger maternal age (Schuz *et al*, 1999). We did find any association with maternal age. Paternal age has been infrequently examined as a risk factor for childhood ALL, with reports of both a positive association (Kaye *et al*, 1991) and no association (Shaw *et al*, 1984; Westergaard *et al*, 1997; Hemminki *et al*, 1999). We demonstrated a strong relationship between ALL and paternal age of 35 years or more. However, paternal age is highly correlated with maternal age and inclusion of the two variables in the same model may have an unpredictable effect on observed relative risks. In our univariable analyses paternal age was more strongly related to risk of ALL than was maternal age, indicating that higher paternal age may be a risk factor for childhood ALL independently of maternal age.

Gestational age and previous history of miscarriage

We found gestation beyond 40 weeks to be associated with a substantial decrease in the risk of childhood ALL. Most previous reports failed to find an association between ALL and length of gestation (Kaye *et al*, 1991; Kaatsch *et al*, 1996; McKinney *et al*, 1999), although one study has shown a similar, but not statistically significant, relationship (Dockerty *et al*, 1999). Some studies have reported positive associations between history of previous foetal loss and childhood leukaemia (van Steensel-Moll *et al*, 1985; Kaye *et al*, 1991), others found no association (Schuz *et al*, 1999) while in the current study, as in a Chinese case-control study (Shu *et al*, 1988) an inverse association was found. Possible mechanisms for a positive association include environmental exposures and genetic predisposition but the issues of selective participation and recall bias have not been fully addressed. As our findings are free from such biases they may therefore reflect the true relationship between prior history of miscarriage and risk of childhood ALL.

Socioeconomic status and household density

As in this study, some investigators have shown that higher socio-economic position, is associated with higher risk of ALL (McWhirter, 1982; Kaye *et al*, 1991; Dockerty *et al*, 1999) but others have failed to observe this association (Shaw *et al*, 1984; van Steensel-Moll *et al*, 1985, 1986; Shu *et al*, 1988; Kaatsch *et al*, 1996; McKinney *et al*, 1999; Rosenbaum *et al*, 2000) and some have found the opposite pattern, both for childhood ALL (Brondum *et al*, 1999; Dockerty *et al*, 1999) and infant leukaemia (Ross *et al*, 1997). In some of these studies subjects from higher socio-economic and better educational backgrounds are overrepresented among controls, reducing the ability to detect a positive association between socio-economic position and childhood ALL. This was not a problem in the current study and the availability of data on social class recorded at the time of birth adds to the robustness of our findings. The positive relationship with social class is further supported by the strong inverse association we demonstrated between household density and risk of childhood ALL. This finding is in keeping with the inverse (although not statistically significant), relationship between household density and childhood ALL which was observed in a case control study from New Zealand (Dockerty *et al*, 1999). However, as far as we are aware, this is the first time

the association has been demonstrated in a cohort study and our findings are likely to reflect the true relationship between socio-economic position, household density and risk of childhood ALL.

Our finding of around a two-fold greater risk of childhood ALL among children born into surroundings where there is little overcrowding supports the hypothesis that childhood ALL is caused by an inappropriate immunological response to a common infection or infections in children whose immune systems are not 'programmed' by early exposure to these infections (Greaves and Alexander, 1993; Greaves, 1997). In support of this hypothesis, several studies have reported that early exposure to infection (van Steensel-Moll *et al*, 1986; Kaatsch *et al*, 1996; McKinney *et al*, 1999) is associated with a reduced risk of childhood ALL. However, contrary to what may be expected if this hypothesis is correct, we and several other investigators (Shaw *et al*, 1984; Shu *et al*, 1988; Kaye *et al*, 1991; McKinney *et al*, 1999; Schuz *et al*, 1999; Rosenbaum *et al*, 2000) did not find that later-born infants had a lower risk of ALL than first-born infants. Also we did not demonstrate a relationship between risk of ALL and number of children in the household, while being born into a household containing three or more adults appeared to be protective. If early exposure to infections in childhood protects against ALL, transmission of infections from adults to children may be more important.

CONCLUSION

This population based cohort study is the first such study to demonstrate that being born into a household of high socio-economic position or low household density is a risk factor for the development of childhood ALL. There is also evidence that being born into an extended family may be protective. These findings offer support for the theory that modulation of the immune system by early exposure to infection is important in reducing future risk of ALL. Further investigation of this hypothesis is warranted, including the examination of infections that are likely to be transmitted, within households, from adults to children.

ACKNOWLEDGEMENTS

We wish to thank the haematologists and oncologists treating patients with ALL in Northern Ireland who have over the years provided data on their patients to relevant registers. We would also like to thank the midwives, health visitors and staff of the Northern Ireland Child Health System for the efforts in collecting the birth record data. Thanks are also due to the Department of Health and Social Services and Public Safety for allowing us access to the Child Health System Data and for the provision of support to the Northern Ireland Cancer Registry. We appreciate the work of Tom Wylie and Gerard Savage from the Northern Ireland Cancer Registry in managing this data and we also appreciate the statistical assistance provided by Dr Chris Patterson, Department of Epidemiology and Public Health, The Queen's University of Belfast.

REFERENCES

- Blatt J (2000) IGF1 and Leukemia. *Pediatr Hematol Oncol* **17**: 199–201
- Brondum J, Shu XO, Steinbuch M, Severson RK, Potter JD, Robison LL (1999) Parental cigarette smoking and the risk of acute leukemia in children. *Cancer* **85**: 1380–1388
- Dockerty JD, Skegg DC, Elwood JM, Herbison GP, Becroft DM, Lewis ME (1999) Infections, vaccinations, and the risk of childhood leukemia. *Br J Cancer* **80**: 1483–1489
- Eshet R, Zaizov R, Freud E, Laron Z (2000) Decreased insulin-like growth factor-1 receptor sites on circulating mononuclear cells from children with acute leukemia. *Pediatr Hematol Oncol* **17**: 253–260
- Greaves MF, Alexander FE (1993) An infectious aetiology for common acute lymphoblastic leukemia in childhood? *Leukemia* **7**: 349–360
- Greaves MF (1997) Aetiology of acute leukemia. *Lancet* **349**: 344–349

- Greaves M (1999) Molecular genetics, natural history and the demise of childhood leukemia. *Eur J Cancer Prev* **35**: 173–185
- Hemminki K, Kyyronen P, Vaitinen P (1999) Parental age as a risk factor of childhood leukemia and brain cancer in offspring. *Epidemiology* **10**: 271–275
- Kaatsch P, Kaletch U, Krummenauer F, Meinert R, Miesner A, Haaf G, Michaelis J (1996) Case-control study on childhood leukemia in Lower Saxony, Germany. Basic considerations, methodology and summary of results. *Klin Paediatr* **208**: 179–185
- Kaye SA, Robison LL, Smithson WA, Gunderson P, King FL, Neglia JP (1991) Maternal reproductive history and birth characteristics in childhood acute lymphoblastic leukemia. *Cancer* **68**: 1351–1355
- Kinlen LJ (1995) Epidemiological evidence for an infective basis in childhood leukemia. *Br J Cancer* **71**: 1–5
- Kogan MD (1995) Social causes of low birth weight. *J R Soc Med* **88**: 611–615
- McKinney PA, Cartwright RA, Saiu JM, Mann JR, Stiller CA, Draper GJ, Hartley AL, Hopton PA, Birch JM, Waterhouse JA (1987) The inter-regional epidemiological study of childhood cancer (IRESCC): a case-control study of aetiological factors in leukemia and lymphoma. *Arch Dis Child* **62**: 279–287
- McKinney PA, Juszcak E, Findlay E, Smith K, Thomson CS (1999) Pre- and perinatal risk factors for childhood leukemia and other malignancies: a Scottish case control study. *Br J Cancer* **80**: 1844–1851
- McWhirter WR (1982) The relationship of incidence of childhood lymphoblastic leukemia to social class. *Br J Cancer* **46**: 640–645
- Mogren I, Damber L, Tavelin B, Hogberg U (1999) Characteristics of pregnancy and birth and malignancy in the offspring (Sweden). *Cancer Causes Control* **10**: 85–94
- Petridou E, Dessypris N, Spanos E, Mantzoros C, Skalkidou A, Kalmanti M, Koliouskas D, Kosmidis H, Panagiotou JP, Piperopoulou F, Tzortatou F, Trichopoulos D (1999) Insulin-like growth factor-I and binding protein-3 in relation to childhood leukemia. *Int J Cancer* **80**: 494–496
- Robison LL, Codd M, Gunderson P, Neglia JP, Smithson WA, King FL (1987) Birth weight as a risk factor for acute lymphoblastic leukemia. *Pediatr Hematol Oncol* **4**: 63–72
- Roman E, Ansell P, Bull D (1997) Leukemia and non-Hodgkin's lymphoma in children and young adults: are prenatal and neonatal factors important determinants of disease. *Br J Cancer* **76**: 406–415
- Rosenbaum PF, Buck GM, Brecher ML (2000) Early child-care and preschool experiences and risk of childhood acute lymphoblastic leukemia. *Am J Epidemiol* **152**: 1136–1144
- Ross JA, Perentesis JP, Robison LL, Davies SM (1996) Big babies and infant leukemia: a role for insulin-like growth factor-1? *Cancer Causes Control* **7**: 553–559
- Ross JA, Potter JD, Shu XO, Reaman GH, Lampkin B, Robison LL (1997) Evaluating the relationships among maternal reproductive history, birth characteristics, and infant leukemia – a report from the Children's Cancer Group. *Ann Epidemiol* **7**: 172–179
- Schuz J, Kaatsch P, Meinert R, Michaelis J (1999) Association of childhood cancer with factors related to pregnancy and birth. *Int J Epidemiol* **28**: 631–639
- Shaw G, Lavey R, Jackson R, Austin D (1984) Association of childhood leukemia with maternal age, birth order, and paternal occupation. *Am J Epidemiol* **119**: 788–795
- Shu XO, Gao YT, Brinton LA, Linet MS, Tu JT, Zheng W, Fraumeni JF (1988) A population-based case-control study of childhood leukemia in Shanghai. *Cancer* **62**: 635–644
- Van Steensel-Moll HA, Valkenburg HA, Vandenbroucke JP, van Zanen GE (1985) Are maternal fertility problems related to childhood leukemia. *Int J Epidemiol* **14**: 555–559
- Van Steensel-Moll HA, Valkenburg HA, van Zanen GE (1986) Childhood leukemia and infectious diseases in the first year of life: a register-based case-control study. *Am J Epidemiol* **124**: 590–594
- Westergaard T, Andersen PK, Pedersen JB, Olsen JH, Frisch M, Sorensen HT, Wohlfahrt J, Melbye M (1997) Birth characteristics, sibling patterns, and acute leukemia risk in childhood: a population-based cohort study. *J Natl Cancer Inst* **89**: 939–947
- Wiemels JL, Cazzaniga G, Daniotti M, Eden OB, Addison GM, Masera G, Saha V, Biondi A, Greaves MF (1999) Prenatal origin of acute lymphoblastic leukemia in children. *Lancet* **354**: 1499–1503
- Yeazel MW, Ross JA, Buckley JD, Woods WG, Ruccione K, Robison LL (1997) High birth weight and risk of specific childhood cancers: a report from the Children's Cancer Group. *J Pediatr* **131**: 671–677