Review

Chemotherapy for malignant pleural mesothelioma: past results and recent developments

S Tomek^{*,1}, S Emri², K Krejcy³ and C Manegold⁴

¹Department of Medicine I, Clinical Division of Oncology, Vienna, Austria; ²Department of Chest Disease, Hacettepe University School of Medicine, Ankara, Turkey; ³Eli Lilly Regional Operations GesmbH Area Medical Centre, Vienna, Austria; ⁴Clinical Division of Oncology, Department of Medicine, Thoraxklinik Heidelberg, Germany

This review summarises the results of previously conducted clinical trials, and subsequently presents data arising from all phase II–III studies on chemotherapy of malignant pleural mesothelioma (MPM) published since the last relevant overview. While response rates exceeding 30% have barely been achieved with established cytotoxic drugs in MPM therapy, novel chemotherapeutic agents and their combinations appear more promising. This applies especially to the antimetabolites, and in particular to pemetrexed that produced response rates of up to 45% in combination with platinum compounds. Raltitrexed combined with oxaliplatin has also been shown to be effective, and gemcitabine – applied as a single agent or in combination with cisplatin – as well as vinorelbine appear to improve the quality of life in patients presenting with MPM. Data can now be more precisely analysed by increasingly implemented randomised studies, applying a standardised staging system, and distinguishing prognostic groups. While chemotherapy for MPM remains a challenging task, important steps have clearly been made in the past years to combat this aggressive disease. The publication of pemetrexed with cisplatin phase III results in a peer-reviewed journal may soon establish a standard of care.

British Journal of Cancer (2003) **88**, 167–174. doi:10.1038/sj.bjc.6600673 www.bjcancer.com

© 2003 Cancer Research UK

Keywords: malignant pleural mesothelioma; chemotherapy

Although MPM aetiology is well known, therapeutic success with this disease has been unsatisfactory. Standard malignant pleural mesothelioma (MPM) therapy is still deficient, and decisions for surgical, radiotherapy or multimodal procedures are made on a case-by-case basis. In the majority of cases, a palliative treatment approach remains the only choice. The possibilities to resect MPM curatively are rare and only given in early-stage disease. Yet even at this point, resection is a matter of dispute because of historically high morbidity and mortality rates, relapse tendencies and disappointing long-term survival rates (Rusch and Venkatraman, 1996; Boutin *et al*, 1998). Irradiation for MPM assists in repelling tumour growth and temporarily relieving pain, but not attaining appreciably lengthened overall survival time (Kutcher *et al*, 1987; Soubra *et al*, 1990; Bissett *et al*, 1991). There is also no clear evidence that multimodality protocols involving surgery, radiotherapy and chemotherapy can improve survival (Antman *et al*, 1980; Hilaris *et al*, 1984; Linden *et al*, 1996). Although success in MPM chemotherapy has been limited, it now gives rise to promise by the use of an established and standardised staging system, with consideration given to prognostic factors and development of novel cytostatic agents.

ROLE OF CHEMOTHERAPY

Chemotherapy for MPM continues to be challenging. A multitude of cytotoxic drugs have been tested both as single agents and in combination chemotherapy regimens. The rates of objective tumour regression have only been between 10 and 30% with cytotoxic monotherapy, having no significant impact on median survival. Combination chemotherapy has shown no clear advantage over single-agent therapy (Linden *et al*, 1996).

The evaluation of systemic chemotherapy in MPM has been problematic for several reasons. Owing to the rarity of this disease, only a few randomised studies based on large numbers of participants have been implemented to provide statistically significant statements regarding response to a particular therapy.

Furthermore, inadequate imaging procedures and nonuniform staging systems have complicated data interpretation. As performed in early studies, restaging by means of chest radiography failed to determine response to a given therapy accurately. Response rates have been more reliable since CT scanning has been recognised as a diagnostic tool. In 1995, the International Mesothelioma Interest Group (IMIG) proposed a novel TNM staging system, which was designed to record data concerning the natural history of the disease and was validated in two large surgical series of mesothelioma patients (Rusch and Venkatraman, 1996; Pass *et al*, 1998). The universal application of this system has ever since allowed for a more prudent evaluation of results emerging from clinical studies (Pass *et al*, 1998; Steele and Rudd, 2000).

Early clinical trials of MPM patients included heterogeneous groups of patients with divergent risk factors and were therefore

*Correspondence: Dr S Tomek, Department of Medicine I, Clinical Division of Oncology, University Hospital Vienna, Waehringer Strasse 18-20, A-1090 Vienna, Austria; E-mail: Sandra.Tomek@akh-wien.ac.at
Received 15 May 2002; revised 30 September 2002; accepted 9 October 2002

often not powerful enough in assessing therapeutic efficacy of a particular treatment. In 1998, the European Organization for Research and Treatment of Cancer (EORTC) identified several prognostic variables for the course of the disease. In a multivariate analysis of the EORTC, poor prognosis was associated with the sarcomatous histologic subtype, male gender, poor performance status and a high white blood cell count (Curran *et al*, 1998). Likewise, the Cancer and Leukemia Group B (CALGB) analysed several pretreatment factors pooled from seven phase II studies that were predictive of poor survival and defined six prognostic groups. Poor prognosis was seen in patients with the following criteria: age older than 75 years, poor performance status, chest pain, dyspnoea, weight loss, high white blood cell count, elevated platelet count, low haemoglobin, elevated serum lactate dehydrogenase levels, pleural effusion and nonepithelial histology.

Owing to the establishment of prognostic scoring systems, identification of risk groups facilitated improved study design by evaluating more homogeneous patient groups or risk-group stratification in the course of randomisation (Curran *et al*, 1998).

The following sections summarise conventional single-agent and combination chemotherapy strategies. Results emerging from recent clinical studies with novel cytotoxic agents and their combinations are then presented, which have been published since the last review of chemotherapy in MPM.

Tables 1–3 summarise phase II–III single-agent and combination chemotherapy trials for MPM including more than 15 patients, which have been conducted since 1995.

Single-agent chemotherapy

Doxorubicin is the most frequently investigated chemotherapeutic agent in the treatment of MPM. Studies have failed to corroborate evidence, however, for encouraging response rates to doxorubicin of up to 20% that had been reported in earlier studies (Aisner and Wiernik, 1981; Antman and Corson, 1985). Likewise, newer anthracyclines such as epirubicin, detorubicin, pirarubicin and mitoxantrone have shown low levels of efficacy and have offered no clinically relevant advantage over doxorubicin (Colbert *et al*, 1985; Eisenhauer *et al*, 1986; Sridhar *et al*, 1989; Kaukel *et al*, 1990; Magri *et al*, 1991; van Breukelen *et al*, 1991; Mattson *et al*, 1992; Magri *et al*, 1992). In summary, the overall response rate produced by anthracyclines applied in MPM appears to be no higher than 15% and median survival does not exceed 8 months.

Apart from anthracyclines, several studies have investigated the platinum compounds, cisplatin and carboplatin. Single-agent cisplatin resulted in a response rate of merely 14.3% and a median survival of 7.5 months (Zidar *et al*, 1988). Studies using the newer compound carboplatin resulted in similar response rates ranging between 6 and 16% (Mbidde *et al*, 1986; Raghavan *et al*, 1990; Vogelzang *et al*, 1990).

The alkylating agents cyclophosphamide and mitomycin have shown low-level activity in MPM therapy (Bajorin *et al*, 1987; Sorensen *et al*, 1985). Promising results initially arising from high-dose ifosfamide therapy (Alberts *et al*, 1988) could not be confirmed by subsequent studies (Falkson *et al*, 1992; Zidar

Table 1 Series of ≥ 15 patients with MPM treated with single-agent chemotherapy since 1995

Single agent	First author (year)	No. of patients	Responders		95% confidence interval (%)	Median survival (months)
			No	%		
Anthracyclines and related compounds						
Liposomal doxorubicin	Baas (2000)	32	2	6	0–20	13
Liposomal doxorubicin	Oh/ (2000)	24	0	0	na	9.3
Liposomal danaurubicin	Steele (1998)	11	0	0	na	6.1
Platinum compounds						
ZD0473	Giaccone (2001)	10	Two regressions of evaluable disease		na	na
Alkylating agents						
Ifosfamide	Andersen (1999)	26	1	4	0–11	10
Etoposide i.v.	Sahmoud (1997)	49	2	4	1–15	7.3
Etoposide p.o.	Sahmoud (1997)	45	3	7	2–20	9.5
Topoisomerase interactive agents						
Camptothecin analogues						
Irinotecan	Kindler (2000)	28	0	0	10–55	7.9
Topotecan	Maksymiuk (1998)	22	0	0	na	8
Antimicrotubule agents						
Vinca alkaloids						
Vinorelbine	Steele (2000)	64	12	21	10–44	13.4
Vincristine	Martensson (1989)	23	0	0	0–14	7
Vinblastine	Cowan (1988)	20	0	0	0–16	3
Taxanes						
Docetaxel	Belani (1999)	19	1	5	0–26	na
Docetaxel	Vorobiof (2000)	22	3	14	7–46	12
Paclitaxel	v. Meerbeck (1996)	25	0	0	0–15	9.8
Paclitaxel	Vogelzang (1999)	35	Three regressions of evaluable disease		2–10	5
Antimetabolites						
Edatrexate	Kindler (1999)	20	5	25	9–49	9.6
Edatrexate+LV-rescue	Kindler (1999)	38	6	16	6–31	6.6
Gemcitabine	Kindler (2001)	17	0	0	3–13	4.1
Gemcitabine	van Meerbeck (1999)	27	2	7	1–24	8
Gemcitabine	Bischoff (1998)	16	5	31	na	na
Pemetrexed	Scagliotti (2001)	62	9	6	na	10.7

na=not applicable.

Table 2 Series of ≥ 15 patients with MPM treated with combination chemotherapy since 1995

Agent	First author (year)	No. of patients	Responders		95% confidence interval (%)	Median survival
			No.	%		
Doxorubicin-containing combinations						
Doxorubicin+cisplatin+mitomycin	Pennucci (1997)	23	5	21	7–42	11
Doxorubicin+cisplatin+mitomycin+bleomycin	Breau (1991)	25	11	44	27–63	na
Cisplatin-containing combinations						
Cisplatin+DHAC	Samuels (1998)	29	5	17	5–30	6.4
Cisplatin+etoposide	Eisenhauer (1988)	26	3	12	4–30	na
Cisplatin+gemcitabine	Byrne (1999)	21	10	48	26–69	10.3
Cisplatin+gemcitabine	Novak (2002)	52	17	33	20–46	11.2
Cisplatin+gemcitabine	Van Haarst (2000)	22	4	15	na	10
Cisplatin+Irinotecan	Nakano (1999)	15	4	27	8–55	7.1
Cisplatin+mitomycin+vinblastine	Middleton (1998)	39	8	20	na	6
Cisplatin+pemetrexed (phase I)	Thodtman (1999)	11	5	45	na	na
Cisplatin+paclitaxel	Fizazi (2000)	18	1	6	0–24	12
Other combinations						
Methotrexate+mitoxantrone+mitomycin	Pinto (2001)	22	6	32	12–51	13.5
Carboplatin+gemcitabine	Aversa (1998)	18	3	16	na	8.6
Carboplatin+pemetrexed (phase I)	Hughes (2002)	25	8	32	na	15
Oxaliplatin+raltitrexed	Fizazi (2000)	30	9	30	15–49	na
Oxaliplatin+vinorelbine	Steele (2000)	17	2	12	na	na
Docetaxel+irinotecan	Knuuttila (2000)	15	0	0	2–45	8.5

na=not applicable.

Table 3 Series of randomized phase II – III studies in patients with MPM treated with chemotherapy

Agent	First author (year)	No. of patients	Responders		95% confidence interval (%)	Median survival (months)
			No.	%		
Doxorubicin vs cyclophosphamide	Sorensen (1985)	32	0	0	0–19	na
			0	0	0–19	
Doxorubicin+cyclophosphamide vs doxorubicin+cyclophosphamide+DTIC	Samson (1987)	76	4	11	6–21	7.5
			5	13	6–21	6.3
Cisplatin+doxorubicin vs cisplatin+mitomycin	Chahinian (1993)	70	5	14	5–30	
			5	26	12–43	7.7
Cisplatin+etoposide vs carboplatin	White SC (2000)	25	1	8	na	4.7
			0	0		5.4
Onconase vs doxorubicin	Vogelzang (2000)	154	na	na	na	7.7
						8.2
Cisplatin vs cisplatin+pemetrexed	Vogelzang (2002)	456	na*	na*	na*	na*
Ongoing trials agent						
		No. of patients planned				
Doxorubicin vs doxorubicin+onconase		300				
Cisplatin vs cisplatin+raltitrexed		240				
Pemetrexed vs best supportive care		240				
Vinorelbine vs MVP vs best supportive care		840				
Cisplatin+gemcitabine vs cisplatin+gemcitabine+bevacizumab		106				

* Vogelzang N.: <http://www.asco.org/ascoMainConstructor/1,47468,,12|002351,00.asp?cat=General+Oncology>.

na=not applicable.

et al, 1992; Icli *et al*, 1996; Krarup-Hansen, 1996; Andersen *et al*, 1999).

The older vinca alkaloids vinblastine, vincristine and vindesine have demonstrated no activity in the treatment of MPM (Kelsen *et al*, 1983; Boutin *et al*, 1987; Cowan *et al*, 1988; Martensson & Sorenson, 1989). Likewise, poor results were shown with oral as well as IV etoposide (Tammilehto *et al*, 1994; Sahnoud *et al*, 1997).

The antifolates methotrexate and edatrexate have been the only single agents to produce comparatively better results. A Norwegian study reported a satisfactory response rate of 37% for high-dose

methotrexate and a median survival of 11 months, for trial subjects with an epithelial subtype drawing a particular benefit from this chemotherapy option (Solheim *et al*, 1992). Sequential multicentre phase II studies conducted by the CALGB have evaluated the activity of the folate antagonist edatrexate, with and without leucovorin rescue. Edatrexate produced 25% overall response rates but proved to be relatively toxic. Leucovorin rescue in the control arm led to decreased toxicities, but may also have reduced the agent's efficacy (Kindler *et al*, 1999). Other antimetabolites like fluorouracil, dihydro-5-azacytidine (DHAC), dideazafolic acid and trimetrexate have shown minor or no activity in the treatment of

MPM (Harvey *et al*, 1984; Vogelzang *et al*, 1994; Vogelzang *et al*, 1997; Samuels *et al*, 1998).

Combination chemotherapy

Doxorubicin-based regimens Doxorubicin is the anthracycline most frequently included in chemotherapeutic regimens. Initially encouraging response rates experienced with the doxorubicin–cisplatin combination in two phase II studies carried out in Germany (RR: 46%) and Italy (RR: 25%) (Henss *et al*, 1988; Ardizzoni *et al*, 1991) failed to be confirmed by a subsequent, randomised CALGB study of doxorubicin–cisplatin vs mitomycin–cisplatin treatment (Chahinian *et al*, 1993). The doxorubicin–cisplatin doublet was only able to produce a 14% response rate and proved inferior to the mitomycin–cisplatin combination (RR: 26%). Median survival duration from study entry was 7.7 and 8.8 months, respectively, with no significant differences between treatments. An Italian group administered the triplet doxorubicin–cisplatin–mitomycin to 24 MPM patients and reported a response rate of 20.9%. Thus, the observed level of activity was similar to that obtained with the respective doublets (Pennucci *et al*, 1997).

The combination of doxorubicin, cisplatin and cyclophosphamide was tested by another prospective trial and produced a similar rate of response (Shin *et al*, 1995). A multi-institutional randomised study was designed to compare the activity of doxorubicin–cyclophosphamide with a triplet consisting of these agents in addition to dacarbazine. With a response rate of 13%, the triplet did not prove superior to the doublet (RR: 11%) (Samson *et al*, 1987).

The combination of doxorubicin and ifosfamide has also been investigated within two studies. One study giving doxorubicin and ifosfamide every 3 weeks achieved a response rate of 12.5% (Carmichael *et al*, 1989). Based on the same trial design, another group applied dose-escalated doxorubicin at 75 mg m^{-2} , with a 32% response rate (Dirix *et al*, 1994). However, the 7-month median survival was poor and toxicity high, limiting the value of this schedule in the treatment of MPM.

Disappointing response rates have also been shown by the anthracyclines epirubicin and rubidazole as combined with ifosfamide and dacarbazine (Zidar *et al*, 1983; Magri *et al*, 1992).

Cisplatin-based regimens Apart from combinations with anthracyclines, cisplatin has also been tested in numerous other chemotherapy regimens. Three trials evaluated cisplatin combined with etoposide, showing response rates from 12 to 24% (Eisenhauer *et al*, 1988; Planting *et al*, 1995; White *et al*, 2000). The doublet combination of cisplatin and DHAC attained an insufficient response rate of 17.3% (Samuels *et al*, 1998). Moderate antitumour activity (RR: 25%) was achieved by the doublet cisplatin–mitomycin in one arm of CALGB 8435, but median survival was poor (7.7 months) (Chahinian *et al*, 1993). A 25% response rate and a 13-month mean duration of response have been indicated for the cisplatin–vinblastine combination (Tsavaris *et al*, 1994). The cisplatin–mitomycin–C–vinblastine triplet did not prove to be superior to the doublets (RR: 23%), yet produced a symptomatic benefit for 63% of the patients with particularly good response for pain (Middleton *et al*, 1998).

Likewise, other treatments combining such conventional cytotoxics as methotrexate with vincristine and mitomycin with vindesine have failed to prove effective (Dimitrov *et al*, 1982; Gridelli *et al*, 1992).

NOVEL CYTOTOXICS

Anthracyclines

Liposomal anthracyclines

Liposomal doxorubicin. Within a phase II study, the EORTC treated 35 patients with liposomal doxorubicin (Caelyx[®]) at a

dosage of 45 mg m^{-2} every 4 weeks (Baas *et al*, 2000). The drug was well tolerated, but the results were disappointing. Only two of the 31 patients (6%) showed a partial response and the median survival was 13 months.

Similar results were reported by Oh *et al*, who administered 50 mg m^{-2} of liposomal doxorubicin (Doxil[®]) every 4 weeks to 24 patients with pleural mesothelioma (Oh *et al*, 2000). With no objective responses and a median survival of only 37 weeks, the drug was considered inactive in this indication.

Liposomal daunorubicin. Liposomal daunorubicin (LD) was given to 13 patients at 120 mg m^{-2} every 3 weeks, but no responses were produced (Steele *et al*, 2001).

Platinum compounds

ZD0473

A phase II trial (Giaccone *et al*, 2001) assessed the efficacy of the new generation platinum compound ZD0473 at 120 mg m^{-2} on day 1 of a 21-day cycle in mesothelioma patients that had relapsed after previous platinum-based chemotherapy. Two of 10 (20%) patients, experienced tumour shrinkage and five of 10 (50%) subjects had stable disease.

Topoisomerase interactive agents

Camptothecin analogues

Irinotecan. The activity of single-agent irinotecan (125 mg m^{-2} given weekly for 4 weeks, every 6 weeks) in malignant mesothelioma was investigated by the CALGB (Kindler *et al*, 2000). In 28 patients evaluable for analysis, no complete or partial responses were observed and the median overall survival was 7.9 months, indicating that irinotecan, at least in this dose and schedule, had no antitumour activity and considerable toxicity (leucopenia, neutropenia, diarrhoea).

Furthermore, the combination of irinotecan and cisplatin was evaluated by a Japanese group (Nakano *et al*, 1999). A total of 15 chemo-naïve MPM patients were treated with irinotecan at 60 mg m^{-2} on days 1, 8 and 15 and cisplatin at 60 mg m^{-2} on day 1, repeated every 28 days. A response rate of 26.7% (four partial responses) was observed and the median survival after chemotherapy was 28.3 weeks. In contrast to the trial conducted by the CALGB, Nakano *et al* reported definite activity with only mild toxicity of this combination regimen and concluded that it warrants further clinical evaluation.

Recently, Verschraegen *et al* (2001) published retrospective data of the same combination in 10 patients with peritoneal mesothelioma. Six courses of irinotecan 50 mg m^{-2} (day 1, 8 and 15) and cisplatin 50 mg m^{-2} (day 1) were administered every 4 weeks, either intraperitoneally or intravenously, and were well tolerated. The authors reported that 70% of the patients improved on treatment. Owing to the observed clinical benefit, a phase II trial is currently planned.

Topotecan. In order to evaluate the efficacy and toxicity of the camptothecin analogue topotecan in the treatment of MPM, the North Central Cancer Treatment Group treated 22 MPM patients with topotecan 1.5 mg m^{-2} daily for 5 days at 3-week intervals. 18 patients (86%) experienced grade 3 or 4 neutropenia. There were no objective responses seen and the median survival for all patients was 230 days (Maksymiuk *et al*, 1998).

Antimicrotubule agents

Vinca alkaloids

Vinorelbine. Steele *et al* (2000) administered vinorelbine to 64 patients with MPM at a weekly 30 mg m^{-2} dose for 6 weeks. A response rate of 21% (12 out of 64 patients) was observed and 63% of patients experienced disease stabilisation. Furthermore, the

patients' quality of life was increased with respect to lung-related symptoms and physical well-being in general.

The same study group conducted a trial to evaluate the activity of vinorelbine at 30 mg m^{-2} on days 1 and 8 and oxaliplatin 130 mg m^{-2} on day 1 every 3 weeks in 21 MPM patients (Steele *et al*, 2000). Only two partial responses had been observed and toxicity had been significant. These inferior results could perhaps result from the large number of stage IV patients, the higher proportion of less favorable subtypes, and the inclusion of several participants showing a low performance status. At any rate, the addition of oxaliplatin to vinorelbine seems to bring no advantage, thus discouraging further studies in this doublet.

Recently, a British randomised phase III study has been initiated, comparing the efficacy of single-agent vinorelbine (30 mg m^{-2} up to 60 mg m^{-2}) vs MPV (mitomycin 8 mg m^{-2} , vinblastin 6 mg m^{-2} , cisplatin 50 mg m^{-2}) vs active best supportive care alone.

Taxanes

Paclitaxel. A total of 25 patients with MPM were given paclitaxel intravenously at a dose of 200 mg m^{-2} as a 3 h infusion every 3 weeks in a phase II study conducted by the EORTC Lung Cancer Cooperative Group (van Meerbeeck *et al*, 1996). No major objective responses were seen and the median survival time was only 39 weeks.

These disappointing results were confirmed by the CALGB, which administered paclitaxel at a slightly higher dose of 250 mg m^{-2} , given as a 24-hour infusion every 3 weeks plus filgastrim (G-CSF) support to 35 patients with MPM (Vogelzang *et al*, 1999). Only three (9%) regressions of evaluable disease were observed and the median survival was 5 months.

Fizazi *et al* (2000) conducted a phase II study of paclitaxel (200 mg m^{-2}) and cisplatin (100 mg m^{-2}), given to 18 patients on day 1 every 3 weeks. With only one partial response and a response rate of 6%, the paclitaxel–cisplatin doublet was considered ineffective and the trial was stopped early.

Bednar (1999) evaluated the combination of paclitaxel at 175 mg m^{-2} and carboplatin at an AUC of 5–6 in seven patients with malignant mesothelioma and reported more encouraging results. One patient with peritoneal malignant mesothelioma had a pathologically proven CR for a duration of 20 months. The overall median survival was 15 months from diagnosis and 12 months from the onset of treatment.

Docetaxel. The Eastern Cooperative Oncology Group assigned 19 patients with malignant mesothelioma to a phase II study of docetaxel 100 mg m^{-2} , administered every 21 days. Only one patient achieved a partial response, with an overall response rate of 5%. The study group thus concluded that docetaxel was not effective in the treatment of malignant mesothelioma (Belani *et al*, 1999).

Vorobiof *et al* (2000), who also evaluated the activity of docetaxel using the same treatment schedule, reported only three partial remissions and six minor responses (25% or less reduction in tumour burden) in 22 MPM patients with median survival barely exceeding 12 months.

A clinical study further tested docetaxel at 60 mg m^{-2} in combination with irinotecan at 190 mg m^{-2} on day 1 of a 3-weekly cycle in 15 MPM patients (Knuutila *et al*, 2000). No objective responses were achieved, and median survival was 8.5 months only. The study was finally discontinued owing to high toxicity and deficient activity levels.

Antimetabolites

Antifolates

Raltitrexed. The EORTC has recently completed a phase II study of the quinazoline antifolate raltitrexed (ZD1694, Tomudex) at 3 mg m^{-2} as a single agent in the treatment of malignant mesothelioma, which is now in final analysis.

From 1999 to 2000, the Institut Gustave Roussy has treated 70 patients with a combination of raltitrexed (3 mg m^{-2}) and oxaliplatin (130 mg m^{-2}). Preliminary results show an encouraging response rate of 25% (14 of 57 patients evaluable for efficacy) and acceptable toxicity (Fizazi *et al*, 2000).

Based on these promising results, the EORTC initiated a randomised phase III study, evaluating the efficacy of the doublet raltitrexed (3 mg m^{-2})–cisplatin (80 mg m^{-2}) vs cisplatin (80 mg m^{-2}) alone.

Pemetrexed. A phase I study conducted in 1999 first investigated the activity of the multitargeted antifolate pemetrexed (LY231514, ALIMTA[®]) combined with cisplatin in patients presenting with solid tumours (Thodtmann *et al*, 1999). Of 11 mesothelioma patients evaluable for analysis, five experienced a partial remission (RR: 45%).

These promising data resulted in the largest phase III study ever conducted in patients with MPM, which was initiated in March 1999. The results of this single-blind trial were first presented at the ASCO annual meeting in May 2002. A total of 456 patients were randomised to receive either cisplatin–pemetrexed (cisplatin 75 mg m^{-2} and pemetrexed 500 mg m^{-2} on day 1, every 21 days) or cisplatin monotherapy in the control group. The combination of pemetrexed–cisplatin resulted in superior median overall survival, response rate, lung function and subjective quality-of-life measures (Vogelzang N http://www.asco.org/asco/ascoMainConstructor/1,47468,_12002351,00.asp?cat=General+Oncology). The principal investigator concluded that based on these very promising data, pemetrexed–cisplatin should now be considered standard front-line therapy for patients with MPM.

Another phase I pemetrexed study demonstrated clinical activity in combination with carboplatin. A total of 27 patients were treated with various dose levels of pemetrexed and carboplatin on day 1 of a 21-day schedule (Hughes *et al*, 2002). In 25 evaluable patients, eight (32%) achieved a partial remission and 14 experienced stable disease at various dose levels. Furthermore, a symptomatic improvement was documented in 19 cases.

A two-stage phase II trial has just been completed by Scagliotti and collaborators, investigating single-agent pemetrexed in the treatment of MPM. A total of 62 patients were given pemetrexed at 500 mg m^{-2} on day 1, every 3 weeks. Scagliotti *et al* (2001) reported a response rate of 14.5% and a median survival of 10.7 months.

Furthermore, a randomised phase III trial comparing pemetrexed plus best supportive care vs best supportive care alone in previously treated patients with MPM is currently ongoing.

Nucleoside analogues

Gemcitabine. The activity of single-agent gemcitabine at 1250 mg m^{-2} on days 1, 8 and 15 on a 28-day schedule was evaluated by the EORTC-Lung Cancer Group in 27 chemotherapy-naïve subjects with MPM (van Meerbeeck *et al*, 1999). Two partial responses were achieved for an overall response rate of 7%. An additional 56% experienced disease stabilisation and overall median survival was 8 months.

The CALGB administered high-dose gemcitabine at 1500 mg m^{-2} on days 1, 8 and 15 in a 28-day cycle to 17 participants (Kindler *et al*, 2001). Only one minor regression and six cases of stable disease were reported in 13 assessable patients, with an overall median survival of 4.1 months. In contrast to these data, Bischoff *et al* (1998) reported an encouraging response rate of 31% (5/16) with the same dose and schedule as the van Meerbeeck study (gemcitabine 1250 mg m^{-2} on days 1, 8 and 15, every 28 days). Furthermore, seven additional patients reported a symptom relief via decreased pain or dyspnoea.

An Australian study group evaluated the doublet gemcitabine–cisplatin in 21 patients presenting with advanced MPM (Byrne *et al*, 1999). The subjects received gemcitabine at 1000 mg m^{-2} on day 1, 8 and 15 and cisplatin at 100 mg m^{-2} on day 1 of a 28-day cycle. This combination chemotherapy produced an encouraging

response rate of 47.6% and an overall survival of 41 weeks. Furthermore, nine of 10 responding patients experienced significant relief of chest pain and dyspnoea. Subsequently, a multicentre study was initiated, evaluating the same chemotherapy regimen in 53 MPM patients (Novak *et al*, 2002). A response rate of 33% was achieved and the median survival time was 11.2 months. Response to treatment was accompanied by significantly improved global quality of life and respiratory function.

Contrary to the Australian trials, however, only four (15%) of 22 assessable subjects experienced a partial response within a European multicentre phase II study (Van Haarst *et al*, 2000) of gemcitabine at 1250 mg m⁻² on days 1 and 8, in addition to cisplatin at 80 mg m⁻² on day 1. The discrepancy between these studies may possibly result from the different treatment schedules, patient selection criteria and methodology applied in treatment evaluation.

In turn, Aversa *et al* (1998) evaluated the activity of the gemcitabine-carboplatin combination. A total of 20 patients were given gemcitabine at 1000 mg m⁻² on days 1, 8 and 15 and carboplatin (AUC=5) on day 1, every 28 days for a median of 4.5 cycles. In 18 assessable subjects, a response rate of 16% and an 8.6-month median survival rate was achieved.

Recently, investigators at the University of Chicago have initiated a multicenter, randomised phase II trial of cisplatin-gemcitabine and the vascular endothelial growth factor (VEGF)

inhibitor bevacizumab, which has shown preliminary evidence of activity in MPM.

CONCLUSION

With rates of objective tumour regression ranging from 10 to 30% for cytotoxic monotherapy, diffuse pleural mesothelioma was considered to date to be widely chemoresistant. The most favourable responses to conventional chemotherapeutic agents were reported by the antimetabolites methotrexate and edatrexate.

In the past few years, a number of novel cytotoxic agents have been introduced into clinical oncology, the activity of which has also been tested in MPM therapy. Initial results produced by monotherapy based on new antimetabolites, along with platin combinations, provide encouragement. Preliminary results of the raltitrexed-oxaliplatin combination have shown promising activity. Single-agent gemcitabine and the drug combined with cisplatin appear to decrease symptoms associated with tumor load. In particular, pemetrexed with cisplatin has demonstrated superior median overall survival, response rate, lung function and quality-of-life measures in the largest randomised trial in MPM. These promising data suggest that finally effective chemotherapy exists for this aggressive disease and that pemetrexed-cisplatin will become a new systemic therapy standard.

REFERENCES

- Aisner J, Wiernik PH (1981) Chemotherapy in the treatment of malignant mesothelioma. *Semin Oncol* 8: 335-343
- Alberts AS, Falkson G, Van Zyl L (1988) Malignant pleural mesothelioma: phase II pilot study of ifosfamide and mesna. *J Natl Cancer Inst* 80: 698-700
- Andersen MK, Krarup-Hansen A, Martensson G, Winther-Nielsen H, Thylen A, Damgaard K, Olling S, Wallin J (1999) Ifosfamide in malignant mesothelioma: a phase II study. *Lung Cancer* 24: 39-43
- Antman KH, Blum RH, Greenberger JS, Flowerdew G, Skarin AT, Canellos GP (1980) Multimodality therapy for malignant mesothelioma based on a study of natural history. *Am J Med* 68: 356-362
- Antman KH, Corson JM. (1985) Benign and malignant pleural mesothelioma. *Clin Chest Med* 6: 127-140
- Ardizzone A, Rosso R, Salvati F, Fusco V, Cinquegrana A, De Palma M, Serrano J, Pennucci MC, Soresi E, Crippa M *et al* (1991) Activity of doxorubicin and cisplatin combination chemotherapy in patients with diffuse malignant pleural mesothelioma. An Italian Lung Cancer Task Force (FONICAP) Phase II study. *Cancer* 67: 2984-2987
- Aversa SL, Crcuri C, DePangher V *et al* (1998) Carboplatin and Gemcitabine chemotherapy for malignant pleural mesothelioma (MPM): A phase II study of the GSTPV. *Ann of Oncol* 9: 117
- Baas P, van Meerbeeck J, Groen H, Schouwink H, Burgers S, Daamen S, Giaccone G (2000) Caelyx in malignant mesothelioma: a phase II EORTC study. *Ann Oncol* 11: 697-700
- Bajorin D, Kelsen D, Mintzer DM (1987) Phase II trial of mitomycin in malignant mesothelioma. *Cancer Treat Rep* 71: 857-858
- Bednar ME, Chahinian P (1999) Paclitaxel and carboplatin for malignant mesothelioma. *Ann Meet Am Soc Clin Oncol* A1916 18: 496a
- Belani CP, Adak S, Aisner S, Stella PJ, Levitan N, Johnson DH (1999) Docetaxel for malignant mesothelioma: phase II study of the eastern Cooperative Oncology Group (ECOG 2595). *Proc Am Soc Clin Oncol* 1999 A1829, Vol 18, p.p. 474a
- Bischoff HG, Manegold C, Knopp M, Blatter J, Drings P (1998) Gemcitabine (Gemzar®) may reduce tumor load and tumor associated symptoms in malignant pleural mesothelioma. *Proc Ann Meet Am Soc Clin Oncol* 17: A1784
- Bissett D, Macbeth FR, Cram I (1991) The role of palliative radiotherapy in malignant mesothelioma. *Clin Oncol (R Coll Radiol)* 3: 315-317
- Boutin C, Irissou M, Guerin JC, Roegel E, Paramelle B, Brambilla C, Jeannin L, Dabouis G, Le Caer H, Viallat JR (1987) Phase II trial of vindesine in malignant pleural mesothelioma. *Cancer Treat Rep* 71: 205-206
- Boutin C, Schlessner M, Frenay C, Astoul P (1998) Malignant pleural mesothelioma. *Eur Respir J* 12: 972-981
- Breau JL, Boaziz C, Morere JJF (1991) Combination therapy with cisplatin, adramycin, bleomycin and mitomycin C plus systemic and intra-pleural hyaluronidase in 25 consecutive cases of stages II, III pleural mesothelioma. Presented at the first International Mesothelioma Conference, Paris, France
- Byrne MJ, Davidson JA, Musk AW, Dewar J, van Hazel G, Buck M, de Klerk NH, Robinson BW (1999) Cisplatin and gemcitabine treatment for malignant mesothelioma: a phase II study. *J Clin Oncol* 17: 25-30
- Carmichael J, Cantwell BM, Harris AL (1989) A phase II trial of ifosfamide/mesna with doxorubicin for malignant mesothelioma. *Eur J Cancer Clin Oncol* 25: 911-912
- Chahinian AP, Antman K, Goutsou M, Corson JM, Suzuki Y, Modeas C, Herndon II JE, Aisner J, Ellison RR, Leone L *et al* (1993) Randomised phase II trial of cisplatin with mitomycin or doxorubicin for malignant mesothelioma by the Cancer and Leukemia Group B. *J Clin Oncol* 11: 1559-1565
- Colbert N, Vannetzel JM, Izrael V, Schlienger M, Milleron B, Blanchon F, Herman D, Akoun G, Roland J, Chatelet F *et al* (1985) A prospective study of doxorubicin in malignant mesothelioma. *Cancer* 56: 2170-2174
- Cowan JD, Green S, Lucas J, Weick JK, Balcerzak SP, Rivkin SE, Coltman CA, Baker LH (1988) Phase II trial of five day intravenous infusion vinblastine sulfate in patients with diffuse malignant mesothelioma: a Southwest Oncology Group study. *Invest New Drugs* 6: 247-248
- Curran D, Sahnoud T, Therasse P, van Meerbeeck J, Postmus PE, Giaccone G (1998) Prognostic factors in patients with pleural mesothelioma: the European Organization for Research and Treatment of Cancer experience. *J Clin Oncol* 16: 145-152
- Dimitrov NV, Egner J, Balceva E, Suhrlund LG (1982) High-dose methotrexate with citrovorum factor and vincristine in the treatment of malignant mesothelioma. *Cancer* 50: 1245-1247
- Dirix LY, van Meerbeeck J, Schrijvers D, Corthouts B, Prove A, van Marck E, Vermeire P, van Oosterom AT (1994) A phase II trial of dose-escalated doxorubicin and ifosfamide/mesna in patients with malignant mesothelioma. *Ann Oncol* 5: 653-655
- Eisenhauer EA, Evans WK, Murray N, Kocha W, Wierzbicki R, Wilson K (1988) A phase II study of VP-16 and cisplatin in patients with unresectable malignant mesothelioma. An NCI Canada Clinical Trials Group Study. *Invest New Drugs* 6: 327-329

- Eisenhauer EA, Evans WK, Raghavan D, Desmeules MJ, Murray NR, Stuart-Harris R, Wilson KS (1986) Phase II study of mitoxantrone in patients with mesothelioma: a National Cancer Institute of Canada Clinical Trials Group Study. *Cancer Treat Rep* 70: 1029–1030
- Falkson G, Hunt M, Borden EC, Hayes JA, Falkson CI, Smith TJ (1992) An extended phase II trial of ifosfamide plus mesna in malignant mesothelioma. *Invest New Drugs* 10: 337–343
- Fizazi K, Caliendo R, Soulie P, Fandi A, Daniel C, Bedin A, Doubre H, Viala J, Rodier J, Trandafir L, Le Chevalier T, Cvitkovic E, Armand J, Ruffie P (2000) Combination raltitrexed (Tomudex(R))-oxaliplatin: a step forward in the struggle against mesothelioma? The Institut Gustave Roussy experience with chemotherapy and chemo-immunotherapy in mesothelioma. *Eur J Cancer* 36: 1514–1521
- Giaccone G, O'Brien M, Byrne MJ, Van Steenkiste J, Cosaert J (2001) Phase II Trial of ZD0473 in Patients with Mesothelioma Relapsing After One Prior Chemotherapy Regimen. *Proc Ann. Meet Am Soc Clin Oncol* A2781 20: 257b
- Gridelli C, Pepe R, Airoma G, Incoronato P, Rossi A, Palazzolo G, Bianco AR (1992) Mitomycin C and vindesine: an ineffective combination chemotherapy in the treatment of malignant pleural mesothelioma. *Tumori* 78: 380–382
- Harvey VJ, Slevin ML, Ponder BA, Blackshaw AJ, Wrigley PF (1984) Chemotherapy of diffuse malignant mesothelioma. Phase II trials of single-agent 5-fluorouracil and adriamycin. *Cancer* 54: 961–964
- Henss H, Fiebig HH, Schildge J, Arnold H, Hasse J (1988) Phase-II study with the combination of cisplatin and doxorubicin in advanced malignant mesothelioma of the pleura. *Onkologie* 11: 118–120
- Hilaris BS, Nori D, Kwong E, Kutcher GJ, Martini N (1984) Pleurectomy and intraoperative brachytherapy and postoperative radiation in the treatment of malignant pleural mesothelioma. *Int J Radiat Oncol Biol Phys* 10: 325–331
- Hughes A, Calvert P, Azzabi A, Plummer A, Johnsons R, Rusthoven R, Griffin M, Fishwick K, Boddy AV, Verril M, Calvert H (2002) Phase I clinical and pharmacokinetic study of pemetrexed and carboplatin in patients with malignant pleural mesothelioma. *J Clin Oncology* 20: 3533–3544
- Icli F, Karaoguz H, Hasturk S, Kurt B, Akbulut H, Dincol D, Demirkazik A, Cay F, Akyar S (1996) Two dose levels of ifosfamide in malignant mesothelioma. *Lung Cancer* 15: 207–213
- Kaukel E, Koschel G, Gatzemeyer U, Salewski E (1990) A phase II study of pirarubicin in malignant pleural mesothelioma. *Cancer* 66: 651–654
- Kelsen D, Gralla R, Cheng E, Martini N (1983) Vindesine in the treatment of malignant mesothelioma: a phase II study. *Cancer Treat Rep* 67: 821–822
- Kindler HL, Belani CP, Herndon II JE, Vogelzang NJ, Suzuki Y, Green MR (1999) Edatrexate (10-ethyl-deaza-aminopterin) (NSC #626715) with or without leucovorin rescue for malignant mesothelioma. Sequential phase II trials by the cancer and leukemia group B. *Cancer* 86: 1985–1991
- Kindler HL, Herndon JE, Vogelzang NJ, Green MR. (2000). CPT-11 in Malignant Mesothelioma: A Phase II Trial by the Cancer and Leukemia Group B (CALGB 9733). *Proc Ann. Meet Am Soc Clin Oncol* 27: A1978. Vol. 19, pp. 505a
- Kindler HL, Millard F, Herndon II JE, Vogelzang NJ, Suzuki Y, Green MR (2001) Gemcitabine for malignant mesothelioma: A phase II trial by the Cancer and Leukemia Group B. *Lung Cancer* 31: 311–317
- Knuuttila A, Ollikainen T, Halme M, Mali P, Kivisaari L, Linnainmaa K, Jekunen A, Mattson K (2000) Docetaxel and irinotecan (CPT-11) in the treatment of malignant pleural mesothelioma—a feasibility study. *Anticancer Drugs* 11: 257–261
- Krup-Hansen A (1996) Studies concerning high dose ifosfamide to patients suffering from malignant mesothelioma. *Lung Cancer* 16: 101–102
- Kutcher GJ, Kestler C, Greenblatt D, Brenner H, Hilaris BS, Nori D (1987) Technique for external beam treatment for mesothelioma. *Int J Radiat Oncol Biol Phys* 13: 1747–1752
- Linden CJ, Mercke C, Albrechtsson U, Johansson L, Ewers SB (1996) Effect of hemithorax irradiation alone or combined with doxorubicin and cyclophosphamide in 47 pleural mesotheliomas: a nonrandomised phase II study. *Eur Respir J* 9: 2565–2572
- Magri MD, Foladore S, Veronesi A, Serra C, Nicotra M, Tommasi M, Grandi G, Monfardini S, Bianchi C (1992) Treatment of malignant mesothelioma with epirubicin and ifosfamide: a phase II cooperative study. *Ann Oncol* 3: 237–238
- Magri MD, Veronesi A, Foladore S, De Giovanni D, Serra C, Crismancich F, Tuveri G, Nicotra M, Tommasi M, Morassut S et al (1991) Epirubicin in the treatment of malignant mesothelioma: a phase II cooperative study. The North-Eastern Italian Oncology Group (GOCCNE)—Mesothelioma Committee. *Tumori* 77: 49–51
- Maksymiuk AW, Marschke Jr RF, Tazelaar HD, Grill J, Nair S, Marks RS, Brooks BJ, Mailliard JA, Burton GM, Jett JR (1998) Phase II trial of topotecan for the treatment of mesothelioma. *Am J Clin Oncol* 21: 610–613
- Martensson G, Sorenson S (1989) A phase II study of vincristine in malignant mesothelioma—a negative report. *Cancer Chemother Pharmacol* 24: 133–134
- Mattson K, Giaccone G, Kirkpatrick A, Evrard D, Tammilehto L, van Breukelen FJ, Planteydt HT, van Zandwijk N (1992) Epirubicin in malignant mesothelioma: a phase II study of the European Organization for Research and Treatment of Cancer Lung Cancer Cooperative Group. *J Clin Oncol* 10: 824–828
- Mbidde EK, Harland SJ, Calvert AH, Smith IE (1986) Phase II trial of carboplatin (JM8) in treatment of patients with malignant mesothelioma. *Cancer Chemother Pharmacol* 18: 284–285
- Middleton GW, Smith IE, O'Brien ME, Norton A, Hickish T, Priest K, Spencer L, Ashley S. (1998) Good symptomatic relief with palliative MVP (mitomycin-C, vinblastine and cisplatin) chemotherapy in malignant mesothelioma. *Ann Oncol* 9: 269–273
- Nakano T, Chahinian AP, Shinjo M, Togawa N, Tonomura A, Miyake M, Ninomiya K, Yamamoto T, Higashino K (1999) Cisplatin in combination with irinotecan in the treatment of patients with malignant pleural mesothelioma: a pilot phase II clinical trial and pharmacokinetic profile. *Cancer* 85: 2375–2384
- Novak A, Byrne M, Williamson R (2002) Multicentre phase II study of cisplatin and gemcitabine for malignant mesothelioma. *Br J Cancer* 87: 491–496
- Oh Y, Perez-Soler R, Fossella FV, Glisson BS, Kurie J, Walsh GL, Truong M, Shin DM (2000) Phase II study of intravenous Doxil in malignant pleural mesothelioma. *Invest New Drugs* 18: 243–245
- Pass HI, Temack BK, Kranda K, Steinberg SM, Feuerstein IR (1998) Preoperative tumor volume is associated with outcome in malignant pleural mesothelioma. *J Thorac Cardiovasc Surg* 115: 310–317 discussion 317–318
- Pennucci MC, Ardizzoni A, Pronzato P, Fioretti M, Lanfranco C, Verna A, Giorgi G, Vigani A, Frola C, Rosso R (1997) Combined cisplatin, doxorubicin, and mitomycin for the treatment of advanced pleural mesothelioma: a phase II FONICAP trial. Italian Lung Cancer Task Force. *Cancer* 79: 1897–1902
- Pinto C, Marino A, Guaraldi M, Melotti B, Piana E, Martoni A, Pannuti F (2001) Combination chemotherapy with mitoxantrone, methotrexate, and mitomycin (MMM regimen) in malignant pleural mesothelioma: a phase II study. *Am J Clin Oncol* 24(2): 143–147
- Planting AS, van der Burg ME, Goey SH, Schellens JH, van den Bent MJ, de Boer-Dennert M, Stoter G, Verweij J (1995) Phase II study of a short course of weekly high-dose cisplatin combined with long-term oral etoposide in pleural mesothelioma. *Ann Oncol* 6: 613–615
- Raghavan D, Gianoutsos P, Bishop J, Lee J, Young I, Corte P, Bye P, McCaughan B (1990) Phase II trial of carboplatin in the management of malignant mesothelioma. *J Clin Oncol* 8: 151–154
- Rusch VW, Venkatraman E (1996) The importance of surgical staging in the treatment of malignant pleural mesothelioma. *J Thorac Cardiovasc Surg* 111: 815–825 discussion 825–826
- Sahmoud T, Postmus PE, van Pottelsberghe C, Mattson K, Tammilehto L, Splinter TA, Planting AS, Sutedia T, van Pawel J, van Zandwijk N, Baas P, Rozenaals KJ, Schrijver M, Kirkpatrick A, Van Glabbeke M, Ardizzoni A, Giaccone G (1997) Etoposide in malignant pleural mesothelioma: two phase II trials of the EORTC Lung Cancer Cooperative Group. *Eur J Cancer* 33: 2211–2215
- Samson MK, Wasser LP, Borden EC, Wanebo HJ, Creech RH, Phillips M, Baker LH (1987) Randomised comparison of cyclophosphamide, imidazole carboxamide, and adriamycin versus cyclophosphamide and adriamycin in patients with advanced stage malignant mesothelioma: a Sarcoma Intergroup Study. *J Clin Oncol* 5: 86–91
- Samuels BL, Herndon II JE, Harmon DC, Carey R, Aisner J, Corson JM, Suzuki Y, Green MR, Vogelzang NJ (1998) Dihydro-5-azacytidine and cisplatin in the treatment of malignant mesothelioma: a phase II study by the Cancer and Leukemia Group B. *Cancer* 82: 1578–1584
- Scagliotti G, Shin D, Kindler H, Johnson D, Keppler U (2001) Phase II Study of ALIMTA (pemetrexed disodium, MTA) Single Agent in Patients with Malignant Pleural Mesothelioma. *Eur J Cancer* 37 (Suppl. 6): 20

- Shin DM, Fossella FV, Umsawasdi T, Murphy WK, Chasen MH, Walsh G, Komaki R, McMurtrey MJ, Hong WK (1995) Prospective study of combination chemotherapy with cyclophosphamide, doxorubicin, and cisplatin for unresectable or metastatic malignant pleural mesothelioma. *Cancer* **76**: 2230–2236
- Solheim OP, Saeter G, Finnanger AM, Stenwig AE (1992) High-dose methotrexate in the treatment of malignant mesothelioma of the pleura. A phase II study. *Br J Cancer* **65**: 956–960
- Sorensen PG, Bach F, Bork E, Hansen HH (1985) Randomised trial of doxorubicin versus cyclophosphamide in diffuse malignant pleural mesothelioma. *Cancer Treat Rep* **69**: 1431–1432
- Soubra M, Dunscombe PB, Hodson DI, Wong G (1990) Physical aspects of external beam radiotherapy for the treatment of malignant pleural mesothelioma. *Int J Radiat Oncol Biol Phys* **18**: 1521–1527
- Sridhar KS, Hussein AM, Feun LG, Zubrod CG (1989) Activity of pirarubicin (4'-O-tetrahydropyranyladriamycin) in malignant mesothelioma. *Cancer*, **63**: 1084–1091
- Steele JP, O'Doherty CA, Shamash J, Evans MT, Gower NH, Tischkowitz MD, Rudd RM (2001) Phase II trial of liposomal daunorubicin in malignant pleural mesothelioma. *Ann Oncol* **12**: 497–499
- Steele JP, Rudd RM (2000) Malignant mesothelioma: predictors of prognosis and clinical trials. *Thorax* **55**: 725–726
- Steele JP, Shamash J, Evans MT, Gower NH, Tischkowitz MD, Rudd RM (2000) Phase II study of vinorelbine in patients with malignant pleural mesothelioma. *J Clin Oncol* **18**: 3912–3917
- Tammilehto L, Maasilta P, Mantyla M, Salo J, Mattson K (1994) Oral etoposide in the treatment of malignant mesothelioma. A phase II study. *Ann Oncol* **5**: 949–950
- Thodtmann R, Depenbrock H, Blatter J, Johnson RD, van Oosterom A, Hanauske AR (1999) Preliminary results of a phase I study with MTA (LY231514) in combination with cisplatin in patients with solid tumors. *Semin Oncol* **26**: 89–93
- Tsavaris N, Mylonakis N, Karvounis N, Bacoyiannis C, Briasoulis E, Skarlos D, Pavlidis N, Stamatelos G, Kosmidis P (1994) Combination chemotherapy with cisplatin-vinblastine in malignant mesothelioma. *Lung Cancer* **11**: 299–303
- van Breukelen FJ, Mattson K, Giaccone G, van Zandwijk N, Planteydt HT, Kirkpatrick A, Dalesio O (1991) Mitoxantrone in malignant pleural mesothelioma: a study by the EORTC Lung Cancer Cooperative Group. *Eur J Cancer* **27**: 1627–1629
- Van Haarst JW, Burgers JA, Manegold CH et al (2000) Multicenter phase II study of gemcitabine and cisplatin in malignant pleural mesothelioma (MPM). Program and abstracts of the 9th World Conference on Lung Cancer; September 11–15, Tokyo, Japan. Abstract 56
- van Meerbeeck JP, Baas P, Debruyne C, Groen HJ, Manegold C, Ardizzoni A, Gridelli C, van Marck EA, Lentz M, Giaccone G (1999) A Phase II study of gemcitabine in patients with malignant pleural mesothelioma. European Organization for Research and Treatment of Cancer Lung Cancer Cooperative Group. *Cancer* **85**: 2577–2582
- van Meerbeeck J, Debruyne C, van Zandwijk N, Postmus PE, Pennucci MC, van Breukelen F, Galdermans D, Groen H, Pinson P, van Glabbeke M, van Marck E, Giaccone G (1996) Paclitaxel for malignant pleural mesothelioma: a phase II study of the EORTC Lung Cancer Cooperative Group. *Br J Cancer* **74**: 961–963
- Verschraegen C, Le D, Kudelka A, Kavanagh J, Hunt K (2001) Cisplatin and Irinotecan (CPT-11) for Peritoneal Mesothelioma. *Proc Ann Meet Am Soc Clin Oncol* **A2110** **20**: 90
- Vogelzang NJ, Goutsou M, Corson JM, Suzuki Y, Graziano S, Aisner J, Cooper MR, Coughlin KM, Green MR (1990) Carboplatin in malignant mesothelioma: a phase II study of the Cancer and Leukemia Group B. *Cancer Chemother Pharmacol* **27**: 239–242
- Vogelzang NJ, Herndon II JE, Cirrincione C, Harmon DC, Antman KH, Corson JM, Suzuki Y, Citron ML, Green MR (1997) Dihydro-5-azacytidine in malignant mesothelioma. A phase II trial demonstrating activity accompanied by cardiac toxicity. Cancer and Leukemia Group B. *Cancer* **79**: 2237–2242
- Vogelzang NJ, Herndon II JE, Miller A, Strauss G, Clamon G, Stewart FM, Aisner J, Lyss A, Cooper MR, Suzuki Y, Green MR (1999) High-dose paclitaxel plus G-CSF for malignant mesothelioma: CALGB phase II study 9234. *Ann Oncol* **10**: 597–600
- Vogelzang NJ, Weissman LB, Herndon II JE, Antman KH, Cooper MR, Corson JM, Green MR (1994) Trimetrexate in malignant mesothelioma: A Cancer and Leukemia Group B Phase II study. *J Clin Oncol* **12**: 1436–1442
- Vorobiof DA, Chasen MR, Abratt R, Rapoport B, Cronje N, Fourie L, McMichael G, Hacking D (2000) Taxotere in malignant pleural mesothelioma. A Phase II clinical trial. Program and abstracts of the 9th World Conference on Lung Cancer; September 11–15, Tokyo, Japan. Abstract 58.
- White SC, Anderson H, Jayson GC, Ashcroft L, Ranson M, Thatcher N (2000) Randomised phase II study of cisplatin-etoposide versus infusional carboplatin in advanced non-small-cell lung cancer and mesothelioma. *Ann Oncol* **11**: 201–206
- Zidar BL, Benjamin RS, Frank J, Lane M, Baker LH (1983) Combination chemotherapy for advanced sarcomas of bone and mesothelioma utilizing rubidazole and DTIC: a Southwest Oncology Group Study. *Am J Clin Oncol* **6**: 71–74
- Zidar BL, Green S, Pierce HI, Roach RW, Balcerzak SP, Militello L (1988) A phase II evaluation of cisplatin in unresectable diffuse malignant mesothelioma: a Southwest Oncology Group Study. *Invest New Drugs* **6**: 223–226
- Zidar BL, Metch B, Balcerzak SP, Pierce HI, Militello L, Keppen MD, Berenberg JL (1992) A phase II evaluation of ifosfamide and mesna in unresectable diffuse malignant mesothelioma. A Southwest Oncology Group study. *Cancer* **70**: 2547–2551