REFERENCES

- Fukui M, Koyama M, Nakagawa Y, Itoh Y, Nakamura N, Kondo M (2000) Castration and diabetes. *Diabetes Care* **23**: 1032–1033
- Haffner SM, Valdez RA, Mykkanen L, Stern MP, Katz MS (1994) Decreased testosterone and dehydroepiandrosterone sulfate concentrations are associated with increased insulin and glucose concentrations in nondiabetic men. *Metabolism* **43**: 599–603
- Hsing AW, Chua Jr S, Gao YT, Gentschein E, Chang L, Deng J, Stanczyk FZ (2001) Prostate cancer risk and serum levels of insulin and leptin: a population-based study. *J Natl Cancer Inst* **93**: 783–789
- Jansson PA, Eliasson B, Lindmark S, Eriksson JW (2002) Endocrine abnormalities in healthy first-degree relatives of type 2 diabetes patients—potential role of steroid hormones and leptin in the development of insulin resistance. *Eur J Clin Invest* **2002**: 172–178
- Lehrer S, Diamond EJ, Stagger S, Stone NN, Stock RG (2002) Serum insulin level, disease stage, prostate specific antigen (PSA) and Gleason score in prostate cancer. *Br J Cancer* **87**: 726–728
- Marin P, Holmang S, Jonsson L, Sjostrom L, Kvist H, Holm G, Lindstedt G, Bjorntorp P (1992) The effects of testosterone treatment on body composition and metabolism in middle-aged obese men. *Int J Obes Relat Metab Disord* **16**: 991–997
- Nowicki M, Bryc W, Kokot F (2001) Hormonal regulation of appetite and body mass in patients with advanced prostate cancer treated with combined androgen blockade. *J Endocrinol Invest* **24**: 31–36
- Rizza RA (2000) Androgen effect on insulin action and glucose metabolism. *Mayo Clin Proc* **75**(Suppl): S61–S64
- Stattin P, Bylund A, Rinaldi S, Biessy C, Dechaud H, Stenman UH, Egevad L, Riboli E, Hallmans G, Kaaks R (2000) Plasma insulin-like growth factor-I, insulin-like growth factor-binding proteins, and prostate cancer risk: a prospective study. *J Natl Cancer Inst* **92**: 1910–1917
- Thomas T, Burguera B, Melton III LJ, Atkinson EJ, O'Fallon WM, Riggs BL, Khosla S (2000) Relationship of serum leptin levels with body composition and sex steroid and insulin levels in men and women. *Metabolism* **49**: 1278–1284

Reply: Prostate cancer, insulin and androgen deprivation therapy

S Lehrer^{*,1,2}

¹Department of Radiation Oncology, Mount Sinai School of Medicine, New York, USA; ²Veterans Affairs Medical Center, Bronx, New York, NY 10029, USA

British Journal of Cancer (2003) **89**, 1815. doi:10.1038/sj.bjc.6601352 www.bjcancer.com
© 2003 Cancer Research UK

Sir,

Stattin and Kaaks's data show that insulin levels increase with tumour stage, which is what we reported. They do not have a large enough sample size (sufficient statistical power) to conclude that there is no difference in the insulin levels of the groups in the table. I suggest that Stattin and Kaaks collect a larger sample after determining the sample size; they need to provide 90% power of detecting a significant ($P = 0.05$ two-tailed) result. Also, they might

want to measure fasting insulin before breakfast to reduce the variability of the data, since insulin levels fluctuate throughout the day and after meals. In the meantime, I refer Stattin, Kaaks and all interested readers to the article by Altman and Bland (1995), which eloquently explains the pitfalls of concluding clinical insignificance when differences between sets of small sample size are found to be statistically nonsignificant.

REFERENCE

- Altman DG, Bland JM (1995) Absence of evidence is not evidence of absence. *BMJ* **311**: 485

*Correspondence: S Lehrer; E-mail: stevenlehrer@hotmail.com