

Phase I study of pegylated liposomal doxorubicin and the multidrug-resistance modulator, valspodar

PM Fracasso^{*1}, KA Blum¹, MK Ma¹, BR Tan¹, LP Wright¹, SA Goodner¹, CL Fears¹, W Hou¹, MA Arquette¹, J Picus¹, A Denes¹, JE Mortimer¹, L Ratner¹, SP Ivy² and HL McLeod¹

¹Alvin J Siteman Cancer Center and the Department of Medicine, Washington University School of Medicine, St Louis, MO 63110, USA; ²Cancer Therapy Evaluation Program, Division of Cancer Treatment and Diagnosis, National Cancer Institute, Bethesda, MD, USA

Valspodar, a P-glycoprotein modulator, affects pharmacokinetics of doxorubicin when administered in combination, resulting in doxorubicin dose reduction. In animal models, valspodar has minimal interaction with pegylated liposomal doxorubicin (PEG-LD). To determine any pharmacokinetic interaction in humans, we designed a study to determine maximum tolerated dose, dose-limiting toxicity (DLT), and pharmacokinetics of total doxorubicin, in PEG-LD and valspodar combination therapy in patients with advanced malignancies. Patients received PEG-LD 20–25 mg m⁻² intravenously over 1 h for cycle one. In subsequent 2-week cycles, valspodar was administered as 72 h continuous intravenous infusion with PEG-LD beginning at 8 mg m⁻² and escalated in an accelerated titration design to 25 mg m⁻². Pharmacokinetic data were collected with and without valspodar. A total of 14 patients completed at least two cycles of therapy. No DLTs were observed in six patients treated at the highest level of PEG-LD 25 mg m⁻². The most common toxicities were fatigue, nausea, vomiting, mucositis, palmar plantar erythrodysesthesia, diarrhoea, and ataxia. Partial responses were observed in patients with breast and ovarian carcinoma. The mean (range) total doxorubicin clearance decreased from 27 (10–73) ml h⁻¹ m⁻² in cycle 1 to 18 (3–37) ml h⁻¹ m⁻² with the addition of valspodar in cycle 2 ($P=0.009$). Treatment with PEG-LD 25 mg m⁻² in combination with valspodar results in a moderate prolongation of total doxorubicin clearance and half-life but did not increase the toxicity of this agent.

British Journal of Cancer (2005) 93, 46–53. doi:10.1038/sj.bjc.6602653 www.bjcancer.com

Published online 7 June 2005

© 2005 Cancer Research UK

Keywords: doxil; multidrug resistance; pegylated liposomal doxorubicin; PSC 833; valspodar

Resistance to chemotherapeutic agents is an obstacle to the successful treatment of malignancies. Resistance has been attributed, in part, to the expression of the *MDR1* gene and its protein product, P-glycoprotein (Gottesman *et al*, 2002). P-glycoprotein, a member of the ATP-binding cassette superfamily of transmembrane transporters, prevents the intracellular accumulation of many natural product-derived cytotoxic agents (Gottesman *et al*, 2002). As a result, targeted inhibition of P-glycoprotein by agents administered in combination with chemotherapeutic agents has been attempted (Ford *et al*, 1996; Sikic, 1999; Leonard *et al*, 2003). One such agent, valspodar (PSC 833), a cyclosporine D analogue, reverses P-glycoprotein-mediated resistance *in vitro* at concentrations of 1000 ng ml⁻¹ (Twentyman and Bleehen, 1991). In phase I studies, combinations of valspodar with single agent etoposide, doxorubicin, paclitaxel, or vinblastine are feasible with toxicities consisting of reversible cerebellar ataxia, myelosuppression, and hyperbilirubinemia (Boote *et al*, 1996; Giaccone *et al*, 1997; Fracasso *et al*, 2000; Bates *et al*, 2001, 2004; Chico *et al*, 2001; Minami *et al*, 2001). Responses and disease stabilisation have been

noted in patients with carcinomas of the ovary, lung, and kidney (Fracasso *et al*, 2000; Bates *et al*, 2001; Chico *et al*, 2001).

Valspodar has been shown to decrease the clearance of etoposide, doxorubicin, paclitaxel, and vinblastine, and these interactions have necessitated dose reductions of the anticancer agents when given as single agents in combination with valspodar (Boote *et al*, 1996; Giaccone *et al*, 1997; Fracasso *et al*, 2000; Bates *et al*, 2001, 2004; Chico *et al*, 2001; Minami *et al*, 2001). Liposomal encapsulated anticancer agents may be better suited to combination therapy with valspodar therapy, as one such agent, liposomal doxorubicin, appears to have minimal pharmacokinetic interactions with valspodar in mice, in comparison to free doxorubicin (Krishna and Mayer, 1997; Krishna *et al*, 2000). Additional *in vivo* studies have demonstrated that the elimination of liposomal doxorubicin from the plasma via renal and hepatic mechanisms is minimally affected by valspodar (Krishna *et al*, 1999). These animal studies would suggest that the FDA-approved anticancer agent, pegylated liposomal doxorubicin (PEG-LD) (Doxil[®]) may be administered at full dose in combination with valspodar to patients with malignancies, thus allowing for equivalent anticancer drug exposure with the added benefit of P-glycoprotein inhibition and no additional toxicity.

Given the activity of PEG-LD in various malignancies, including Kaposi's sarcoma and breast and ovarian carcinoma (Northfelt *et al*, 1998; Gordon *et al*, 2000; Campos *et al*, 2001; Sparano and Winer, 2001), and its minimal interactions with valspodar in

*Correspondence: Dr PM Fracasso, Department of Medicine, Washington University School of Medicine, 660 South Euclid St, Box 8056, St Louis, MO 63110, USA; E-mail: fracasso@wustl.edu

Received 23 February 2005; revised 28 April 2005; accepted 9 May 2005; published online 7 June 2005

animal studies (Krishna and Mayer, 1997; Krishna *et al*, 1999, 2000), we proposed a phase I study with this combination. This study was designed to determine the maximum tolerated dose (MTD) and dose-limiting toxicity (DLT) using a one-patient-per-cohort accelerated titration design as proposed by Simon *et al* (1997). In addition, this study evaluated the toxicities of the combination of PEG-LD and valspodar and investigated the effects of valspodar on the pharmacokinetics of doxorubicin. Treatment responses were observed, but were not a primary end point.

PATIENTS AND METHODS

Patient eligibility

This study was initially written and approved for treatment of patients with AIDS-related Kaposi's sarcoma. However, given the decline in Kaposi's sarcoma with the use of highly active antiretroviral therapy (HAART) therapy and the known activity of PEG-LD in breast and ovarian malignancies, this protocol was amended to include patients with a histologically documented resistant or recurrent malignancies. A life expectancy > 4 months, age ≥ 18 years, and laboratory parameters (haemoglobin ≥ 8 g dl⁻¹, absolute neutrophil count ≥ 1000 cells μ l⁻¹, platelet count $\geq 75\,000$ cells μ l⁻¹, creatinine ≤ 2.0 mg dl⁻¹, SGOT $\leq 2 \times$ the institutional upper limit of normal (ULN), and bilirubin $< 1.5 \times$ the institutional ULN) were required. Prior cytotoxic or radiation therapy had to be completed 4 weeks prior to enrollment. Prior chemotherapy with PEG-LD and doxorubicin was permitted. Patients with congestive heart failure or neuropathy (motor or sensory), or with a history of prior PEG-LD or cyclosporine A hypersensitivity were excluded. Any patient receiving medications that are metabolised by the cytochrome P450 system including calcium channel blockers, imidazole antifungal agents, anti-convulsants, or macrolide antibiotics had to discontinue these medications 48 h prior to valspodar therapy, during valspodar administration and up to 48 h after the last dose of valspodar in a cycle. The Cancer Therapy Evaluation Program (T97-0073), Division of Cancer Treatment and Diagnosis, National Cancer Institute (NCI) and the Washington University Human Studies Committee approved this protocol. All patients provided their signed informed consent prior to study entry.

Treatment plan

This study used a treatment cycle of 14 days, similar to the treatment schedule of PEG-LD used by Northfelt *et al*, (1998) for AIDS-related Kaposi's sarcoma. In cycle 1, patients in dose levels 1 through 7 received PEG-LD (Doxil[®], Ortho Biotech Products, L.P., Raritan, NJ, USA) alone at 20 mg m⁻² intravenously over 1 h. Patients in dose levels 8 and 9 received PEG-LD at 22 and 25 mg m⁻² during cycle 1, respectively. Beginning with cycle 2, PEG-LD was administered at a starting dose of 8 mg m⁻² in combination with a fixed dose of valspodar and dose escalated in 2 mg m⁻² increments (dose levels 2–8) to a final dose escalation to 24 mg m⁻² (dose level 9). However, the protocol was amended after the treatment of patient #8 to change the final dose level to PEG-LD 25 mg m⁻² (administered every 2 weeks). Valspodar (PSC 833, Amdray[®], Novartis Pharmaceuticals Corporation, East Hanover, NJ, USA) is typically administered prior to the antineoplastic agent. However, given the long half-life of PEG-LD, valspodar was administered immediately after the PEG-LD infusion as a 1.42 mg kg⁻¹ h⁻¹ infusion over 2 h and then reduced to 0.42 mg kg h⁻¹ (10 mg kg d⁻¹) for an additional 70 h. Administered in this way, valspodar maintains levels above 1000 ng ml⁻¹, the concentration necessary to reverse P-glycoprotein-mediated resistance *in vitro*, for 72 h.

Patients were not treated at all dose levels because we utilised the one-patient-per-cohort accelerated titration design (#4) as proposed by Simon *et al*, (1997) described below. Briefly, this design began with the treatment of one patient at the lowest dose level (PEG-LD 8 mg m⁻²). Inpatient double dose escalation (i.e. 4 mg m⁻² vs usual single dose increments of 2 mg m⁻²) occurred at the completion of each cycle unless the patient experienced a DLT or two instances of grade ≥ 2 nonhaematologic toxicity during any cycle. Provided the patient(s) on the previous dose level had no DLT and less than two instances of grade 2 nonhaematologic toxicity during any cycle, the next patient enrolled began therapy at the new escalated dose level. Inpatient and interpatient dose escalation in the absence of the toxicity as defined above was permitted, such that dose escalation could occur simultaneously for two (or more) patients at different dose levels. Once a DLT or two instances of grade ≥ 2 nonhaematologic toxicity occurred, the accelerated escalation phase was stopped, and the design reverted to a modified Fibonacci scheme with three patients enrolled at this dose level. If zero of three patients experienced DLT, dose escalation resumed in single dose increments (2 mg m⁻²). If one of three experienced DLT, three more patients were enrolled at the same dose level. If two of three patients experienced a DLT, the previous dose level was expanded to six patients. The MTD was that dose in which no greater than one of six patients experienced a DLT.

Dose-limiting toxicity was defined during the second cycle only as grade 4 neutropenia or thrombocytopenia of > 7 days duration or a failure to recover the absolute neutrophil count (ANC) to ≥ 1000 cells μ l⁻¹ or the platelet count to $\geq 75\,000$ cells μ l⁻¹ by day 21. Any grade 3 or 4 nonhaematologic toxicity excluding alopecia, nausea, vomiting, fever, anorexia, or mucositis were also considered dose-limiting. For valspodar, grade 3 or 4 cerebellar dysfunction, despite a prior 25% dose reduction in valspodar, was also considered a DLT.

Dose modifications and delays were permitted as follows: an ANC < 1000 cells μ l⁻¹ or a platelet count < 75 000 cells μ l⁻¹ by day 1 of the subsequent cycle necessitated a 1 week treatment delay. PEG-LD doses were decreased by one dose level if the ANC was < 500 cells μ l⁻¹ for > 7 days (grade 4 neutropenia), if the patient experienced febrile neutropenia, or if the ANC failed to recover to ≥ 1000 cells μ l⁻¹ by the next cycle. In the event of grade 3 skin toxicity, palmar-plantar erythrodysesthesia (PPE), or mucositis, a 25% dose reduction occurred in the PEG-LD and treatment was delayed until the toxicity resolved to a grade 1. In the case of grade 3 cerebellar dysfunction, valspodar was reduced by 25%.

Antitumour response was evaluated after the second cycle of therapy (i.e. 4 weeks) and subsequently, after every four cycles (i.e. every 8 weeks). Responses were defined according to the RECIST criteria (Therasse *et al*, 2000) except for the patient with Kaposi's sarcoma, whose response was defined according to the criteria used by the World Health Association (WHO Handbook, 1979).

Pharmacokinetic monitoring

During the first (PEG-LD alone) and the second (PEG-LD and valspodar) cycles, patients had blood samples obtained at sites contralateral to the infusion site and collected in heparinised tubes before PEG-LD infusion, and 1, 2, 6, 8, 24, 48, and 72 h after the start of the PEG-LD infusion. Plasma samples were stored at -70°C until analyses.

Plasma (0.5 ml) with internal standard (daunorubicin 1000 ng) was combined with 3% (v v⁻¹) triton X-100 (50 μ l), vortexed for 10 s to disrupt the liposome, and vortex mixed with 65% (w v⁻¹) 5-sulphosalicylic acid (50 μ l) for 10 s to extract doxorubicin (Chin *et al*, 2001). The reagent mixture (610 μ l) was then centrifuged at 14 000 r.p.m. for 10 min at room temperature. The extracted supernatant was removed and reconstituted in 75 μ l of 3 M sodium

acetate and a final volume of 50 μ l was injected onto a column using a fluorescence detection method. Briefly, a HPLC pump LC-10AD and controller SCL-10A (Shimadzu Scientific instruments, Columbia, MD, USA) were used to deliver a mobile phase of 50 mM potassium dihydrogen phosphate, pH 3.0: acetonitrile (75:25, v v⁻¹) with 10 mg ml⁻¹ desipramine isocratically at a flow rate of 1.0 ml min⁻¹ to precolumn premier C18 (120A) and a Shimadzu chromegabond[®] TC 18 column (4.6 \times 100 mm, 5 μ m) (Shimadzu Scientific Instruments, Columbia, MD, USA). Desipramine was used to reduce doxorubicin adsorption to the HPLC system (Gabizon *et al*, 1993). Samples were injected with a Shimadzu SIL-10AD autosampler. Fluorescence detector RF 10AXL (Shimadzu Scientific Instruments, Columbia, MD, USA) was used to quantitate total doxorubicin concentrations in the presence of the internal standard daunorubicin, using excitation and emission wavelengths of 470 and 590 nm, respectively. Total doxorubicin concentrations were linear within the concentration range of 25–1000 ng ml⁻¹. The lower limit of quantitation for this assay was 25 ng ml⁻¹. Samples containing concentrations above the highest calibration standard were reanalysed after dilution with blank plasma up to a 1:40 dilution. In all, 20 replicates of a low (35 ng ml⁻¹), intermediate (150 ng ml⁻¹), and high (600 ng ml⁻¹) concentration were repeated on three consecutive days to determine the accuracy and precision of the assay. Intra-assay variability was 1.9, 3.3, and 3.8% at 35, 150, and 600 ng ml⁻¹, respectively. Interassay variability was within 5% across the linear range of the calibration curve. All validation samples were within 10% of the corresponding spiked concentrations.

The disposition of total doxorubicin was fit with both one- and two-compartment model with linear distribution and elimination. The parameters estimated in the linear model included apparent volume of the central compartment (V) and the intercompartmental first-order rate constants K_{12} , K_{21} , and K_{10} . Pharmacokinetic parameters for each set of patient data were fit using an iterative two-stage analysis with maximum-likelihood (ML) estimation, as implemented in ADAPT II software (D'Argenio and Schumitzky, 1979), and the median for each parameter was then used as the revised initial estimates for the next ML iteration. The median for each parameter was updated until the parameter estimates for all parameters were stable (defined as no net change to the second significant digit). Total doxorubicin concentration–time data from each patient was fit individually for each cycle. All modelling of cycle 2 data included any residual total doxorubicin concentrations detectable in the pretreatment sample (time = 0). Each data set was assessed for the goodness of model-curve fit by an estimate of the variance for the predicted values, correlation coefficients, and the Akaike information criterion (AIC) between different models.

Area under the concentration vs time curves ($AUC_{0-\infty}$) was calculated by integration of the simulated concentration–time data from the model estimates. Systemic clearance was calculated as $V \times K_{10}$. The alpha and beta elimination half-lives were calculated using equations generated by ADAPT II software. The differences in total doxorubicin clearance, terminal half-life, or total doxorubicin AUC between cycle 1 and 2 were assessed using Wilcoxon matched pairs test. The correlation between PEG-LD dose and AUC was assessed using the Spearman's rank test. All comparisons were considered significant at $P < 0.05$.

RESULTS

Patients

The characteristics of the 14 patients enrolled from March 1998 until December 2001 are summarised in Table 1. Six patients had received prior therapy with doxorubicin and one patient had received prior therapy with PEG-LD. Pegylated liposomal doxo-

Table 1 Patient characteristics

Characteristics	No. of patients
	14
Age	
Mean	54
Range	34–80
Sex	
Male:female	2:12
ECOG performance status	
0	3
1	9
2	2
Diagnosis	
Breast	4
Colorectal	3
Sarcoma (leiomyosarcoma, Kaposi's sarcoma)	2
Other (hepatoma, ovarian, head and neck, primary peritoneal, renal cell carcinoma)	5
Prior therapy	
Chemotherapy	12
Median # of regimens (range)	3 (0–7)
Prior doxorubicin	6
Mean dose (range) in mg m ⁻²	337 (100–555)
Prior PEG-LD	1
Dose (mg m ⁻²)	279
Radiation therapy	7

PEG-LD = pegylated liposomal doxorubicin.

rubicin and valsopodar was first-line therapy for the patients with hepatoma and Kaposi's sarcoma.

Treatment administered

All patients completed at least two cycles of therapy, with cycle 1 containing PEG-LD alone and cycle 2 containing both PEG-LD and valsopodar (Table 2). A total of 69 combined PEG-LD and valsopodar treatment cycles were administered, (range 1–13). As per study design, inpatient double dose escalation was terminated after patient #2 experienced a DLT (grade 3 PPE) at PEG-LD 24 mg m⁻². This patient with hepatoma received a total of 14 cycles after a 25% dose reduction at cycle 5 and again at cycle 10 (see Table 2). The traditional single-step dose escalation with three patients per dose level was then initiated with patient #3 receiving PEG-LD 20 mg m⁻², the previous dose level for which patient #2 had no significant toxicities. Since no DLTs were observed at PEG-LD 20 and 22 mg m⁻², six patients received PEG-LD 25 mg m⁻² in combination with valsopodar.

In all, 90% of the 69 cycles of combined PEG-LD and valsopodar were administered on schedule. Seven cycles administered to four patients (#2, 6, 12, and 14) were delayed due to PPE. These four patients also underwent dose reductions (as noted in Table 2) following the dose delay. In these four patients, all dose delays and reductions occurred after receiving at least three cycles of PEG-LD (one cycle of PEG-LD alone and several cycles in combination with valsopodar) at a dose of >20 mg m⁻².

Toxicity

Table 3 summarises the haematologic toxicities for all treatment cycles. Only two grade 3 haematologic toxicities were noted. Patient #12 experienced grade 3 neutropenia and PPE after four cycles of treatment with PEG-LD 25 mg m⁻², which led to a

Table 2 Summary of all patients and treatment administered

Patient #	Cycle # and dose of PEG-LD (mg m ⁻²)													
	1 ^a	2	3	4	5	6	7	8	9	10	11	12	13	14
1	20	8	12	16	20									
2	20	16	20	24	18	18	18	18	18	13.5	13.5	13.5	13.5	13.5
3	20	20	20	20										
4	20	20												
5	20	20												
6	20	22	22	16.5	16.5	16.5	16.5	12.4	12.4	12.4	12.4	12.4		
7	22	22	22	22										
8	22	22	22	22										
9	25	25	25	25										
10	25	25												
11	25	25												
12	25	25	25	25	18.75	18.75	18.75	18.75	18.75	18.75	18.75	18.75		
13	25	25												
14	25	25	25	25	25	18.75	18.75	18.75	18.75	18.75	14.06	14.06	14.06	14.06

^aAll patients received PEG-LD (pegylated liposomal doxorubicin) alone for the cycle 1 and PEG-LD in combination with valsopodar for all other cycles.

Table 3 Hematologic toxicities

Patient #	PEG-LD (mg m ⁻²)	No. of PEG-LD and valsopodar cycles ^a	ANC nadir mean (range) (cells μl ⁻¹)	Platelet nadir mean (range) (10 ³ μl ⁻¹)	Haemoglobin mean (range) (g dl ⁻¹)
1–14	20–25	0	3.803 (1.994–7.571)	271 (161–424)	11.2 (6.7–13.8)
1	8	4 ^b	2.631 (1.613–3.602)	220 (195–229)	13.5 (12.8–14.1)
2	16	13 ^c	3.106 (2.109–5.083)	399 (295–529)	12.1 (10.8–12.8)
3–5	20	5	3.730 (2.275–6.255)	311 (66–501)	10.2 (8.3–12.5)
6–8	22	17 ^d	2.826 (1.386–5.039)	265 (113–464)	11.0 (8.6–13.8)
9–14	25	30 ^e	2.298 (0.936–7.056)	298 (137–477)	11.6 (8.7–14.0)

^aTotal number of cycles does not include cycle 1 (PEG-LD alone). ^bPatient #1 had dose escalations of 12, 16, and 20 mg m⁻². ^cPatient #2 received dose escalations of 20 and 24 mg m⁻² before dose reductions of 18 mg m⁻² for five cycles and 13.5 mg m⁻² for five cycles. ^dPatient #6 was dose reduced to 16.5 mg m⁻² for four cycles and 12.4 mg m⁻² for five cycles. ^ePatient #12 was dose reduced to 18.75 mg m⁻² for five cycles and 14.06 mg m⁻² for four cycles. Patient #14 was dose reduced to 18.75 mg m⁻² for eight cycles. PEG-LD = pegylated liposomal doxorubicin.

Table 4 Most common nonhaematologic toxicities

Toxicity	Cycle 1 ^a					All other cycles ^b				
	Grade					Grade				
	0	1	2	3	4	0	1	2	3	4
<i>Constitutional</i>										
Fatigue/malaise	10	2	1	1	—	13	1	—	—	—
<i>Dermatology</i>										
PPE	14	—	—	—	—	9	—	—	4	1
<i>Gastrointestinal</i>										
Diarrhoea	10	2	2	—	—	10	2	1	1	—
Nausea/vomiting	9	4	1	—	—	4	6	3	1	—
Stomatitis/dysphagia	14	—	—	—	—	7	5	2	—	—
<i>Infection w/o neutropenia</i>										
Infection w/o neutropenia	10	1	2	1	—	12	1	1	—	—
<i>Other</i>										
PEG-LD infusion reaction	13	—	—	1	—	14	—	—	—	—
Valsopodar-related events	—	—	—	—	—	5	8	—	1	—

^aPEG-LD (pegylated liposomal doxorubicin) alone. ^bPEG-LD given in combination with valsopodar.

subsequent dose delay and reduction, and patient #4 experienced grade 3 anemia requiring a red blood cell transfusion after cycle 1. In addition, patients #5, 6, 8, and 14 with grade 2 anemia received red blood cell transfusions, and patients #4, 5, 13, and 14 received erythropoietin.

Table 4 summarises the most common nonhaematologic toxicities. Grade 3 and 4 events were uncommon and consisted primarily of PPE. Patient #9 developed the only grade 4 toxicity, PPE, after receiving four cycles of PEG-LD 25 mg m⁻². Patients #2, 6, 12, and 14 experienced grade 3 PPE after they received more

than three cycles of therapy at PEG-LD at doses exceeding 20 mg m^{-2} . Other grade 3 events included fatigue, infection without neutropenia (cellulitis), diarrhoea, nausea/vomiting, PEG-LD infusion reaction, and valsopodar-related ataxia. The PEG-LD reaction consisted of flushing, tightness, and abdominal pain. This resolved after administration of dexamethasone and diphenhydramine and did not occur with subsequent cycles after premedication with these agents. During the valsopodar infusion, nine patients developed symptoms consisting of perioral numbness and tingling, paresthesias, chest discomfort, dizziness, light-headedness, or ataxia. These symptoms were mild (grade 1) and resolved shortly after the completion of the infusion except in patient #13 who developed grade 3 ataxia which resolved several days later. This patient was never retreated with PEG-LD and valsopodar due to progressive disease.

Response

Two of the 14 patients achieved a partial response (PR). Patient #6 who had received seven prior chemotherapy regimens for metastatic breast carcinoma achieved a PR lasting for 9 weeks. Patient #12 with ovarian carcinoma who had previously received PEG-LD achieved a partial response for 19 weeks. After 12 cycles of therapy, this patient requested that she be removed from the study, as the infusion pump required for administration of valsopodar was inconvenient for her lifestyle. A patient with Kaposi's sarcoma had a near PR (49% decrease in target skin lesions). This patient had s.d. for 12 weeks and after five cycles of therapy, this patient refused additional therapy. An additional four patients with breast (two patients), hepatoma, primary peritoneal, and rectal carcinoma had stable disease (range 4–22 weeks). Patient #2 with hepatoma received 13 cycles of the combination therapy with a best response of stable disease. This patient was removed from study to undergo curative surgical resection, although, at the time of resection, this was not possible.

Pharmacokinetics

All patients had total doxorubicin pharmacokinetics performed either alone and in combination with valsopodar. Initial evaluation with a one-compartment model was used based on a previous study of PEG-LD disposition in children (Marina *et al*, 2002) and was able to describe total doxorubicin pharmacokinetics in the current study. However, both the AIC score and the goodness of fit were inferior to the two-compartment model (data not shown). Figure 1 depicts a best-fit line for total doxorubicin concentration–time data in patient #1 (PEG-LD 8 mg m^{-2}) during cycles 1 and 2. Maximum-likelihood analysis was updated and two iterations were required to achieve stable estimates of the median parameters. The mean parameters (CV%) observed in the study include $K_{10} = 0.02 \text{ h}^{-1}$ (61%), $V = 1.48 \text{ l m}^{-2}$ (29%), $K_{12} = 0.21$ (140%) h^{-1} , and $K_{21} = 0.64 \text{ h}^{-1}$ (69%). Total doxorubicin pharmacokinetic parameters by patient are listed in Table 5. The interindividual variability in total doxorubicin clearance was seven-fold ($10\text{--}73 \text{ mL h m}^{-2}$) in cycle 1 and over 12-fold ($3\text{--}37 \text{ mL h}^{-1} \text{ m}^{-2}$) in cycle 2. The correlation between dose and total doxorubicin clearance was weak ($R_s = 0.06$). The mean (range) total doxorubicin clearance decreased from 27 ($10\text{--}73$) $\text{mL h}^{-1} \text{ m}^{-2}$ in cycle 1 to 18 ($3\text{--}37$) $\text{mL h}^{-1} \text{ m}^{-2}$ with the addition of valsopodar in cycle 2 ($P = 0.009$). A three-fold range in V was observed in cycle 1 and there was no significant difference in V between cycles 1 and 2 ($P = 0.43$). The estimated mean (range) terminal elimination half-life was 89 h ($40\text{--}336 \text{ h}$) in cycle 2, much longer than estimated in cycle 1 ($P = 0.001$). The AUC was also slightly higher in cycle 2 (mean $1689 \mu\text{g h mL}^{-1}$, range $492\text{--}6257 \mu\text{g h mL}^{-1}$) than in cycle 1 (mean $1087 \mu\text{g h mL}^{-1}$, range $343\text{--}2408 \mu\text{g h mL}^{-1}$; $P = 0.04$). In this small set of patients, neither toxicity nor response was clearly related to total doxorubicin AUC or clearance.




Figure 1 Plasma doxorubicin concentration vs time plot in patient 1 after receiving intravenous doxorubicin at 20 mg m^{-2} over 60 min in cycle 1. A second intravenous doxorubicin dose at 8 mg m^{-2} was started 14 days (336 h) after the first dose in combination with valsopodar at a dosing rate of $1.42 \text{ mg kg}^{-1} \text{ h}^{-1}$ over 2 h, followed by $0.42 \text{ mg kg}^{-1} \text{ h}^{-1}$ over an additional 70 h. Symbols represent measured plasma doxorubicin concentrations, and the solid line represents the best fit from the maximum likelihood estimation using ADAPT II software.

DISCUSSION

Clinical trials in solid tumour malignancies using the P-glycoprotein modulator, valsopodar, combined with single agent etoposide, doxorubicin, paclitaxel, or vinblastine have required dose reductions of the antineoplastic agents (Boote *et al*, 1996; Giaccone *et al*, 1997; Fracasso *et al*, 2000; Bates *et al*, 2001, 2004; Chico *et al*, 2001; Minami *et al*, 2001). These dose reductions are necessary in order to reduce toxicity due to the decreased clearance and increased AUC of these agents when given in combination with valsopodar. A theoretical concern has been that these dose reductions may limit drug exposure to the tumour, thereby decreasing tumour cell kill. Combining valsopodar with PEG-LD may overcome this limitation as animal studies have revealed no pharmacokinetic interactions between these drugs (Krishna and Mayer, 1997; Krishna *et al*, 1999, 2000). In addition, since liposomes reduce the exposure of entrapped anticancer agent to susceptible tissues while increasing the drug delivery to tumours, PEG-LD may provide maximal drug intensity with minimal toxicity (Krishna and Mayer, 1999). In this study, we demonstrated that valsopodar can be given safely with acceptable toxicity at PEG-LD 25 mg m^{-2} every 2 weeks. There was no DLT at this dose level after cycle 2 (PEG-LD and valsopodar). No further dose escalation was attempted as we reasoned that PEG-LD 25 mg m^{-2} every 2 weeks equaled the cumulative dose of 50 mg m^{-2} every 4 weeks, the approved dose of PEG-LD used in the treatment of breast and ovarian carcinoma. More importantly, most patients required dose reductions due to PPE or neutropenia after multiple cycles of this combination at this dose level as well as at lower dose levels. When administered on this schedule, the pharmacokinetics of total doxorubicin in combination with valsopodar was moderately affected.

The pharmacokinetic data collected in this study suggested that valsopodar does moderately increase plasma levels and decrease the clearance of total doxorubicin. Nevertheless, this reduced clearance seemed to have little effect on the toxicity profile of PEG-LD when given at the 25 mg m^{-2} dose. Large interpatient variability (seven-fold in cycle 1 and 12-fold in cycle 2) in total doxorubicin clearance was evident, which was consistent with that found in other adult studies (4–40 fold) (Gabizon *et al*, 1991, 1994;

Table 5 Patient pharmacokinetic parameters

Patient #	PEG-LD (mg m ⁻²)		Clearance (ml h ⁻¹ m ⁻²)		Terminal half-life (h)		AUC _{0-∞} (μg h ml ⁻¹)	
	Cycle 1	Cycle 2	Cycle 1	Cycle 2	Cycle 1	Cycle 2	Cycle 1	Cycle 2
1	20		32		44		622	
	8		17		101		492	
2	20		12		63		1693	
	16		13		88		1301	
3	20		25		44		789	
	20		19		65		1041	
4	20		18		55		1115	
	20		17		65		1175	
5	20		17		48		1165	
	20		19		52		1082	
6	20		26		50		784	
	22		3		336		6257	
7	22		38		40		581	
	22		20		48		1090	
8	22		38		43		579	
	22		32		45		690	
9	25		22		63		1122	
	25		12		102		2470	
10	25		29		48		870	
	25		20		60		1299	
11	25		73		34		343	
	25		37		40		684	
12	25		11		73		2279	
	25		11		83		2348	
13	25		29		48		869	
	25		27		55		945	
14	25		10		99		2408	
	25		10		101		2767	
Mean Cycle 1		1.4	27		54		1087	
CV%		32	60		31		58	
Mean Cycle 2		1.5	18		89		1689	
CV%		27	49		84		88	

PEG-LD = pegylated liposomal doxorubicin.

Amantea *et al*, 1997). One patient (#6) had a dramatic decrease in clearance in cycle 2 with resulting high AUC. The patient was on no other known inhibitors of cytochrome P450 and the mechanism for this observation is not clear. The long half-life is not surprising for a pegylated liposomal drug and increased doxorubicin release was previously demonstrated in preclinical and clinical studies with this composition (Papahadjopoulos *et al*, 1991; Gabizon, 1992; Huang *et al*, 1994; Amantea *et al*, 1997). During the second cycle in which PEG-LD and valsopodar are combined, the total doxorubicin half-life is even longer. This apparent increase in total doxorubicin half-life during second cycle could be in part an artefact of 'carry-over' from cycle 1 rather than purely a pharmacokinetic interaction between valsopodar and doxorubicin. However, three patients (Patients #7, 9, and 11) who had no residual plasma doxorubicin prior to their second cycle, all consistently had 50%

reduction in total doxorubicin clearance in cycle 2 supporting the conclusion that this finding was not an estimation artefact from the residual drug. Therefore, the reduction in total doxorubicin clearance with the addition of valsopodar in cycle 2 was likely a result of a pharmacokinetic interaction, similar to that seen in other models where the clearance of single agent cytotoxic agents such as etoposide, doxorubicin, paclitaxel, and vinblastine was significantly reduced by valsopodar administration (Boote *et al*, 1996; Giaccone *et al*, 1997; Fracasso *et al*, 2000; Bates *et al*, 2001, 2004; Chico *et al*, 2001; Minami *et al*, 2001). An influence of repeated administrations of PEG-LD on this interaction could not be assessed, as all patients received valsopodar in combination with PEG-LD after cycle 1.

Despite this effect on total doxorubicin clearance, valsopodar did not seem to significantly alter the toxicities of PEG-LD. Patients

with the highest total doxorubicin AUC ($>2000 \mu\text{g h ml}^{-1}$) after cycle 2 of therapy had similar toxicities to other patients. Myelosuppression, an uncommon toxicity with PEG-LD, did not appear to be exacerbated by valsopodar. Only one patient experienced grade 3 neutropenia and no episodes of neutropenic fever occurred. As would be expected, dermatologic and gastrointestinal events were more commonly observed. PPE occurred more often in patients receiving greater than three cycles of PEG-LD and valsopodar and often led to dose reductions. As is the case with single agent studies with PEG-LD, this toxicity is more common after cumulative doses of PEG-LD. Nausea and vomiting occurred more frequently in patients after the first cycle when the combination therapy was administered as opposed to the first cycle when only PEG-LD was given as a single agent. This is a known toxicity with valsopodar, although it is possible that the combination exacerbated this toxicity. Nevertheless, the majority of these events were grade 1–2 and controllable with antiemetic therapy.

In conclusion, the combination of PEG-LD 25 mg m^{-2} given with a 72 h infusion of valsopodar can be administered every 2 weeks with acceptable toxicity. The pharmacokinetic interaction is

moderate. Responses were noted in patients with breast and ovarian carcinoma at this dose of PEG-LD. However, the 72 h infusion makes this an inconvenient regimen. In addition, there are more potent third-generation, highly specific, P-glycoprotein modulators in clinical trials. One such trial in Europe, a phase 2 study with docetaxel and the oral agent, zosuquidar, in women with metastatic breast carcinoma, is ongoing. Should this be a positive study, additional studies with PEG-LD and this agent should be considered in the treatment of breast carcinoma.

ACKNOWLEDGEMENTS

We thank the nurses and data managers at the Siteman Cancer Center for caring for the patients on this trial and Dr Teresa J Vietti for her editorial assistance. This work was supported by the St Louis Men's Group Against Cancer and a Clinical Oncology Career Development Award CDA-96-27 from the American Cancer Society (Atlanta, GA) (to PMF) and P30 CA091842 (Pharmacology Core of the Siteman Cancer Center).

REFERENCES

- Amantea MA, Forrest A, Northfelt DW, Mamelok R (1997) Population pharmacokinetics and pharmacodynamics of pegylated-liposomal doxorubicin in patients with AIDS-related Kaposi's sarcoma. *Clin Pharmacol Ther* **61**: 301–311
- Bates S, Bakke S, Kang M, Robey R, Zhai S, Thambi P, Chen C, Patil S, Smith T, Steinberg S, Merino M, Goldspiel B, Meadows B, Stein W, Choyke P, Balis F, Figg W, Fojo T (2004) A phase I/II study of infusional vinblastine with the P-glycoprotein antagonist valsopodar (PSC 833) in renal cell carcinoma. *Clin Cancer Res* **10**: 4724–4733
- Bates S, Kang M, Meadows B, Bakke S, Choyke P, Merino M, Goldspiel B, Chico I, Smith T, Chen C, Robey R, Bergan R, Figg WD, Fojo T (2001) A phase I study of infusional vinblastine in combination with the P-glycoprotein antagonist PSC 833 (valsopodar). *Cancer* **92**: 1577–1590
- Boote DJ, Dennis IF, Twentyman PR, Osborne RJ, Laburte C, Hensel S, Smyth JF, Brampton MH, Bleehen NM (1996) Phase I study of etoposide with SDZ PSC 833 as a modulator of multidrug resistance in patients with cancer. *J Clin Oncol* **14**: 610–618
- Campos SM, Penson RT, Mays AR, Berkowitz RS, Fuller AF, Goodman A, Matulonis UA, Muzikansky A, Seiden MV (2001) The clinical utility of liposomal doxorubicin in recurrent ovarian cancer. *Gynecol Oncol* **81**: 206–212
- Chico I, Kang MH, Bergan R, Abraham J, Bakke S, Meadows B, Rutt A, Robey R, Choyke P, Merino M, Goldspiel B, Smith T, Steinberg S, Figg WD, Fojo T, Bates S (2001) Phase I study of infusional paclitaxel in combination with the P-glycoprotein antagonist PSC 833. *J Clin Oncol* **19**: 832–842
- Chin DL, Lum BL, Duran GE, Sikic BI (2001) A rapid and sensitive HPLC assay of pegylated (polyethylene glycol) liposomal doxorubicin by UV detection. *Proc Am Assoc Cancer Res* **42**: 383
- D'Argenio DZ, Schumitzky A (1979) A program package for simulation and parameter estimation in pharmacokinetic systems. *Comput Programs Biomed* **9**: 115–134
- Ford J, Yang J-M, Hait W (1996) P-glycoprotein-mediated multidrug resistance: experimental and clinical strategies for its reversal. In *Drug Resistance*, Hait W (ed) pp 3–38. Norwell: Kluwer Academic Publishers
- Fracasso PM, Westerveldt P, Fears CA, Rosen DM, Zuhowski EG, Cazenave LA, Litchman M, Egorin MJ (2000) Phase I study of paclitaxel in combination with multidrug resistance modulator, PSC 833 (valsopodar), in refractory malignancies. *J Clin Oncol* **18**: 1124–1134
- Gabizon A (1992) Selective tumor localization and improved therapeutic index of anthracyclines encapsulated in long-circulating liposomes. *Cancer Res* **52**: 891–896
- Gabizon A, Catane R, Uziely B, Kaufman B, Safra T, Cohen R, Martin F, Huang A, Barenholz Y (1994) Prolonged circulation time and enhanced accumulation in malignant exudates of doxorubicin encapsulated in polyethylene-glycol coated liposomes. *Cancer Res* **54**: 987–992
- Gabizon A, Chisin R, Amselem S, Druckmann S, Cohen R, Goren D, Fromer I, Peretz T, Sulkes A, Barenholz Y (1991) Pharmacokinetic and imaging studies in patients receiving a formulation of liposome-associated adriamycin. *Br J Cancer* **64**: 1125–1132
- Gabizon AA, Barenholz Y, Bialer M (1993) Prolongation of the circulation time of doxorubicin encapsulated in liposomes containing a polyethylene glycol-derivatized phospholipid: pharmacokinetic studies in rodents and dogs. *Pharmacol Res* **10**: 703–708
- Giaccone G, Linn SC, Welink J, Catimel G, Stieltjes H, van der Vijgh WJ, Eelink C, Vermorken JB, Pinedo HMA (1997) A dose-finding and pharmacokinetic study of reversal of multidrug resistance with SDZ PSC 833 in combination with doxorubicin in patients with solid tumors. *Clin Cancer Res* **3**: 2005–2015
- Gordon AN, Granai CO, Rose PG, Hainsworth J, Lopez A, Weissman C, Rosales R, Sharpington T (2000) Phase II study of liposomal doxorubicin in platinum- and paclitaxel-refractory epithelial ovarian cancer. *J Clin Oncol* **18**: 3093–3100
- Gottesman M, Fojo T, Bates S (2002) Multidrug resistance in cancer: role of ATP-dependent transporters. *Nat Rev Cancer* **2**: 48–58
- Huang S, Stauffer P, Hong K, Guo J, Phillips T, Huang A, Papahadjopoulos D (1994) Liposomes and hyperthermia in mice: increased tumor uptake and therapeutic efficacy of doxorubicin in sterically stabilized liposomes. *Cancer Res* **54**: 2186–2191
- Krishna R, Mayer LD (1997) Liposomal doxorubicin circumvents PSC 833-free drug interactions, resulting in effective therapy of multidrug-resistant solid tumors. *Cancer Res* **57**: 5246–5253
- Krishna R, Mayer LD (1999) The use of liposomal anticancer agents to determine the roles of drug pharmacodistribution and P-glycoprotein (PGP) blockage in overcoming multidrug resistance (MDR). *Anticancer Res* **19**: 2885–2892
- Krishna R, McIntosh N, Riggs KW, Mayer LD (1999) Doxorubicin encapsulated in sterically stabilized liposomes exhibits renal and biliary clearance properties that are independent of valsopodar (PSC 833) under conditions that significantly inhibit nonencapsulated drug excretion. *Clin Cancer Res* **5**: 2939–2947
- Krishna R, St-Louis M, Mayer L (2000) Increased intracellular drug accumulation and complete chemosensitization achieved in multidrug-resistant solid tumors by co-administering valsopodar (PSC 833) with sterically stabilized liposomal doxorubicin. *Int J Cancer* **85**: 131–141
- Leonard G, Fojo T, Bates S (2003) The role of ABC transporters in clinical practice. *Oncologist* **8**: 411–424
- Marina NM, Cochrane D, Harney E, Zomorodi K, Blaney S, Winick N, Bernstein M, Link MP (2002) Dose escalation and pharmacokinetics of pegylated liposomal doxorubicin (Doxil) in children with solid tumors: a Pediatric Oncology Group study. *Clin Cancer Res* **8**: 413–418

- Minami H, Ohtsu T, Fujii H, Igarashi T, Kuniaki I, Uchiyama-Kokubu N, Aizawa T, Watanabe T, Uda Y, Tanigawara Y, Sasaki Y (2001) Phase I study of intravenous PSC-833 and doxorubicin: reversal of multidrug resistance. *Jpn J Clin Oncol* **92**: 220–230
- Northfelt D, Dezube B, Thommes J, Miller B, Fischl M, Friedman-Kien A, Kaplan L, Du Mond C, Mamelok R, Henry D (1998) Pegylated-liposomal doxorubicin versus doxorubicin, bleomycin, and vincristine in the treatment of AIDS-related Kaposi's sarcoma: results of a randomized phase III clinical trial. *J Clin Oncol* **16**: 2445–2451
- Papahadjopoulos D, Allen T, Gabizon A, Mayhew E, Matthay K, Huang S, Lee K, Woodle M, Lasic D, Redemann C, Martin F (1991) Sterically stabilized liposomes: improvements in pharmacokinetics and antitumor therapeutic efficacy. *Proc Natl Acad Sci* **88**: 11460–11464
- Sikic BI (1999) Modulation of multidrug resistance: a paradigm for translational clinical research. *Oncology (Huntington)* **13**: 183–187
- Simon R, Freidlin B, Rubinstein L, Arbuck S, Collins J, Christian M (1997) Accelerated titration designs for phase I clinical trials in oncology. *J Natl Cancer Inst* **89**: 1138–1147
- Sparano J, Winer E (2001) Liposomal anthracyclines for breast cancer. *Semin Oncol* **28**: 32–40
- Therasse P, Arbuck SG, Eisenhauer EA, Wanders J, Kaplan RS, Rubinstein L, Verweij J, Van Glabbeke M, van Oosterom AT, Christian MC, Gwyther SG (2000) New guidelines to evaluate the response to treatment in solid tumors. *J Natl Cancer Inst* **92**: 205–216
- Twentyman P, Bleehen N (1991) Resistance modification by PSC 833, a novel non-immunosuppressive cyclosporin. *Eur J Cancer* **27**: 1639–1642