EGFR mutation status in tumour-derived DNA from pleural effusion fluid is a practical basis for predicting the response to gefitinib

H Kimura^{1,2,6}, Y Fujiwara^{3,6}, T Sone², H Kunitoh³, T Tamura³, K Kasahara² and K Nishio^{*,1,4,5}

¹Shien-Lab, National Cancer Center Hospital, Tsukiji 5-1, Chuo-ku, Tokyo, Japan; ²Respiratory Medicine, Kanazawa University Hospital, Takara-machi 13-1, Kanazawa, Ishikawa, Japan; ³Medical Oncology, National Cancer Center Hospital, Tsukiji 5-1, Chuo-ku, Tokyo, Japan; ⁴Pharmacology Division, National Cancer Center Research Institute, Tsukiji 5-1-1, Chuo-ku, Tokyo, Japan; ⁵Center for Medical Genomics, National Cancer Center Research Institute, Tsukiji 5-1-1, Chuo-ku, Tokyo, Japan

Epidermal growth factor receptor (EGFR) mutations are strong determinants of tumour response to EGFR tyrosine kinase inhibitors in non-small-cell lung cancer (NSCLC). Pleural effusion is a common complication of lung cancer. In this study, we assessed the feasibility of detection of EGFR mutations in samples of pleural effusion fluid. We obtained 43 samples, which was the cell-free supernatant of pleural fluid, from Japanese NSCLC patients, and examined them for EGFR mutations. The epidermal growth factor receptor mutation status was determined by a direct sequencing method (exons 18–21 in EGFR). EGFR mutations were detected in 11 cases (E746_A750del in seven cases, E746_T751del insA in one case, L747_T751del in one case, and L858R in two cases). The EGFR mutations were observed more frequently in women and non-smokers. A comparison between the EGFR mutant status and the response to gefitinib in the 27 patients who received gefitinib revealed that all seven patients with partial response and one of the seven patients with stable disease had an EGFR mutation. No EGFR mutations were detected in the patients with progressive disease. The results suggest that DNA in pleural effusion fluid can be used to detect EGFR mutations and that the EGFR mutation status may be useful as a predictor of the response to gefitinib.

British Journal of Cancer (2006) 95, 1390–1395. doi:10.1038/sj.bjc.6603428 www.bjcancer.com

Published online 24 October 2006

© 2006 Cancer Research UK

Keywords: pleural effusion; EGFR; mutation; gefitinib

Lung cancer is a major cause of cancer-related mortality worldwide and is expected to remain a major health problem for the foreseeable future (Parkin *et al*, 2005). Most patients have advanced disease at the time of diagnosis. The initial therapy for advanced non-small-cell lung cancer (NSCLC) is systemic chemotherapy with a two-drug combination regimen, which often includes a platinum agent, but the median survival of patients treated with such regimens has ranged from only 8 to 10 months. Little improvement in the efficacy of chemotherapy has been made in the last 20 years (Breathnach *et al*, 2001; Kelly *et al*, 2001; Schiller *et al*, 2002).

Targeting epidermal growth factor receptor (EGFR) is an appealing strategy for the treatment of NSCLC, because EGFR has been found to be expressed, sometimes strongly, in NSCLC (Franklin *et al*, 2002). Gefitinib ('Iressa', AstraZeneca) is a small molecule and selective EGFR tyrosine kinase inhibitor that has shown antitumour activity in NSCLC patients as a single agent in phase II trials (Fukuoka *et al*, 2003). Adding gefitinib to chemotherapy in phase III studies of patients with untreated advanced NSCLC did not significantly improve the outcome over chemotherapy alone (Giaccone *et al*, 2004; Herbst *et al*, 2004), and a possible explanation for the failure to observe any added benefit in these trials is that the patients had not been screened or selected for their ability to derive any clinical benefit from an EGFR inhibitor.

An association between mutations in sites of EGFR tyrosine kinase in NSCLC and hyper-responsiveness to gefitinib has recently been reported (Lynch *et al*, 2004; Paez *et al*, 2004). The mutations consisted of small in-frame deletions or substitutions clustered around the ATP-binding site in exons 18, 19, and 21 of EGFR, and the mutations increased the affinity of the enzyme for ATP and gefitinib (Lynch *et al*, 2004). Some investigators subsequently found that EGFR mutations are one of the strong determinants of tumour response to EGFR tyrosine kinase inhibitors (Pao *et al*, 2004; Han *et al*, 2005). The investigators

*Correspondence: Dr K Nishio, Shien-Lab, National Cancer Center Hospital, Tsukiji 5-1, Chuo-ku, Tokyo, Japan;
E-mail: knishio@gan2.res.ncc.go.jp

Research support: H Kimura received support as an Awardee of a Research Resident Fellowship from the Foundation for Promotion of Cancer Research (Japan) for the 3rd Term Comprehensive 10-Year Strategy for Cancer Control.

⁶These authors contributed equally to this work

Received 13 June 2006; revised 14 September 2006; accepted 15 September 2006; published online 24 October 2006

used surgical tissue to detect the *EGFR* mutations in their studies, but most patients who require gefitinib therapy are diagnosed at an advanced stage of the disease and are inoperable. As it is often difficult to obtain a sufficient tumour sample from patients with inoperable NSCLC to detect *EGFR* mutations by direct sequencing, a method of detecting *EGFR* mutations in other specimens needed to be established.

Malignant pleural effusion is a common complication of lung cancer. It is present in approximately 15% of patients at the time of diagnosis (Pass *et al*, 2005) and in 10–50% of patients during the course of the disease (Fenton and David Richardson, 1995). In about half of NSCLC patients with a pleural effusion, the effusion fluid is cytologically positive at the first time examined, and ultimately most effusions are determined to be malignant. As pleural effusion fluid sampling is usually easy, non-invasive, and repeatable, we hypothesised that tumour-derived DNA in the pleural effusion fluid of NSCLC patients would be a source of useful information on the status of the *EGFR* gene and could allow prediction of the response to gefitinib. Some investigators have reported that pleural effusion fluid is a useful clinical specimen for searching for point mutations in oncogenes, such as *K-ras*, *rho A*, *p53*, and *FHIT* (Nakamoto *et al*, 2001; Lee *et al*, 2004). As the two trials were small, the results regarding the sensitivity and specificity of detection of the mutations in pleural effusion as a diagnostic method were unclear. Detection of *EGFR* mutations in pleural effusion fluid has been described in one case report, and the patient responded to gefitinib (Huang *et al*, 2005). The results in that patient encouraged us to hypothesise that the *EGFR* mutation status determined in pleural effusion fluid is useful for predicting the responsiveness to *EGFR* tyrosine kinase inhibitors.

In the present study, we attempted to detect *EGFR* mutations in pleural effusion fluid and to clarify the usefulness of their detection as a predictor of the response to gefitinib.

PATIENTS AND METHODS

Patients

The subjects were NSCLC patients who had a pleural effusion at the time of diagnosis. The diagnosis of NSCLC was based on the histological or cytological findings, and the histological type was determined according to the WHO criteria (Travis *et al*, 1999). Patients' records consisted of age, gender, smoking habit, histological type, and treatment. Smoking status was collected from the patients' records. Patients were divided into three groups according to their smoking status: never smokers (<100 cigarettes/lifetime), former smokers (≥ 100 cigarettes/lifetime, no smoking at present), and current smokers (≥ 100 cigarettes/lifetime). The response of the patients treated with gefitinib was evaluated every 4 or 8 weeks in accordance with the 'Response Evaluation Criteria in Solid Tumours (RECIST)' guidelines. (Therasse *et al*, 2000). Partial response (PR) and stable disease (SD) were confirmed by a sustained 4-week follow-up. This study was approved by the Institutional Review Board of the National Cancer Center Hospital and of Kanazawa University Hospital, and written informed consent was obtained from all participants. No research results were entered into the patient's records or released to the patient or the patient's physician.

Collection of pleural effusion fluid and DNA purification

The pleural effusion fluid was collected into heparinised tubes between 29 March 2005 and 30 January 2006. No particular collection method was used. A 2-ml sample of the fluid was centrifuged at 250 g for 10 min at room temperature, and the

supernatant was collected and stored at -80°C until DNA extraction. DNA was extracted from 1 ml of the supernatant with a Qiaamp DNA Mini Kit (Qiagen, Hilden, Germany) according to the blood and body fluid spin protocol in the manufacturer's instructions, with the following protocol modifications. The same column was used repeatedly until the whole sample had been processed. The DNA obtained was eluted in 50 μl of sterile bi-distilled buffer, and the extracted DNA was stored at -20°C until used. The amounts of DNA extracted were estimated with spectrophotometry.

Polymerase chain reaction amplification and direct sequencing

Exons 18, 19, 20, and 21 of the *EGFR* gene were amplified by polymerase chain reaction (PCR). The primers were designed based on the report by Lynch *et al* (2004). Genomic PCR of 1 μl of template DNA was performed in 25 μl volumes containing 0.75 U of Ampli Taq Gold DNA polymerase (Perkin-Elmer, Roche Molecular Systems Inc., Branchburg, NJ, USA), 2.5 μl of PCR buffer, 0.8 μM dNTP, 0.5 μM of each primer, and different concentrations of MgCl_2 , depending on the polymorphic marker. The first PCR analyses were performed in a volume of 25 μl by 25 cycles consisting of a denaturation step at 94°C for 45 s, a primer annealing step at 58°C for 30 s, and an elongation step at 72°C for 30 s. The final step at 72°C was extended for 10 min. Nested PCR was performed with 20 cycles under the same conditions as the first PCR. Sequencing of each sample was performed in duplicate with an ABI prism 310 (Applied Biosystems, Foster City, CA, USA). PCR products were sequenced in both sense and antisense directions. *Epidermal growth factor receptor* mutations detected in the initial round of sequencing were confirmed by subsequent rounds of independent PCR and sequencing reactions. Only specimens in which a mutation was identified in both rounds were recorded as mutation-positive. The sequences were compared with the GenBank-archived human sequence for *EGFR* (accession number: AY588246). The nucleic acid and protein coordinates used to name the mutations are based on NM_005228.3 and NP_005219.2, respectively.

Statistical analyses

This study was carried out as exploratory research for detecting *EGFR* mutations from pleural effusion fluid and clarifying the relationship between the mutation status and clinical manifestations. The number of enrolled patients was therefore not precalculated. Patient characteristics, including gender, tumour histology, and smoking habit were tabulated according to their mutation status. Fisher's exact test was used to test for associations between the presence of *EGFR* mutations and the patients' characteristics. The relationship between response to gefitinib and the mutation status was evaluated individually.

RESULTS

Patients and pleural effusion specimens

Forty-three patients were enrolled in this study (Table 1). Two hundred and sixty-two patients were seen with stage IIIB and IV at our institutions in the period of this study. Forty-three of the 262 patients were enrolled in this study. The enrolled patients were not all of the patients with pleural effusion because written informed consent was not obtained from any patients with pleural effusion. Their median age was 62 years (range, 39–82 years), and there were 21 females (53.8%) and 17 never smokers (43.6%). The histological and/or cytological diagnosis was adenocarcinoma in 39 patients, and squamous cell

Table 1 Patient characteristics and *EGFR* mutation status

	(n)	EGFR mutation (n)
No. of patients	43	11 (25.6%)
Age (years)		
Median	63	
Range	39–82	
Gender		
Male	22	4 (18.2%)
Female	21	7 (33.3%)
Smoking habit		
Current	9	2 (22.2%)
Former	16	2 (12.5%)
Never	18	7 (38.9%)
Histology		
Adenocarcinoma	39	11 (28.2%)
Squamous cell carcinoma	1	0 (0%)
Large cell carcinoma	1	0 (0%)
Unclassified	2	0 (0%)
No. of patients treated with gefitinib	27	8 (29.6%)
PR	7	7 (14.3%)
SD	7	1 (0%)
PD	13	0 (0%)

EGFR = epidermal growth factor receptor; PD = progressive disease; PR = partial response; SD = stable disease.

carcinoma and large cell carcinoma in one each, and unclassified NSCLC in two patients. Non-small-cell lung cancer cells in the pleural effusion samples of 40 of the patients were identified cytologically. There were no malignant cells in the pleural effusion fluid of the other three patients. We have no data of the proportion of malignant cells and normal cells. Twenty-seven patients were treated with gefitinib (250 mg day⁻¹) and evaluated for a response. Eight of the 27 patients were treated with gefitinib as an initial treatment and the other 19 patients were treated with the agent as a second or third line. The others were treated with systematic chemotherapy, including a platinum agent. The results of the evaluation showed that seven of the 27 patients who received gefitinib therapy had a PR and seven had SD. The other 13 patients had progressive disease (PD). DNA was extracted from all 43 samples of pleural effusion fluid. Amounts of the DNA extracted were detectable from 27 samples at a concentration up to 144.0 ng ml⁻¹. Amounts from 16 samples were under the detectable limit.

Detection of *EGFR* mutations in pleural effusion fluid

Direct sequencing of PCR products in exons 18–21 of *EGFR* in the pleural effusion fluid of all patients allowed their mutation status to be determined. Heterozygous mutations were identified in 11 (25.6%) of the 43 patients (Table 1). Nine mutations were deletional mutations located in exon 19 (E746_A750del in seven, L746_T751del insA in one, L747_T751del in one), and two were substitution mutations located in exon 21 (L858R) (Table 2) (Figure 1). No mutations were detected in exon 18 or 20. The E746_A750 deletion and L858R substitution mutations were the most common (9 out of 11 mutations, 81.8%) and are well-known hotspot mutations described previously (Kosaka *et al*, 2004; Pao *et al*, 2004). No more than one mutation was identified per patient, and no *EGFR* mutations were detected in pleural effusion fluid that did not contain malignant cells.

Table 2 Site of mutations in exons 18–21 of *EGFR*

Nucleotide changes	Amino-acid changes	No. of patients
2481_2495del	E746_A750del	6
2482_2496del	E746_A750del	1
2483_2497del	E746_T753del insA	1
2486_2500del	L747_T751del	1
2819T > G	L858R	2

EGFR = epidermal growth factor receptor; del = deletion; ins = insertion. The numbering of the mutation sites was based on NM_005228.3 (nucleotide) and NP_005219.2 (amino acid).

Epidermal growth factor receptor mutation status and patients' characteristics

EGFR mutations were detected more frequently in the samples from females (7 out of 21, 33.3% of females, 4 out of 18, 22.2% of males; $P=0.310$) and non-smokers (7 out of 17, 41.1% of non-smokers, 4 out of 22, 18.1% of current or former smokers; $P=0.156$), although the differences were not statistically significant (Table 3). Of the 11 mutations, 63.6% were in women and 63.6% were in non-smokers. All of the patients with mutations had adenocarcinoma. No *EGFR* mutations were found in any of the patients with squamous carcinoma or large cell carcinoma. A comparison between the *EGFR* mutant status and the response to gefitinib showed that all seven patients with a PR and one of the seven patients with SD had an *EGFR* mutation. No *EGFR* mutations were detected in any of the patients with PD (Table 4). We have no response data from the 16 patients who had never treated with gefitinib, and we have not evaluated the relationship between the response to chemotherapy and the *EGFR* mutation status in pleural effusion fluid.

DISCUSSION

This is the first report of an analysis of the *EGFR* mutation status in DNA obtained from the pleural effusion fluid of a series of NSCLC patients and evaluation of the relationship between the mutation status and the clinical response to gefitinib. It is interesting that all patients who achieved a PR to gefitinib had the *EGFR* mutations. We hypothesised that the mutation status in DNA extracted from pleural effusion fluid would allow prediction of the clinical outcome of gefitinib therapy in NSCLC patients, and we therefore expected the pleural effusion fluid to be a practical source of DNA for detection of *EGFR* mutations. The sites of *EGFR* mutations found in this study are identical to those reported in previous studies (Kosaka *et al*, 2004; Pao *et al*, 2004). The main mutations found were in-frame deletions in exon 19 and the missense mutation L858R in exon 21. No patients had more than one mutation. It was possible to determine the mutation status of *EGFR* by using the DNA in only 1.0 ml of pleural effusion fluid, even though the concentration of the extracted DNA specimens was in most cases below the concentration detectable by spectrophotometry (data not shown). The results of the comparison between the mutation status and clinical manifestations in this study confirmed the finding in previous studies that *EGFR* mutations are frequently present in small sub-groups of NSCLC patients, such as females and never smokers, although the differences were not statistically significant. It is well known that *EGFR* mutations are frequently observed in adenocarcinomas. As 36 of the 39 patients (92.3%) enrolled in this study had adenocarcinoma, we could not evaluate differences in the frequency of the *EGFR* mutations according to the histological type. Pleural effusion occurs in lung carcinoma of all cell types, but

Figure 1 The wave figures of the nucleotide sequence of the *EGFR* gene with heterozygous mutations obtained by direct sequencing (see 'Patients and Methods') are shown. Horizontal arrows in both the sense and the antisense directions are shown to demonstrate the two breakpoints of the deletion. The patients in **A**, **B**, and **C** have in-frame deletions in exon 19 (Figure **A**, E746_A750del; **B**, E746_T753del insA; **C**, L747_T751del; **D**, L858R). The double peaks (vertical arrows) represent the heterozygous missense mutations resulting in an amino acid substitution of L858R in exon 19 (Figure **D**).

appears to be most frequent in adenocarcinoma (Chernow and Shahn, 1997).

This study had several limitations. First, we could not compare the results of the *EGFR* mutation status in the pleural effusion fluid to the mutation status in tumour tissue. Forty of the 43 patients

enrolled were cytologically diagnosed as having NSCLC from pleural effusion fluid specimens. As the DNA extracted from pleural effusion fluid consisted of DNA derived from both tumour cells and normal cells, the *EGFR* mutation status needs to be evaluated in a pair of DNA specimens from the tumour and pleural

Table 3 Frequency of *EGFR* mutations in DNA from the pleural effusion fluid of NSCLC patients according to (A) gender, (B) histology, (C) smoking habit, and (D) response to gefitinib

(A) Gender and <i>EGFR</i> mutation status		
	<i>EGFR</i> mutation	
	+	–
Female	7	14
Male	4	18
$P = 0.310$		
(B) Histology and <i>EGFR</i> mutation status		
	<i>EGFR</i> mutation	
	+	–
Ad	11	28
Non-Ad	0	4
$P = 0.558$		
(C) Smoking habit and <i>EGFR</i> mutation status		
	<i>EGFR</i> mutation	
	+	–
Never	7	11
Current/former	4	21
$P = 0.156$		

Ad = adenocarcinoma; EGFR = epidermal growth factor receptor; + = mutation-positive; – = no mutations. (A)(B)(C); a total of 43 samples were evaluated.

effusion fluid to confirm the usefulness of the mutation status determined from pleural effusion fluid. However, it is sometimes difficult to obtain tumour samples from patients with advanced NSCLC, and even more so from patients diagnosed as having NSCLC using methods other than the histological examination of tumour tissue, such as on the basis of pleural effusion or sputum cytology. Second, direct sequencing may be not able to provide satisfactory results for detection of *EGFR* mutations in mixed samples of mutated and wild DNA. Although direct sequencing has generally been used to detect *EGFR* mutations in previous studies, detection of a mutation by this method requires at least 30% of the mutated DNA in a sample (Bosari *et al*, 1995; Fan *et al*, 2001). Lung cancers are very heterogeneous, and as normal cells, such as inflammatory cells or mesothelial cells, are contained in the pleural effusion fluid of lung cancer patients, in addition to tumour cells, a small amount of mutated DNA in pleural effusion fluid can be missed by direct sequencing. Unfortunately, we have no data at the present time on whether *EGFR* mutations were detectable in pleural effusion samples with either a few malignant cells, a small proportion of malignant cells with normal mesothelial cells, or cytologically negative samples. To establish a method for the detection of *EGFR* mutations from pleural effusion fluid, the mutation detectable proportion of malignant cells to normal cells in pleural fluid should be elucidated. We are planning an additional study using cytological examination to clarify the mutation detectable proportion as a next step. When pleural fluid is used as the material for detection of *EGFR* mutations, a patient with an *EGFR* mutation may be diagnosed as having wild-type *EGFR* because of the two limitations described above. Although we expected a high frequency of detection of *EGFR* mutations in this study because of the high proportion of adenocarcinomas (92.3%), we detected *EGFR* mutations in only 28.2% of the patients enrolled, a lower frequency than in two previous reports on Japanese NSCLC patients (Takano *et al*, 2005; Asano *et al*, 2006). Patients with false-negative results, meaning that no *EGFR* mutations were detected in a patient with an *EGFR* mutation, were not excluded from this study. Some investigators have tried to improve the sensitivity of detection of

Table 4 *EGFR* mutation status in patients who received gefitinib therapy

Age (years)	Gender	Smoking	Histology	<i>EGFR</i> mutation status	Response to gefitinib
62	F	Never	Ad	E747_P753insS	PR
58	F	Never	Ad	E746_A750del	PR
80	F	Never	Ad	E746_A750del	PR
61	M	Never	Ad	E746_A750del	PR
65	M	Former	Ad	E746_A750del	PR
60	M	Current	Ad	E746_A750del	PR
66	F	Never	Ad	E747_T750del	PR
76	F	Never	Ad	Wild	SD
57	F	Former	Ad	Wild	SD
40	F	Never	Ad	Wild	SD
72	F	Never	Ad	Wild	SD
58	F	Former	Ad	Wild	SD
66	F	Never	Ad	Wild	SD
65	F	Former	Ad	L858R	SD
39	F	Never	Ad	Wild	PD
69	M	Former	Ad	Wild	PD
72	F	Never	Ad	Wild	PD
74	F	Never	Ad	Wild	PD
67	M	Former	Ad	Wild	PD
62	M	Former	SCC	Wild	PD
59	F	Current	Ad	Wild	PD
77	M	Current	Ad	Wild	PD
82	F	Never	Ad	Wild	PD
66	F	Never	Ad	Wild	PD
56	M	Current	Ad	Wild	PD
61	M	Former	Ad	Wild	PD
65	M	Former	Ad	Wild	PD

Ad = adenocarcinoma; EGFR = epidermal growth factor receptor; F = female; M = male; NSCLC = unclassified NSCLC; PD = progressive disease; PR = partial response; SCC = squamous cell carcinoma; SD = stable disease.

EGFR mutations in samples containing a mixture of tumour and normal cells. Wookey *et al* (2005) reported findings that the ARMS method was superior to the direct sequencing method and WAVE method for detecting *EGFR* mutations. Other groups have reported that LightCycler PCR assay (Sasaki *et al*, 2005), SSCP assay (Marchetti *et al*, 2005), and enriched PCR assay (Asano *et al*) are more sensitive than direct sequencing and are more rapid. A standardised method of detecting *EGFR* mutations needs to be established as soon as possible.

The final limitation in the present study is that it remains unclear whether there is any survival benefits associated with gefitinib therapy in those patients enrolled with *EGFR* mutations. The relationship between the *EGFR* mutation status determined in pleural effusion fluid and the gefitinib response in a portion of the patients enrolled supports the pleural effusion fluid *EGFR* mutation status as useful for predicting the response to gefitinib. The relationship between the *EGFR* mutation status determined in the pleural effusion fluid and the gefitinib response in the remaining patients and the survival benefit of gefitinib therapy in the patients with *EGFR* mutations are currently being evaluated, and confirmation of the results is expected in the very near future.

In conclusion, our results suggest that the DNA in pleural effusion fluid can be used to detect *EGFR* mutations and that the *EGFR* mutation status determined may be useful as a predictive factor of response to gefitinib.

ACKNOWLEDGEMENTS

We gratefully acknowledge the excellent statistical assistance of Tetsutaro Hamano.

REFERENCES

- Asano H, Toyooka S, Tokumo M, Ichimura K, Aoe K, Ito S, Tsukuda K, Ouchida M, Aoe M, Katayama H, Hiraki A, Sugi K, Kiura K, Date H, Shimizu N (2006) Detection of EGFR gene mutation in lung cancer by mutant-enriched polymerase chain reaction assay. *Clin Cancer Res* 12: 43–48
- Bosari S, Marchetti A, Buttitta F, Graziani D, Borsani G, Loda M, Bevilacqua G, Coggi G (1995) Detection of p53 mutations by single-strand conformation polymorphisms (SSCP) gel electrophoresis. A comparative study of radioactive and nonradioactive silver-stained SSCP analysis. *Diagn Mol Pathol* 4: 249–255
- Breathnach OS, Freidlin B, Conley B, Green MR, Johnson DH, Gandara DR, O'Connell M, Shepherd FA, Johnson BE (2001) Twenty-two years of phase III trials for patients with advanced non-small-cell lung cancer: sobering results. *J Clin Oncol* 19: 1734–1742
- Chernow B, Shahn SA (1997) Carcinomatous involvement of the pleura. *Am J Med* 63: 695–702
- Fan X, Furnari FB, Cavenee WK, Castresana JS (2001) Non-isotopic silver-stained SSCP is more sensitive than automated direct sequencing for the detection of PTEN mutations in a mixture of DNA extracted from normal and tumor cells. *Int J Oncol* 18: 1023–1026
- Fenton KN, David Richardson J (1995) Diagnosis and management of malignant pleural effusions. *Am J Surg* 170: 69–74
- Franklin WA, Vee R, Hirsch FR, Helfrich BA, Bunn Jr PA (2002) Epidermal growth factor receptor family in lung cancer and premalignancy. *Semin Oncol* 29: 3–14
- Fukuoka M, Yano S, Giaccone G, Tamura T, Nakagawa K, Douillard JY, Nishiaki Y, Vansteenkiste J, Kudoh S, Rischin D, Eek R, Horai T, Noda K, Takata I, Smit E, Averbuch S, Macleod A, Feyereislova A, Dong RP, Baselga J (2003) Multi-institutional randomized phase II trial of gefitinib for previously treated patients with advanced non-small-cell lung cancer (The IDEAL 1 Trial) [corrected]. *J Clin Oncol* 21: 2237–2246
- Giaccone G, Herbst RS, Manegold C, Scagliotti G, Rosell R, Miller V, Natale RB, Schiller JH, Von Pawel J, Pluzanska A, Gatzemeier U, Grous J, Ochs JS, Averbuch SD, Wolf MK, Rennie P, Fandi A, Johnson DH (2004) Gefitinib in combination with gemcitabine and cisplatin in advanced non-small-cell lung cancer: a phase III trial – INTACT 1. *J Clin Oncol* 22: 777–784
- Han SW, Kim TY, Hwang PG, Jeong S, Kim J, Choi IS, Oh DY, Kim JH, Kim DW, Chung DH, Im SA, Kim YT, Lee JS, Heo DS, Bang YJ, Kim NK (2005) Predictive and prognostic impact of epidermal growth factor receptor mutation in non-small-cell lung cancer patients treated with gefitinib. *J Clin Oncol* 23: 2493–2501
- Herbst RS, Giaccone G, Schiller JH, Natale RB, Miller V, Manegold C, Scagliotti G, Rosell R, Oliff I, Reeves JA, Wolf MK, Krebs AD, Averbuch SD, Ochs JS, Grous J, Fandi A, Johnson DH (2004) Gefitinib in combination with paclitaxel and carboplatin in advanced non-small-cell lung cancer: a phase III trial – INTACT 2. *J Clin Oncol* 22: 785–794
- Huang MJ, Lim KH, Tzen CY, Hsu HS, Yen Y, Huang BS (2005) EGFR mutations in malignant pleural effusion of non-small cell lung cancer: a case report. *Lung Cancer* 49: 413–415
- Kelly K, Crowley J, Bunn Jr PA, Presant CA, Grevstad PK, Moinpour CM, Ramsey SD, Wozniak AJ, Weiss GR, Moore DF, Israel VK, Livingston RB, Gandara DR (2001) Randomized phase III trial of paclitaxel plus carboplatin versus vinorelbine plus cisplatin in the treatment of patients with advanced non-small-cell lung cancer: a Southwest Oncology Group trial. *J Clin Oncol* 19: 3210–3218
- Kosaka T, Yatabe Y, Endoh H, Kuwano H, Takahashi T, Mitsudomi T (2004) Mutations of the epidermal growth factor receptor gene in lung cancer: biological and clinical implications. *Cancer Res* 64: 8919–8923
- Lee JH, Hong YS, Ryu JS, Chang JH (2004) p53 and FHIT mutations and microsatellite alterations in malignancy-associated pleural effusion. *Lung Cancer* 44: 33–42
- Lynch TJ, Bell DW, Sordella R, Gurubhagavatula S, Okimoto RA, Brannigan BW, Harris PL, Haserlat SM, Supko JG, Haluska FG, Louis DN, Christiani DC, Settleman J, Haber DA (2004) Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *N Engl J Med* 350: 2129–2139
- Marchetti A, Martella C, Felicioni L, Barassi F, Salvatore S, Chella A, Camplese PP, Iarussi T, Mucilli F, Mezzetti A, Cuccurullo F, Sacco R, Buttitta F (2005) EGFR mutations in non-small-cell lung cancer: analysis of a large series of cases and development of a rapid and sensitive method for diagnostic screening with potential implications on pharmacologic treatment. *J Clin Oncol* 23: 857–865
- Nakamoto M, Teramoto H, Matsumoto S, Igishi T, Shimizu E (2001) K-ras and rho A mutations in malignant pleural effusion. *Int J Oncol* 19: 971–976
- Paez JG, Janne PA, Lee JC, Tracy S, Greulich H, Gabriel S, Herman P, Kaye FJ, Lindeman N, Boggon TJ, Naoki K, Sasaki H, Fujii Y, Eck MJ, Sellers WR, Johnson BE, Meyerson M (2004) EGFR mutations in lung cancer: correlation with clinical response to gefitinib therapy. *Science* 304: 1497–1500
- Pao W, Miller V, Zakowski M, Doherty J, Politi K, Sarkaria I, Singh B, Heelan R, Rusch V, Fulton L, Mardis E, Kupfer D, Wilson R, Kris M, Varmus H (2004) EGF receptor gene mutations are common in lung cancers from “never smokers” and are associated with sensitivity of tumors to gefitinib and erlotinib. *Proc Natl Acad Sci USA* 101: 13306–13311
- Parkin DM, Bray F, Ferlay J, Pisani P (2005) Global cancer statistics, 2002. *CA Cancer J Clin* 55: 74–108
- Pass HI, Johnson DH, Minna JD, Turrissi AT (2005) Lung cancer principles and practice. In *Clinical Presentation of Non-Small Cell Carcinoma of the Lung* Antoinette J, Wozniak, SMG (eds) pp 291–303. Philadelphia: Lippincott Williams & Wilkins
- Sasaki H, Endo K, Konishi A, Takada M, Kawahara M, Iuchi K, Matsumura A, Okumura M, Tanaka H, Kawaguchi T, Shimizu T, Takeuchi H, Yano M, Fukai I, Fujii Y (2005) EGFR Mutation status in Japanese lung cancer patients: genotyping analysis using Light Cycler. *Clin Cancer Res* 11: 2924–2929
- Schiller JH, Harrington D, Belani CP, Langer C, Sandler A, Krook J, Zhu J, Johnson DH, the Eastern Cooperative Oncology Group (2002) Comparison of four chemotherapy regimens for advanced non-small-cell lung cancer. *N Engl J Med* 346: 92–98
- Takano T, Ohe Y, Sakamoto H, Tsuta K, Matsuno Y, Tateishi U, Yamamoto S, Nokihara H, Yamamoto N, Sekine I, Kunitoh H, Shibata T, Sakiyama T, Yoshida T, Tamura T (2005) Epidermal growth factor receptor gene mutations and increased copy numbers predict gefitinib sensitivity in patients with recurrent non-small-cell lung cancer. *J Clin Oncol* 23: 6829–6837
- Therasse P, Arbuck SG, Eisenhauer EA, Wanders J, Kaplan RS, Rubinstein L, Verweij J, Van Glabbeke M, van Oosterom AT, Christian MC, Gwyther SG (2000) New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada. *J Natl Cancer Inst* 92: 205–216
- Travis W, Colbey TJ, Corrin B (1999) *Histologic Typing of Tumors of Lung and Pleura: World Health Organization International Classification of Tumors* 3rd edn New York, NY: Springer Verlag
- Wookey A, Ellison G, Donald E (2005) Comparison of methods for the detection of mutations in the epidermal growth factor receptor gene. *In 96th Ann Meet Am Assoc Cancer Res* 46: 1250, abstr. 5287